

**CHLOPHEDIANOL HYDROCHLORIDE, DEXCHLORPHENIRAMINE MALEATE, AND PSEUDOEPHEDRINE HYDROCHLORIDE- chlophedianol hydrochloride, dexchlorpheniramine maleate, and pseudoephedrine hydrochloride liquid**  
**Westminster Pharmaceuticals, LLC**

-----  
**chlophedianol hydrochloride, dexchlorpheniramine maleate, and pseudoephedrine hydrochloride**

***Drug Facts***

<b><i>Active ingredients (in each 5 mL teaspoonful)</i></b>		<b><i>Purpose</i></b>
Chlophedianol Hydrochloride 12.5 mg		Cough Suppressant
Dexchlorpheniramine Maleate 1 mg		Antihistamine
Pseudoephedrine Hydrochloride 30 mg		Nasal Decongestant

**Uses**

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces congestion
- reduces swelling of nasal passages

**Warnings**

**Do not exceed recommended dosage.**

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus)

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland
- heart disease
- high blood pressure
- thyroid disease
- diabetes

**Ask a doctor before use if you are** taking sedatives or tranquilizers.

**When using this product**

- excitability may occur, especially in children
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.
- new symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

**Do not exceed recommended dosage.**

adults and children 12 years of age and over:	2 teaspoonfuls every 6 hours, not to exceed 8 teaspoonfuls in 24 hours.
children 6 to under 12 years of age:	1 teaspoonful every 6 hours, not to exceed 4 teaspoonfuls in 24 hours.
children under 6 years of age:	consult a doctor

**Other Information**

- Store at room temperature 68°-86°F (20°-30°C) with excursions of 59°-86°F (15°-30°C)

**Inactive ingredients**

citric acid anhydrous, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol solution, sucralose.

**Questions or Comments?**

Call weekdays from 9 AM to 5 PM EST at 1-844-7294. You may also report serious side effects to this phone number.

**PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label**

NDC 69367-183-16

Chlophedianol HCl,  
Dexchlorpheniramine  
Maleate and  
Pseudoephedrine HCl

Cough Suppressant  
Antihistamine  
Nasal Decongestant

Sugar Free, Alcohol Free,  
Dye Free

Tutti Frutti Flavor

EACH 5 mL (1 TEASPOONFUL) CONTAINS:

Chlophedianol Hydrochloride 12.5 mg  
Dexchlorpheniramine Maleate 1 mg  
Pseudoephedrine Hydrochloride 30 mg

16 fl. oz. (473 mL)

Westminster  
Pharmaceuticals

Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing.

**THIS BOTTLE IS NOT TO BE DISPENSED TO CONSUMER.**

Dispense in a tight, light-resistant container with a child resistant cap.

**Manufactured for:**

Westminster Pharmaceuticals LLC.

Nashville TN, 37217

Rev. 05/21

US Patent # 9,050,289

NDC 69367-183-16

**Chlophedianol HCl,  
Dexchlorpheniramine  
Maleate and  
Pseudoephedrine HCl**

**Cough Suppressant  
Antihistamine  
Nasal Decongestant**

**Sugar Free, Alcohol Free,  
Dye Free**

**Tutti Frutti Flavor**

**EACH 5 mL (1 TEASPOONFUL) CONTAINS:**  
Chlophedianol Hydrochloride.....12.5 mg  
Dexchlorpheniramine Maleate.....1 mg  
Pseudoephedrine Hydrochloride.....30 mg



16 fl. oz. (473 mL)

**WP Westminster  
Pharmaceuticals**

**Drug Facts** Lift Here

**Active ingredients Purpose  
(in each 5 mL teaspoonful)**

Chlophedianol  
Hydrochloride 12.5 mg.....Cough Suppressant  
Dexchlorpheniramine  
Maleate 1 mg.....Antihistamine  
Pseudoephedrine  
Hydrochloride 30 mg.....Nasal Decongestant

**Uses**

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces congestion
- reduces swelling of nasal passages

**Warnings**

**Do not exceed recommended dosage.**

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Drug Facts (continued)**

**Ask a doctor before use if you have**

- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland
- heart disease
- high blood pressure
- thyroid disease
- diabetes

**Ask a doctor before use if you are taking sedatives or tranquilizers.**

**When using this product**

- excitability may occur, especially in children
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.
- new symptoms occur

**If pregnant or breast-feeding, ask a health professional before use.**

**Drug Facts (continued)**

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

**Do not exceed recommended dosage.**

adults and children 12 years of age and over:	2 teaspoonfuls every 6 hours, not to exceed 8 teaspoonfuls in 24 hours.
children 6 to under 12 years of age:	1 teaspoonful every 6 hours, not to exceed 4 teaspoonfuls in 24 hours.
children under 6 years of age:	consult a doctor

**Other Information**

- Store at room temperature 68°-86°F (20°-30°C) with excursions of 59°-86°F (15°-30°C)

**Inactive ingredients**

citric acid anhydrous, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol solution, sucralose.

**Questions or Comments?**

Call weekdays from 9 AM to 5 PM EST at 1-844-7294. You may also report serious side effects to this phone number.

**Manufactured for:**

Westminster Pharmaceuticals LLC.

Nashville TN, 37217

Rev. 05/21

CHLOPHEDIANOL HYDROCHLORIDE, DEXCHLORPHENIRAMINE  
MALEATE, AND PSEUDOEPHEDRINE HYDROCHLORIDE

chlophedianol hydrochloride, dexchlorpheniramine maleate, and pseudoephedrine hydrochloride  
liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69367-183
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
CHLOPHEDIANOL HYDROCHLORIDE (UNII: 69QQ58998Y) (CHLOPHEDIANOL - UNII:42C50P12AP)	CHLOPHEDIANOL HYDROCHLORIDE	12.5 mg in 5 mL
DEXCHLORPHENIRAMINE MALEATE (UNII: B10YD955QW) (DEXCHLORPHENIRAMINE - UNII:3Q9Q0B929N)	DEXCHLORPHENIRAMINE MALEATE	1 mg in 5 mL
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69367-183-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/29/2021	10/01/2024

Marketing Information			
Marketing	Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
OTC MONOGRAPH DRUG	M012	07/29/2021	10/01/2024

**Labeler -** Westminster Pharmaceuticals, LLC (079516651)

Revised: 7/2021

Westminster Pharmaceuticals, LLC