

CHILDRENS PAIN AND FEVER- acetaminophen suspension
L.N.K. International, Inc.

Quality Plus 44-079

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - sore throat
 - flu
 - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this

product

Ask a doctor before use if your child has

liver disease.

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- **shake well before using**
- mL = milliliter; FL OZ = fluid ounce
- only use the dose cup provided
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
Under 24	Under 2	Ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 mL
48-59	6-8	10 mL
60-71	9-10	12.5 mL
72-95	11	15 mL

*or as directed by a doctor

Other information

- **each 5 mL contains:** sodium 3 mg

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

anhydrous citric acid, FD&C red #40, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions or comments?

1-800-426-9391

Principal display panel

**QUALITY
+PLUS**

NDC 50844-079-36

Compare to active ingredient
in Children's Tylenol®*

**CHILDREN'S
PAIN &
FEVER**

**Acetaminophen
160 mg per 5 mL**

Oral Suspension
Pain Reliever/
Fever Reducer

- Ages 2-11
- Pain/Fever
- Ibuprofen Free
- Aspirin Free

Strawberry
Flavor

ALCOHOL FREE • NO PARABENS

4 FL OZ (118 mL)

**IMPORTANT:
Please see dosing chart or ask a
doctor for your child's correct dose**

DIRECTIONS:

STEP 1.

**Shake well
before using**

STEP 2.
Remove child
protective
cap

STEP 3.
Pour your
child's dose
into the
dosing cup

TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR
MISSING

*This product is not manufactured or distributed by
Johnson & Johnson Corporation, owner of the
registered trademark Children's Tylenol®.
50844 ORG092207936

Distributed by
LNK INTERNATIONAL, INC.
60 Arkay Drive
Hauppauge, NY 11788
USA

QUALITY
+PLUS

CHILDREN'S

PAIN & FEVER

Ages 2-11

4 FL OZ (118 mL)

QUALITY
+PLUS

NDC 50844-079-36

Compare to active ingredient
in Children's Tylenol®

CHILDREN'S

PAIN & FEVER

Acetaminophen
160 mg per 5 mL
Oral Suspension
Pain Reliever/
Fever Reducer



Strawberry
Flavor



- Ages 2-11
- Pain/Fever
- Ibuprofen Free
- Aspirin Free

ALCOHOL FREE • NO PARABENS

4 FL OZ (118 mL)

DIRECTIONS:

STEP 1.
Shake well
before using



STEP 2.
Remove child
protective
cap



STEP 3.
Pour your
child's dose
into the
dosing cup



IMPORTANT:
Please see dosing chart or ask a
doctor for your child's correct dose

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR
MISSING**

Drug Facts

KEEP OUTER PACKAGE FOR
COMPLETE PRODUCT INFORMATION

Active ingredient (in each 5 mL) Purpose
Acetaminophen 160 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - sore throat
 - flu
 - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount

with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

this product does not contain directions or complete warnings for adult use

do not give more than directed

shake well before using

mL = milliliter; FL OZ = fluid ounce

only use the dose cup provided

find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

repeat dose every 4 hours while symptoms last

do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
Under 24	Under 2	Ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 mL
48-59	6-8	10 mL
60-71	9-10	12.5 mL
72-95	11	15 mL

*or as directed by a doctor

Other information

- each 5 mL contains: sodium 3 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients anhydrous citric acid, FD&C red #40, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions or comments? 1-800-426-9391

This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Children's Tylenol®. 50844 ORG092207936

Distributed by
LNK INTERNATIONAL, INC.
60 Arkay Drive
Hauppauge, NY 11788
USA

B-1603-079-36
ORG092207936



No Print / No Varnish
Lot & Expiry Area

Quality Plus 44-079

CHILDRENS PAIN AND FEVER

acetaminophen suspension

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:50844-079

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	STRAWBERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-079-36	1 in 1 CARTON	02/01/2023	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/01/2023	

Labeler - L.N.K. International, Inc. (038154464)**Establishment**

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Revised: 2/2024

L.N.K. International, Inc.