

LIVE BETTER TRIPLE ANTIBIOTIC PLUS PAIN RELIEF- bacitracin zinc, neomycin sulfate, polymyxin b sulfate, and pramoxine hydrochloride ointment

The Great Atlantic & Pacific Tea Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Live Better

Triple Antibiotic plus Pain Relief

Drug Facts

Active ingredients (each gram contains)	Purpose
Bacitracin zinc 500 units	First aid antibiotic
Neomycin sulfate 3.5 mg	First aid antibiotic
Polymyxin B sulfate 10,000 units	First aid antibiotic
Pramoxine hydrochloride 10 mg	External analgesic

Uses

first aid to help prevent infection and for the temporary relief of pain or discomfort in minor:

- cuts
- scrapes
- burns

Warnings

For external use only.

Do not use

- in the eyes
- over large areas of the body
- if you are allergic to any of the ingredients

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- condition persists or gets worse
- you need to use longer than 1 week
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- a rash or other allergic reaction develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Adults and children 2 years of age and older:
 - clean the affected area and dry thoroughly

- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage
- Children under 2 years of age: ask a doctor

Other information

- To open: unscrew cap, pull tab to remove foil seal
- store at 20° to 25°C (68° to 77°F)
- see carton or tube crimp for lot number and expiration date

Inactive ingredient

white petrolatum

Questions?

Call **1-866-923-4914**

DISTRIBUTED BY ONPOINT, INC.
2 PARAGON DRIVE,
MONTVALE, NJ 07645

PRINCIPAL DISPLAY PANEL - 28.4 g Tube Carton**Maximum Strength****Triple Antibiotic Ointment + Pain Relief**

■ BACITRACIN ZINC ■ NEOMYCIN SULFATE ■ POLYMYXIN B SULFATE ■
PRAMOXINE HYDROCHLORIDE

First Aid Antibiotic / Pain Relieving Ointment

NET WT 1 OZ (28.4g)

NDC 51143-027-02

Compare to the active ingredients in Neosporin® + Pain Relief™



Helps Prevent Infection in Minor Cuts, Scrapes, and Burns Plus Maximum Strength Pain Relief.

LPK-7203-0
0113-0
M268

Maximum Strength

Triple Antibiotic Ointment + Pain Relief

■ BACITRACIN ZINC ■ NEOMYCIN SULFATE ■ POLYMYXIN B SULFATE ■ PRAMOXINE HYDROCHLORIDE

First Aid Antibiotic / Pain Relieving Ointment

NET WT 1 OZ (28.4g)

T51



Drug Facts (continued)

Other information

- To open: unscrew cap, pull tab to remove foil seal
- store at 20° to 25°C (68° to 77°)
- see carton or tube crimp for lot number and expiration date

Inactive ingredient white petrolatum

Questions? Call 1-866-923-4014

NO COPY / NO COLOR
THIS FLAP FOR LOT #
AND EXP DATE PRINT



DISTRIBUTED BY ONPOINT, INC.
2 PARAGON DRIVE,
MONTVALE, NJ 07645
© 2013. ALL RIGHTS RESERVED.
Made in Canada

The product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Neosporin® + Pain Relief

Drug Facts (each gram contains)

Bacitracin zinc 500 units.
Neomycin sulfate 3.5 mg.
Polymyxin B sulfate 10,000 units.
Pramoxine hydrochloride 10 mg.

Purpose
First aid antibiotic
First aid antibiotic
External analgesic

Uses
first aid to help prevent infection and for the temporary relief of pain or discomfort in minor:
cuts • scrapes • burns

Warnings
For external use only.
Do not use • in the eyes • over large areas of the body
• if you are allergic to any of the ingredients
Ask a doctor before use if you have
deep or puncture wounds • animal bites • serious burns
Stop use and ask a doctor if
• condition persists or gets worse
• you need to use longer than 1 week
• symptoms persist for more than 1 week, or clear up and occur again within a few days
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
• Adults and children 2 years of age and older:
• clean the affected area and dry thoroughly.
• apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily.
• may be covered with a sterile bandage
• Children under 2 years of age: ask a doctor.

LIVE BETTER TRIPLE ANTIBIOTIC PLUS PAIN RELIEF
bacitracin zinc, neomycin sulfate, polymyxin b sulfate, and pramoxine hydrochloride ointment

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51143-027	
Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
Bacitracin Zinc (UNII: 89Y4M234ES) (Bacitracin - UNII:58H6RWO52I)		Bacitracin	500 [iU] in 1 g	
Neomycin Sulfate (UNII: 057Y626693) (Neomycin - UNII:I16QD7X297)		Neomycin	3.5 mg in 1 g	
Polymyxin B Sulfate (UNII: 19371312D4) (Polymyxin B - UNII:J2VZ07J96K)		Polymyxin B	10000 [iU] in 1 g	
Pramoxine hydrochloride (UNII: 88AYB867L5) (Pramoxine - UNII:068X84E056)		Pramoxine hydrochloride	10 mg in 1 g	
Inactive Ingredients				
Ingredient Name			Strength	
petrolatum (UNII: 4T6H12BN9U)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51143-027-02	1 in 1 CARTON		
1		28.4 g in 1 TUBE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part333B	03/31/2012		

Labeler - The Great Atlantic & Pacific Tea Company (001367366)

Registrant - Taro Pharmaceuticals U.S.A., Inc. (145186370)

Establishment

Name	Address	ID/FEI	Business Operations
Taro Pharmaceuticals Inc.		206263295	MANUFACTURE(51143-027)

Revised: 3/2013

The Great Atlantic & Pacific Tea Company