

**LEVOCETIRIZINE DIHYDROCHLORIDE- levocetirizine dihydrochloride tablet, coated**

**Target Corporation**

-----

**Drug Facts**

**Active ingredient (in each tablet)**

Levocetirizine dihydrochloride USP, 5 mg

**Purpose**

Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use**

- if you have kidney disease
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

**Ask a doctor before use if you have**

- ever had trouble urinating or emptying your bladder

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask doctor if**

- you have trouble urinating or emptying your bladder
- an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended

- if pregnant: ask a health professional before use

### Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

adults 65 years of age and older	<ul style="list-style-type: none"> <li>• ask a doctor</li> </ul>
adults and children 12-64 years of age	<ul style="list-style-type: none"> <li>• take 1 tablet (5 mg) once daily in the evening</li> <li>• do not take more than 1 tablet (5 mg) in 24 hours</li> <li>• 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms</li> </ul>
children 6-11 years of age	<ul style="list-style-type: none"> <li>• take 1/2 tablet (2.5 mg) once daily in the evening</li> <li>• do not take more than 1/2 tablet (2.5 mg) in 24 hours</li> </ul>
children under 6 years of age	<ul style="list-style-type: none"> <li>• do not use</li> </ul>
consumers with kidney disease	<ul style="list-style-type: none"> <li>• do not use</li> </ul>

### Other information

- store between 20° and 25°C (68° and 77°F)
- (Bottle only)- Safety Sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing
- (Blister only)- Safety Sealed: Do not use if seal is broken or if individual blister unit is open or torn

### Inactive ingredients

colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

### Questions or comments?

Call **1-888-375-3784**

### Carton Label

Carton Label



7 00101051

**Drug Facts**

**Active ingredient (in each tablet)**  
Levocetirizine dihydrochloride USP, 5 mg — Antihistamine

**Uses**  
temporarily relieves these symptoms due to hay fever or other respiratory allergies:  
■ runny nose  
■ sneezing  
■ itchy, watery eyes  
■ itching of the nose or throat

**Warnings**  
Do not use  
■ if you have kidney disease  
■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

**Ask a doctor before use if you have**  
■ ever had trouble urinating or emptying your bladder

**When using this product**  
■ drowsiness may occur  
■ avoid alcoholic drinks  
■ alcohol, sedatives, and tranquilizers may increase drowsiness  
■ use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**  
■ you have trouble urinating or emptying your bladder  
■ an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding:**  
■ if breast-feeding, not recommended  
■ if pregnant, ask a health professional before use  
Keep out of reach of children.  
In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Drug Facts (continued)**

**Directions**  
adults 65 years of age and older

■ ask a doctor  
■ take 1 tablet (5 mg) once daily in the evening  
■ do not take more than 1 tablet (5 mg) in 24 hours  
■ 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms

children 6-11 years of age  
■ take 1/2 tablet (2.5 mg) once daily in the evening  
■ do not take more than 1/2 tablet (2.5 mg) in 24 hours

children under 6 years of age  
■ do not use

consumers with kidney disease  
■ do not use

**Other information**

■ store between 20° and 25°C (68° and 77°F)  
■ safety sealed; do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

**Inactive ingredients**

croscollon, silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

**Questions or comments?**  
Call 1-888-375-3784

Read directions and warnings before use. Keep this carton, it has important information.



150101004



Original Prescription Strength  
**Allergy Relief**  
Levocetirizine Dihydrochloride  
Tablets USP, 5 mg  
Antihistamine

NDC 11673-447-35

Compare to active ingredient in Xyzal® Allergy\*

Original Prescription Strength

**Allergy Relief**  
Levocetirizine Dihydrochloride  
Tablets USP, 5 mg  
Antihistamine

Original Prescription Strength

**Allergy Relief**  
Levocetirizine Dihydrochloride  
Tablets USP, 5 mg  
Antihistamine

24-hour relief of sneezing, runny nose, itchy nose or throat, itchy and watery eyes

- Clinically Proven Relief in 60 Minutes
- Clinically Proven Relief For 24 Hours



35 TABLETS



35 Tablets Actual Size

Satisfaction guaranteed – Love it or your money back.

Distributed by Target Corporation  
Minneapolis, MN 55403  
Made in India  
TM & ©2024 Target Brands, Inc.  
\*This product is not manufactured or distributed by Chatham, Inc., (part of the Sanofi Group), distributor of Xyzal® Allergy 24HR Tablets. Xyzal® is a registered trademark of UCB Biopharma SRL.

094 04 1984 R01 C-002262-01-009-0000



8 48985 00057 4

Issued: 02/25

LOT

EXP

Bottle Label

Bottle label

NDC 11673-847-35  
**Original Prescription Strength**  
**Allergy Relief**  
 Levocetirizine Dihydrochloride  
 Tablets USP, 5 mg  
 Antihistamine



35 TABLETS



**Active ingredient (in each tablet) Purpose**

Levocetirizine dihydrochloride USP, 5 mg.....Antihistamine

**Uses** temporarily relieves these symptoms due to hay fever or other respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

**Warnings Do not use** ■ if you have kidney disease ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine **Ask a doctor before use if you have** ■ ever had trouble urinating or emptying your bladder

**When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery **Stop use and ask a doctor if** ■ you have trouble urinating or

(Continued On Back Of Label)

094 04 3964 R00 C-002262-01-009

Distribution by Target Corporation  
 Minneapolis, MN 55403

Made in India

TM & ©2024

Target Brands, Inc.

REV 08/23

150094155



emptying your bladder ■ an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding:** ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** adults 65 years of age and older ■ ask a doctor adults and children 12-64 years of age ■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours ■ 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms children 6-11 years of age ■ take 1/2 tablet (2.5 mg) once daily in

the evening ■ do not take more than 1/2 tablet (2.5 mg) in 24 hours children under 6 years of age ■ do not use consumers with kidney disease ■ do not use

**Other information** ■ store between 20° and 25°C (68° and 77°F) ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

**Inactive ingredients** colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

**Questions and comments?** call 1-888-375-3784

## LEVOCETIRIZINE DIHYDROCHLORIDE

levocetirizine dihydrochloride tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11673-847(NDC:43598-735)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
levocetirizine dihydrochloride (UNII: SOD6A38AGA) (levocetirizine - UNII:6U5EA9RT2O)	levocetirizine dihydrochloride	5 mg

### Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	L
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-847-80	1 in 1 CARTON	12/31/2018	
1		80 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11673-847-16	2 in 1 CARTON	12/01/2020	
2		80 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:11673-847-35	1 in 1 CARTON	12/31/2018	
3		35 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:11673-847-79	2 in 1 BLISTER PACK	12/31/2018	12/31/2018
4		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210375	12/31/2018	

**Labeler** - Target Corporation (006961700)