

**SALICYLIC ACID- medicated corn removers patch**  
**Topco Associates LLC**

-----  
**TopCare Medicated Corn Removers**

***Active ingredient***

Salicylic acid 40%

***Purpose***

Corn Remover

***Uses***

- for the removal of corns
- relieves pain by removing corns

***Warnings***

**For external use only.**

**Do not use**

- if you are diabetic
- have poor blood circulation
- on irritated skin, or any area that is infected or reddened

**Stop Use and Ask a doctor**

if discomfort persists

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

***Directions***

- wash affected area and dry area thoroughly
- if necessary, cut medicated patch to fit corn
- apply adhesive side down of medicated patch onto corn
- cover medicated patch with pad
- after 48 hours, remove medicated patch
- repeat procedure every 48 hours as needed for up to 14 days (until corn is removed)
- may soak corn in warm water for 5 minutes to assist in removal

***Other information***

store between 15°C to 30°C (59°F to 86°F)

***Inactive ingredients***

acrylic adhesive, acrylic polymer, polyethylene, polyvinyl alcohol

***Questions?***

Call 1-888-423-0139

**Principal Display Panel**

TopCare  
Medicated

Corn

Removers

SALICYLIC ACID 40%

- Effective Corn Removal Treatment
- Cushions Against Pressure & Friction

9 PADS | 9 MEDICATED PATCHES

**Drug Facts**

| Active ingredient       | Purpose      |
|-------------------------|--------------|
| Salicylic acid 40%..... | Corn remover |

**Uses**

- for the removal of corns
- relieves pain by removing corns

**Warnings**  
For external use only.

**Do not use**

- if you are diabetic
- if you have poor blood circulation
- on irritated skin, on any area that is infected or reddened

**Stop use and ask a doctor if** discomfort persists

**Keep out of reach of children.**  
If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- wash affected area and dry thoroughly
- if necessary, cut medicated patch to fit corn
- apply adhesive side of medicated patch onto corn
- cover medicated patch with pad
- after 48 hours, remove medicated patch
- repeat procedure every 48 hours as needed for up to 14 days (until corn is removed)
- may soak corn in warm water for 5 minutes to assist in removal

**Other information**  
store between 15° and 30°C (59° and 86°F)

**Inactive ingredients**  
acrylic adhesive, acrylic polymer, polyethylene, polyvinyl alcohol

**Questions?** Call 1-888-423-0139



COMPARE TO DR. SCHOLL'S® ACTIVE INGREDIENT\*

# Medicated Corn Removers


## SALICYLIC ACID 40%

- Effective Corn Removal Treatment
- Cushions Against Pressure & Friction



**9 PADS | 9 MEDICATED PATCHES**

0 36800 05452 3



3

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Dr. Scholl's®. DISAPPROVED BY TOPCO ASSOCIATES LLC. ELKS GROVEVILLE, IL 60037 © TOPCO PHARM 0918 QUESTIONS? 1-888-423-0139 www.topcarehealth.com topcaretopco.com

MADE IN USA WITH U.S. AND IMPORTED PARTS

The TopCare® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.

Scan here for more information



52-003TC-01

## SALICYLIC ACID

medicated corn removers patch

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:36800-021 |
| <b>Route of Administration</b> | TOPICAL        |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength | Strength   |
|---|-------------------|------------|
| <b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ) | SALICYLIC ACID    | 40 mg in 9 |

**Inactive Ingredients**

| Ingredient Name                                     | Strength |
|---|----------|
| <b>POLYVINYL ALCOHOL</b> (UNII: 532B59J990)         |          |
| <b>HIGH DENSITY POLYETHYLENE</b> (UNII: UG00KM4WR7) |          |

**Packaging**

| # | Item Code        | Package Description                               | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:36800-021-09 | 9 in 1 PACKAGE; Type 0: Not a Combination Product | 01/03/2012           |                    |

**Marketing Information**

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M030                                     | 01/03/2012           |                    |

**Labeler** - Topco Associates LLC (006935977)

Revised: 1/2026

Topco Associates LLC