MAXIMUM STRENGTH ANTACID- aluminum hydroxide, magnesium hydroxide, simethicone suspension CARDINAL HEALTH

Idr antacid max cherry

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide (equivalent to dried gel, USP) 400 mg Magnesium hydroxide 400 mg Simethicone 40 mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug.

Antacids may interact with certain prescription drugs.

Stop use and ask a doctor ifsymptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years of age and older: take 2 to 4 teaspoonfuls between meals, at bedtime or as directed by a doctor

- do not exceed 8 teaspoonfuls in a 24 hour period or use the maximum dosage for more than 2 weeks
- children under 12 years of age: ask a doctor

Other information

- each 5 mL teaspoonful contains: magnesium 165 mg, sodium 5 mg
- store at room temperature
- protect from freezing
- keep tightly closed


Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or Comments?

1-800-540-3765

package label



Drug Facts

TAMPER-EVIDENT: Do not use if breakaway band on bottle cap is missing or broken.

Active ingredients Purpose (in each 5 mL teaspoonful)

Uses relieves • heartburn • sour stomach • acid indigestion • the symptoms referred to as gas

Warnings

Simethicone 40 mg ..

Ask a doctor before use if you have

kidney disease
 magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions • shake well before use

- adults and children 12 years of age and older: take 2 to 4 teaspoonfuls between meals, at bedtime or as directed by a doctor
- do not take more than 8 teaspoonfuls in a 24 hour period or use the maximum dosage for more than 2 weeks
- children under 12 years of age: ask a doctor

Other information • each 5 mL teaspoonful contains: magnesium 165 mg, sodium 5 mg • store at room temperature

- protect from freezing
- keep tightly closed

Inactive ingredients benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

DISTRIBUTED BY CARDINAL HEALTH Dublin, Ohio 43017

© 2018 Cardinal Health, All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health.

995-11322-0 REV 0818



MAXIMUM STRENGTH ANTACID

aluminum hydroxide, magnesium hydroxide, simethicone suspension

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:70000-0422

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 5 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y10) (DIMETHICONE - UNII:92RU3N3Y10)	DIMETHICONE	40 mg in 5 mL

Inactive Ingredients				
Ingredient Name	Strength			
BENZYL ALCOHOL (UNII: LKG8494WBH)				
BUTYLPARABEN (UNII: 3QPI1U3FV8)				
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K6790BS311)				
HYPROMELLOSES (UNII: 3NXW29V3WO)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
PROPYLPARABEN (UNII: Z8IX2SC10H)				
WATER (UNII: 059QF0KO0R)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
SORBITOL SOLUTION (UNII: 8KW3E207O2)				

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

l	Packaging				
	#	# Item Code Package Description		Marketing Start Date	Marketing End Date
	1	NDC:70000- 0422-1	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2019	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	01/01/2019	

Labeler - CARDINAL HEALTH (063997360)

Registrant - GCP Laboratories (965480861)

Establishment				
Name	Address	ID/FEI	Business Operations	
GCP Laboratories		965480861	manufacture(70000-0422)	

Revised: 11/2023 CARDINAL HEALTH