

# HEPARIN SODIUM- heparin sodium injection, solution

## McKesson Corporation dba SKY Packaging

### HIGHLIGHTS OF PRESCRIBING INFORMATION

#### Heparin Sodium Injection, USP

These highlights do not include all the information needed to use HEPARIN SODIUM INJECTION safely and effectively. See full prescribing information for HEPARIN SODIUM INJECTION.

### HEPARIN SODIUM injection, for intravenous or subcutaneous use

Initial U.S. Approval: 1939

#### INDICATIONS AND USAGE

Heparin sodium injection is an anticoagulant indicated for ( 1)

- Prophylaxis and treatment of venous thrombosis and pulmonary embolism
- Prevention of postoperative deep venous thrombosis and pulmonary embolism in patients undergoing major abdominothoracic surgery or who, for other reasons, are at risk of developing thromboembolic disease
- Atrial fibrillation with embolization
- Treatment of acute and chronic consumptive coagulopathies (disseminated intravascular coagulation)
- Prevention of clotting in arterial and cardiac surgery
- Prophylaxis and treatment of peripheral arterial embolism
- Use as an anticoagulant in blood transfusions, extracorporeal circulation, and dialysis procedures

#### DOSAGE AND ADMINISTRATION

Recommended Adult Dosages:

- Therapeutic Anticoagulant Effect with Full-Dose Heparin † ( 2.3)

Deep Subcutaneous (Intrafat) Injection  <i>Use a different site for each injection</i>	Initial dose	5,000 units by intravenous injection followed by 10,000 to 20,000 units of a concentrated solution, subcutaneously
	Every 8 hours	8,000 to 10,000 units of a concentrated solution
	or Every 12 hours	15,000 to 20,000 units of a concentrated solution
Intermittent Intravenous Injection	Initial dose	10,000 units, either undiluted or in 50 to 100 mL of 0.9% Sodium Chloride Injection, USP
	Every 4 to 6 hours	5,000 to 10,000 units, either undiluted or in 50 to 100 mL of 0.9% Sodium Chloride Injection, USP
Intravenous Infusion	Initial dose	5,000 units by intravenous injection
	Continuous	20,000 to 40,000 units/24 hours in 1,000 mL of 0.9% Sodium Chloride Injection, USP (or in any compatible solution) for infusion

† Based on 150 lb (68 kg) patient. Adjust dose based on laboratory monitoring.

#### DOSAGE FORMS AND STRENGTHS

Heparin Sodium Injection, USP (porcine), **preservative free:** (3)

- 2 mL single-dose vial contains 2,000 USP units

Heparin sodium injection, USP contains **benzyl alcohol:** (3)

- 10 mL multiple-dose vial contains 50,000 USP units

Heparin sodium injection, USP contains **parabens:** (3)

- 1 mL multiple-dose vial contains 1,000 USP units
- 10 mL multiple-dose vial contains 10,000 USP units
- 30 mL multiple-dose vial contains 30,000 USP units
- 1 mL multiple-dose vial contains 5,000 USP units
- 1 mL multiple-dose vial contains 10,000 USP units
- 5 mL multiple-dose vial contains 50,000 USP units
- 1 mL multiple-dose vial contains 20,000 USP units

#### ----- **CONTRAINDICATIONS** -----

- History of heparin-induced thrombocytopenia (HIT) or heparin-induced thrombocytopenia and thrombosis (HITTS) ( 4)
- Known hypersensitivity to heparin or pork products ( 4)
- In whom suitable blood coagulation tests cannot be performed at appropriate intervals ( 4)
- An uncontrolled bleeding state, except when this is due to disseminated intravascular coagulation ( 4)

#### ----- **WARNINGS AND PRECAUTIONS** -----

- Fatal Medication Errors: Confirm choice of correct strength prior to administration ( 5.1)
- Hemorrhage: Hemorrhage, including fatal events, has occurred in patients receiving heparin. Use caution in conditions with increased risk of hemorrhage ( 5.2)
- HIT and HITTS: Monitor for signs and symptoms and discontinue if indicative of HIT and HITTS ( 5.3)
- Benzyl Alcohol Toxicity: Use preservative-free formulation in neonates and infants
- Monitoring: Blood coagulation tests guide therapy for full-dose heparin. Periodically monitor platelet count, hematocrit, and occult blood in stool in all patients receiving heparin ( 5.5, 5.6)

#### ----- **ADVERSE REACTIONS** -----

Most common adverse reactions are hemorrhage, thrombocytopenia, HIT and HITTS, injection site irritation, general hypersensitivity reactions, and elevations of aminotransferase levels. ( 6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact McKesson Corporation dba SKY Packaging at 1-888-243-4363 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

#### ----- **DRUG INTERACTIONS** -----

Drugs that interfere with coagulation, platelet aggregation or drugs that counteract coagulation may induce bleeding ( 7.2)

#### ----- **USE IN SPECIFIC POPULATIONS** -----

- Pregnancy: Preservative-free formulation recommended. ( 8.1)
- Lactation: Preservative-free formulation recommended. ( 8.2)
- Pediatric Use: Use preservative-free formulation in neonates and infants. ( 8.4)

**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 1/2022**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

2.1 Preparation for Administration

2.2 Laboratory Monitoring for Efficacy and Safety

2.3 Therapeutic Anticoagulant Effect with Full-Dose Heparin

2.4 Pediatric Dosing

2.5 Cardiovascular Surgery

2.6 Low-Dose Prophylaxis of Postoperative Thromboembolism

2.7 Converting to Warfarin

2.8 Converting to Oral Anticoagulants other than Warfarin

2.9 Extracorporeal Dialysis

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

5.1 Fatal Medication Errors

5.2 Hemorrhage

5.3 Heparin-Induced Thrombocytopenia and Heparin-Induced Thrombocytopenia and Thrombosis

5.4 Risk of Serious Adverse Reactions in Infants Due to Benzyl Alcohol Preservative

5.5 Thrombocytopenia

5.6 Coagulation Testing and Monitoring

5.7 Heparin Resistance

5.8 Hypersensitivity

### **6 ADVERSE REACTIONS**

6.1 Postmarketing Experience

### **7 DRUG INTERACTIONS**

7.1 Oral Anticoagulants

7.2 Platelet Inhibitors

7.3 Other Interactions

### **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

### **10 OVERDOSAGE**

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

### **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Heparin sodium injection is indicated for:

- Prophylaxis and treatment of venous thrombosis and pulmonary embolism;
- Prevention of postoperative deep venous thrombosis and pulmonary embolism in

patients undergoing major abdominothoracic surgery or who, for other reasons, are at risk of developing thromboembolic disease;

- Atrial fibrillation with embolization;
- Treatment of acute and chronic consumptive coagulopathies (disseminated intravascular coagulation);
- Prevention of clotting in arterial and cardiac surgery;
- Prophylaxis and treatment of peripheral arterial embolism.
- Anticoagulant use in blood transfusions, extracorporeal circulation, and dialysis procedures.

## **2 DOSAGE AND ADMINISTRATION**

### **2.1 Preparation for Administration**

Confirm the choice of the correct heparin sodium injection vial to ensure that the 1 mL vial is not confused with a "catheter lock flush" vial or other 1 mL vial of incorrect strength [see *Warnings and Precautions (5.1)*]. Confirm the selection of the correct formulation and strength prior to administration of the drug.

Inspect parenteral drug products visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Use only if solution is clear and the seal is intact. Do not use if solution is discolored or contains a precipitate.

When heparin is added to an infusion solution for continuous intravenous (IV) administration, invert the container at least six times to ensure adequate mixing and prevent pooling of the heparin in the solution. Storage of prepared infusion solution should not exceed 4 hours at room temperature or 24 hours at 2° to 8°C (36° to 46°F).

Administer heparin sodium injection by intermittent intravenous injection, intravenous infusion, or deep subcutaneous (intrafat, i.e., above the iliac crest or abdominal fat layer) injection. The intramuscular route of administration should be avoided because of the frequent occurrence of hematoma at the injection site [see *Adverse Reactions (6.1)*].

### **2.2 Laboratory Monitoring for Efficacy and Safety**

Adjust the dosage of heparin sodium injection according to the patient's coagulation test results. Dosage is considered adequate when the activated partial thromboplastin time (aPTT) is 1.5 to 2 times normal or when the whole blood clotting time is elevated approximately 2.5 to 3 times the control value. When initiating treatment with heparin sodium injection by continuous intravenous infusion, determine the coagulation status (aPTT, INR, platelet count) at baseline and continue to follow aPTT approximately every 4 hours and then at appropriate intervals thereafter. When the drug is administered intermittently by intravenous injection, perform coagulation tests before each injection during the initiation of treatment and at appropriate intervals thereafter. After deep

subcutaneous (intrafat) injections, tests for adequacy of dosage are best performed on samples drawn 4 to 6 hours after the injection.

Periodically monitor platelet counts, hematocrit, and occult blood in stool during the entire course of heparin therapy, regardless of the route of administration.

### 2.3 Therapeutic Anticoagulant Effect with Full-Dose Heparin

The dosing recommendations in Table 1 are based on clinical experience. Although dosages must be adjusted for the individual patient according to the results of suitable laboratory tests, the following dosage schedules may be used as guidelines:

**Table 1: Recommended Adult Full-Dose Heparin Regimens for Therapeutic Anticoagulant Effect**

<b>Method of Administration</b>	<b>Frequency</b>	<b>Recommended Dose</b>
Deep Subcutaneous (Intrafat) Injection  Use a different site for each injection to prevent the development of hematoma	Initial dose	5,000 units by intravenous injection, followed by 10,000 to 20,000 units of a concentrated solution, subcutaneously
	Every 8 hours	8,000 to 10,000 units of a concentrated solution
	or Every 12 hours	15,000 to 20,000 units of a concentrated solution
Intermittent Intravenous Injection	Initial Dose	10,000 units, either undiluted or in 50 to 100 mL of 0.9% Sodium Chloride Injection, USP
	Every 4 to 6 hours	5,000 to 10,000 units, either undiluted or in 50 to 100 mL of 0.9% Sodium Chloride Injection, USP
Continuous Intravenous Infusion	Initial Dose	5,000 units by intravenous injection
	Continuous	20,000 to 40,000 units/24 hours in 1,000 mL of 0.9% Sodium Chloride Injection, USP (or in any compatible solution) for infusion

\*Based on 68 kg patient

### 2.4 Pediatric Dosing

Use preservative-free heparin sodium injection in neonates and infants [see *Warnings and Precautions (5.4)*].

There are no adequate and well controlled studies on heparin use in pediatric patients. Pediatric dosing recommendations are based on clinical experience. In general, the following dosage schedule may be used as a guideline in pediatric patients:

Recommended Pediatric Use	
Initial Dose	75 to 100 units/kg (IV bolus over 10 minutes)
Maintenance Dose	<p>Infants: 25 to 30 units/kg/hour;</p> <p>Infants &lt; 2 months have the highest requirements (average 28 units/kg/hour)</p> <p>Children &gt; 1 year of age: 18 to 20 units/kg/hour;</p> <p>Older children may require less heparin, similar to Weight-adjusted adult dosage</p>
Monitoring	Adjust heparin to maintain aPTT of 60 to 85 seconds, assuming this reflects an anti-Factor Xa level of 0.35 to 0.70

## 2.5 Cardiovascular Surgery

Patients undergoing total body perfusion for open-heart surgery should receive an initial dose of not less than 150 units of heparin sodium per kilogram of body weight. Frequently, a dose of 300 units per kilogram is used for procedures estimated to last less than 60 minutes, or 400 units per kilogram for those estimated to last longer than 60 minutes.

## 2.6 Low-Dose Prophylaxis of Postoperative Thromboembolism

The most widely used dosage has been 5,000 units 2 hours before surgery and 5,000 units every 8 to 12 hours thereafter for 7 days or until the patient is fully ambulatory, whichever is longer. Administer the heparin by deep subcutaneous (intrafat, i.e., above the iliac crest or abdominal fat layer, arm, or thigh) injection with a fine (25 to 26-gauge) needle to minimize tissue trauma.

## 2.7 Converting to Warfarin

To ensure continuous anticoagulation when converting from heparin sodium injection to warfarin, continue full heparin therapy for several days until the INR (prothrombin time) has reached a stable therapeutic range. Heparin therapy may then be discontinued without tapering [see *Drug Interactions (7.1)*].

## 2.8 Converting to Oral Anticoagulants other than Warfarin

For patients currently receiving intravenous heparin, stop intravenous infusion of heparin sodium immediately after administering the first dose of oral anticoagulant; or for intermittent intravenous administration of heparin sodium, start oral anticoagulant 0 to 2 hours before the time that the next dose of heparin was to have been administered.

## 2.9 Extracorporeal Dialysis

Follow equipment manufacturers' operating directions carefully. A dose of 25 to 30 units/kg followed by an infusion rate of 1,500 to 2,000 units/hour is suggested based on pharmacodynamic data if specific manufacturers' recommendations are not available.

### 3 DOSAGE FORMS AND STRENGTHS

Heparin sodium injection, USP is available as:

Heparin sodium injection, USP, **preservative free**, is available as follows:

- 2,000 USP units per 2 mL, single-dose vial

Heparin sodium injection, USP contains **benzyl alcohol** and is available as follows:

- 50,000 USP units per 10 mL, multiple-dose vial

Heparin sodium injection, USP contains **parabens** and is available as follows:

- 1,000 USP units per mL, multiple-dose vial
- 10,000 USP units per 10 mL, multiple-dose vial
- 30,000 USP units per 30 mL, multiple-dose vial
- 5,000 USP units per mL, multiple-dose vial
- 10,000 USP units per mL, multiple-dose vial
- 50,000 USP units per 5 mL, multiple-dose vial
- 20,000 USP units per mL, multiple-dose vial

### 4 CONTRAINDICATIONS

The use of heparin sodium injection is contraindicated in patients with the following conditions:

- History of heparin-induced thrombocytopenia and heparin-induced thrombocytopenia and thrombosis [*see Warnings and Precautions (5.3)*];
- Known hypersensitivity to heparin or pork products (e.g., anaphylactoid reactions) [*see Adverse Reactions (6.1)*];
- In whom suitable blood coagulation tests, e.g., the whole blood clotting time, partial thromboplastin time, etc., cannot be performed at appropriate intervals (this contraindication refers to full-dose heparin; there is usually no need to monitor coagulation parameters in patients receiving low-dose heparin);
- An uncontrolled active bleeding state [*see Warnings and Precautions (5.4)*], except when this is due to disseminated intravascular coagulation.

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Fatal Medication Errors

Do not use heparin sodium injection as a “catheter lock flush” product. Heparin sodium injection is supplied in vials containing various strengths of heparin, including vials that contain a highly concentrated solution of 10,000 units in 1 mL. Fatal hemorrhages have occurred in pediatric patients due to medication errors in which 1 mL heparin sodium injection vials were confused with 1 mL “catheter lock flush” vials. Carefully examine all heparin sodium injection vials to confirm the correct vial choice prior to administration of the drug.

#### 5.2 Hemorrhage

Avoid using heparin in the presence of major bleeding, except when the benefits of heparin therapy outweigh the potential risks.

Hemorrhage can occur at virtually any site in patients receiving heparin. Fatal hemorrhages have occurred. Adrenal hemorrhage (with resultant acute adrenal insufficiency), ovarian hemorrhage, and retroperitoneal hemorrhage have occurred during anticoagulant therapy with heparin [see *Adverse Reactions (6.1)*]. A higher incidence of bleeding has been reported in patients, particularly women, over 60 years of age [see *Clinical Pharmacology (12.3)*]. An unexplained fall in hematocrit, fall in blood pressure or any other unexplained symptom should lead to serious consideration of a hemorrhagic event.

Use heparin sodium with caution in disease states in which there is increased risk of hemorrhage, including:

- Cardiovascular - Subacute bacterial endocarditis, severe hypertension.
- Surgical - During and immediately following (a) spinal tap or spinal anesthesia or (b) major surgery, especially involving the brain, spinal cord, or eye.
- Hematologic - Conditions associated with increased bleeding tendencies, such as hemophilia, thrombocytopenia and some vascular purpuras.
- Patients with hereditary antithrombin III deficiency receiving concurrent antithrombin III therapy

- The anticoagulant effect of heparin is enhanced by concurrent treatment with antithrombin III (human) in patients with hereditary antithrombin III deficiency. To reduce the risk of bleeding, reduce the heparin dose during concomitant treatment with antithrombin III (human).

- Gastrointestinal - Ulcerative lesions and continuous tube drainage of the stomach or small intestine.
- Other - Menstruation, liver disease with impaired hemostasis.

### **5.3 Heparin-Induced Thrombocytopenia and Heparin-Induced Thrombocytopenia and Thrombosis**

Heparin-induced thrombocytopenia (HIT) is a serious antibody-mediated reaction. HIT occurs in patients treated with heparin and is due to the development of antibodies to a platelet Factor 4- heparin complex that induce *in vivo* platelet aggregation. HIT may progress to the development of venous and arterial thromboses, a condition referred to as heparin-induced thrombocytopenia with thrombosis (HITT). Thrombotic events may also be the initial presentation for HITT. These serious thromboembolic events include deep vein thrombosis, pulmonary embolism, cerebral vein thrombosis, limb ischemia, stroke, myocardial infarction, mesenteric thrombosis, renal arterial thrombosis, skin necrosis, gangrene of the extremities that may lead to amputation, and possibly death. If the platelet count falls below 100,000/mm<sup>3</sup> or if recurrent thrombosis develops, promptly discontinue heparin, evaluate for HIT and HITT, and, if necessary, administer an alternative anticoagulant.



HIT or HITT can occur up to several weeks after the discontinuation of heparin therapy. Patients presenting with thrombocytopenia or thrombosis after discontinuation of heparin sodium should be evaluated for HIT or HITT.

#### **5.4 Risk of Serious Adverse Reactions in Infants Due to Benzyl Alcohol Preservative**

Serious and fatal adverse reactions including “gaspings syndrome” can occur in neonates and infants treated with benzyl alcohol-preserved drugs, including heparin sodium injection multiple-dose vials. The “gaspings syndrome” is characterized by central nervous system depression, metabolic acidosis, and gasping respirations.

When prescribing heparin sodium injection multiple-dose vials in infants consider the combined daily metabolic load of benzyl alcohol from all sources including heparin sodium injection multiple-dose vials and other drugs containing benzyl alcohol. The minimum amount of benzyl alcohol at which toxicity may occur is not known [see *Use in Specific Populations (8.4)*].

#### **5.5 Thrombocytopenia**

Thrombocytopenia in patients receiving heparin has been reported at frequencies up to 30%. It can occur 2 to 20 days (average 5 to 9) following the onset of heparin therapy. Obtain platelet counts before and periodically during heparin therapy. Monitor thrombocytopenia of any degree closely. If the count falls below 100,000/mm<sup>3</sup> or if recurrent thrombosis develops, promptly discontinue heparin, evaluate for HIT and HITT, and, if necessary, administer an alternative anticoagulant [see *Warnings and Precautions (5.3)*].

#### **5.6 Coagulation Testing and Monitoring**

When using a full dose heparin regimen, adjust the heparin dose based on frequent blood coagulation tests. If the coagulation test is unduly prolonged or if hemorrhage occurs, discontinue heparin promptly [see *Overdosage (10)*]. Periodically monitor platelet counts, hematocrit, and occult blood in stool during the entire course of heparin therapy, regardless of the route of administration [see *Dosage and Administration (2.2)*].

#### **5.7 Heparin Resistance**

Resistance to heparin is frequently encountered in fever, thrombosis, thrombophlebitis, infections with thrombosing tendencies, myocardial infarction, cancer, in postsurgical patients, and patients with antithrombin III deficiency. Close monitoring of coagulation tests is recommended in these cases. Adjustment of heparin doses based on anti-Factor Xa levels may be warranted.

#### **5.8 Hypersensitivity**

Patients with documented hypersensitivity to heparin should be given the drug only in clearly life-threatening situations [see *Adverse Reactions (6.1)*].

Because heparin sodium injection is derived from animal tissue, it should be used with caution in patients with a history of allergy.

## 6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Hemorrhage [*see Warnings and Precautions (5.2)*]
- Heparin-Induced Thrombocytopenia and Heparin-Induced Thrombocytopenia and Thrombosis [*see Warnings and Precautions (5.3)*]
- Risk of Serious Adverse Reactions in Infants Due to Benzyl Alcohol Preservative [*see Warnings and Precautions (5.4)*]
- Thrombocytopenia [*see Warnings and Precautions (5.5)*]
- Heparin Resistance [*see Warnings and Precautions (5.7)*]
- Hypersensitivity [*see Warnings and Precautions (5.8)*]

### 6.1 Postmarketing Experience

The following adverse reactions have been identified during post approval use of heparin sodium injection. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- Hemorrhage—Hemorrhage is the chief complication that may result from heparin therapy [*see Warnings and Precautions (5.2)*]. Gastrointestinal or urinary tract bleeding during anticoagulant therapy may indicate the presence of an underlying occult lesion. Bleeding can occur at any site but certain specific hemorrhagic complications may be difficult to detect including:
  - Adrenal hemorrhage, with resultant acute adrenal insufficiency, has occurred with heparin therapy, including fatal cases.
  - Ovarian (corpus luteum) hemorrhage developed in a number of women of reproductive age receiving short- or long-term heparin therapy.
  - Retroperitoneal hemorrhage
- HIT and HITT, including delayed onset cases [*see Warnings and Precautions (5.3)*].
- Local Irritation – Local irritation, erythema, mild pain, hematoma or ulceration may follow deep subcutaneous (intrafat) injection of heparin sodium. Because these complications are much more common after intramuscular use, the intramuscular route is not recommended.
- Histamine-like reactions – Such reactions have been observed at the site of injections. Necrosis of the skin has been reported at the site of subcutaneous injection of heparin, occasionally requiring skin grafting [*see Warnings and Precautions (5.3)*].
- Hypersensitivity – Generalized hypersensitivity reactions have been reported, with chills, fever and urticaria as the most usual manifestations, and asthma, rhinitis, lacrimation, headache, nausea and vomiting, and anaphylactoid reactions, including shock, occurring less frequently. Itching and burning, especially on the plantar side of the feet, may occur. [*see Warnings and Precautions (5.8)*]
- Elevations of aminotransferases – Significant elevations of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) levels have occurred in patients who have received heparin.
- Others - Osteoporosis following long-term administration of high doses of heparin,

cutaneous necrosis after systemic administration, suppression of aldosterone synthesis, delayed transient alopecia, priapism, and rebound hyperlipemia on discontinuation of heparin sodium have also been reported.

## **7 DRUG INTERACTIONS**

### **7.1 Oral Anticoagulants**

Heparin sodium may prolong the one-stage prothrombin time. Therefore, when heparin sodium is given with dicumarol or warfarin sodium, a period of at least 5 hours after the last intravenous dose or 24 hours after the last subcutaneous dose should elapse before blood is drawn, if a valid prothrombin time is to be obtained.

### **7.2 Platelet Inhibitors**

Drugs such as NSAIDS (including salicylic acid, ibuprofen, indomethacin, and celecoxib), dextran, phenylbutazone, thienopyridines, dipyridamole, hydroxychloroquine, glycoprotein IIb/IIIa antagonists (including abciximab, eptifibatide, and tirofiban), and others that interfere with platelet- aggregation reactions (the main hemostatic defense of heparinized patients) may induce bleeding and should be used with caution in patients receiving heparin sodium. To reduce the risk of bleeding, a reduction in the dose of antiplatelet agent or heparin is recommended.

### **7.3 Other Interactions**

Digitalis, tetracyclines, nicotine or antihistamines may partially counteract the anticoagulant action of heparin sodium. Intravenous nitroglycerin administered to heparinized patients may result in a decrease of the partial thromboplastin time with subsequent rebound effect upon discontinuation of nitroglycerin. Careful monitoring of partial thromboplastin time and adjustment of heparin dosage are recommended during coadministration of heparin and intravenous nitroglycerin.

Antithrombin III (human) – The anticoagulant effect of heparin is enhanced by concurrent treatment with antithrombin III (human) in patients with hereditary antithrombin III deficiency. To reduce the risk of bleeding, a reduced dosage of heparin is recommended during treatment with antithrombin III (human).

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

There are no available data on heparin sodium use in pregnant women to inform a drug-associated risk of major birth defects and miscarriage. In published reports, heparin exposure during pregnancy did not show evidence of an increased risk of adverse maternal or fetal outcomes in humans. No teratogenicity, but early embryo-fetal death was observed in animal reproduction studies with administration of heparin sodium to pregnant rats and rabbits during organogenesis at doses approximately 10 times the maximum recommended human dose (MRHD) of 45,000 units/day ( *see Data*). Consider the benefits and risks of heparin sodium injection for the mother and possible risks to

the fetus when prescribing heparin sodium injection to a pregnant woman.

If available, preservative-free heparin sodium injection is recommended when heparin therapy is needed during pregnancy. There are no known adverse outcomes associated with fetal exposure to the preservative benzyl alcohol through maternal drug administration; however, the preservative benzyl alcohol can cause serious adverse events and death when administered intravenously to neonates and infants [*see Use in Specific Populations (8.4)*].

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

## Data

### *Human Data*

The maternal and fetal outcomes associated with uses of heparin via various dosing methods and administration routes during pregnancy have been investigated in numerous studies. These studies generally reported normal deliveries with no maternal or fetal bleeding and no other complications.

### *Animal Data*

In a published study conducted in rats and rabbits, pregnant animals received heparin intravenously during organogenesis at a dose of 10,000 units/kg/day, approximately 10 times the maximum human daily dose based on body weight. The number of early resorptions increased in both species. There was no evidence of teratogenic effects.

## **8.2 Lactation**

### Risk Summary

If available, preservative-free heparin sodium injection is recommended when heparin therapy is needed during lactation. Benzyl alcohol present in maternal serum is likely to cross into human milk and may be orally absorbed by a breastfed infant. There is no information regarding the presence of heparin sodium injection in human milk, the effects on the breastfed infant, or the effects on milk production. Due to its large molecular weight, heparin is not likely to be excreted in human milk, and any heparin in milk would not be orally absorbed by a breastfed infant. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for heparin sodium injection and any potential adverse effects on the breastfed infant from heparin sodium injection or from the underlying maternal condition [*see Use in Specific Populations (8.4)*].

## **8.4 Pediatric Use**

There are no adequate and well controlled studies on heparin use in pediatric patients. Pediatric dosing recommendations are based on clinical experience [*see Dosage and Administration (2.4)*].

Carefully examine all heparin sodium injection vials to confirm choice of the correct strength prior to administration of the drug. Pediatric patients, including neonates, have died as a result of medication errors in which Heparin Sodium Injection vials have been confused with “catheter lock flush” vials [see *Warnings and Precautions (5.1)*].

### Benzyl Alcohol Toxicity

Use preservative-free heparin sodium injection in neonates and infants.

Serious adverse reactions including fatal reactions and the “gaspings syndrome” occurred in premature neonates and infants in the neonatal intensive care unit who received drugs containing benzyl alcohol as a preservative. In these cases, benzyl alcohol dosages of 99 to 234 mg/kg/day produced high levels of benzyl alcohol and its metabolites in the blood and urine (blood levels of benzyl alcohol were 0.61 to 1.378 mmol/L). Additional adverse reactions included gradual neurological deterioration, seizures, intracranial hemorrhage, hematologic abnormalities, skin breakdown, hepatic and renal failure, hypotension, bradycardia, and cardiovascular collapse. Preterm, low-birth weight infants may be more likely to develop these reactions because they may be less able to metabolize benzyl alcohol.

### **8.5 Geriatric Use**

There are limited adequate and well-controlled studies in patients 65 years and older, however, a higher incidence of bleeding has been reported in patients, particularly women, over 60 years of age [see *Warnings and Precautions (5.2)*]. Patients over 60 years of age may require lower doses of heparin. Lower doses of heparin may be indicated in these patients [see *Clinical Pharmacology (12.3)*].

## **10 OVERDOSAGE**

Bleeding is the chief sign of heparin overdose.

### Neutralization of Heparin Effect

When clinical circumstances (bleeding) require reversal of the heparin effect, protamine sulfate (1% solution) by slow infusion will neutralize heparin sodium. **No more than 50 mg** should be administered, **very slowly**, in any 10-minute period. Each mg of protamine sulfate neutralizes approximately 100 USP heparin units. The amount of protamine required decreases over time as heparin is metabolized. Although the metabolism of heparin is complex, it may, for the purpose of choosing a protamine dose, be assumed to have a half-life of about 1/2 hour after intravenous injection. Because fatal reactions often resembling anaphylaxis have been reported with protamine, it should be given only when resuscitation techniques and treatment of anaphylactoid shock are readily available. For additional information consult the labeling of Protamine Sulfate Injection.


## **11 DESCRIPTION**

Heparin is a heterogeneous group of straight-chain anionic mucopolysaccharides, called

glycosaminoglycans, having anticoagulant properties. Although others may be present, the main sugars occurring in heparin are: (1)  $\alpha$ -L-iduronic acid 2-sulfate, (2) 2-deoxy-2-sulfamino- $\alpha$ -D-glucose 6-sulfate, (3)  $\beta$ -D-glucuronic acid, (4) 2-acetamido-2-deoxy- $\alpha$ -D-glucose and (5)  $\alpha$ -L-iduronic acid. These sugars are present in decreasing amounts, usually in the order (2) > (1) > (4) > (3) > (5), and are joined by glycosidic linkages, forming polymers of varying sizes. Heparin is strongly acidic because of its content of covalently linked sulfate and carboxylic acid groups. In heparin sodium, the acidic protons of the sulfate units are partially replaced by sodium ions.

Heparin sodium injection, USP is a sterile solution of heparin sodium derived from porcine intestinal mucosa, standardized for anticoagulant activity, in water for injection. It is to be administered by intravenous or deep subcutaneous routes. The potency is determined by a biological assay using a USP reference standard based on units of heparin activity per milligram.

Structure of heparin sodium (representative subunits):



Heparin sodium injection, USP (porcine), preservative free, is available as follows:

Each mL of the 1,000 units per mL preparation contains: 1,000 USP Heparin units (porcine); 9 mg sodium chloride; Water for Injection q.s. made isotonic with sodium chloride. Hydrochloric acid and/or sodium hydroxide may have been added for pH adjustment (5.0 to 7.5).

Heparin sodium injection, USP (porcine), preserved with benzyl alcohol, is available as follows:

Each mL of the 5,000 units per mL preparation contains: 5,000 USP Heparin units (porcine); 6 mg sodium chloride; 15 mg benzyl alcohol (as a preservative); Water for Injection q.s. Hydrochloric acid and/or sodium hydroxide may have been added for pH adjustment (5.0 to 7.5).

Heparin Sodium Injection, USP (porcine), preserved with parabens, is available as follows:

Each mL of the 1,000 units per mL preparation contains: 1,000 USP Heparin units (porcine); 9 mg sodium chloride; 1.5 mg methylparaben; 0.15 mg propylparaben; Water for Injection q.s. made isotonic with sodium chloride. Hydrochloric acid and/or sodium hydroxide may have been added for pH adjustment (5.0 to 7.5).

Each mL of the 5,000 units per mL preparation contains: 5,000 USP Heparin units (porcine); 5 mg sodium chloride; 1.5 mg methylparaben; 0.15 mg propylparaben; Water for Injection q.s. Hydrochloric acid and/or sodium hydroxide may have been added for pH adjustment (5.0 to 7.5).

Each mL of the 10,000 units per mL preparation contains: 10,000 USP Heparin units (porcine); 1.5 mg methylparaben; 0.15 mg propylparaben; Water for Injection q.s. Hydrochloric acid and/or sodium hydroxide may have been added for pH adjustment (5.0 to 7.5).

Each mL of the 20,000 units per mL preparation contains: 20,000 USP Heparin units (porcine); 1.5 mg methylparaben; 0.15 mg propylparaben; Water for Injection q.s. Hydrochloric acid and/or sodium hydroxide may have been added for pH adjustment (5.0 to 7.5).

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

Heparin interacts with the naturally occurring plasma protein, Antithrombin III, to induce a conformational change, which markedly enhances the serine protease activity of Antithrombin III, thereby inhibiting the activated coagulation factors involved in the clotting sequence, particularly Xa and IIa. Small amounts of heparin inhibit Factor Xa, and larger amounts inhibit thrombin (Factor IIa). Heparin also prevents the formation of a stable fibrin clot by inhibiting the activation of the fibrin stabilizing factor. Heparin does not have fibrinolytic activity; therefore, it will not lyse existing clots.

### **12.2 Pharmacodynamics**

Various times (activated clotting time, activated partial thromboplastin time, prothrombin time, whole blood clotting time) are prolonged by full therapeutic doses of heparin; in most cases, they are not measurably affected by low doses of heparin. The bleeding time is usually unaffected by heparin.

### **12.3 Pharmacokinetics**

#### **Absorption**

Heparin is not absorbed through the gastrointestinal tract and therefore administered via parenteral route. Peak plasma concentration and the onset of action are achieved immediately after intravenous administration.

#### **Distribution**

Heparin is highly bound to antithrombin, fibrinogens, globulins, serum proteases and lipoproteins. The volume of distribution is 0.07 L/kg.

## Elimination

### *Metabolism*

Heparin does not undergo enzymatic degradation.

### *Excretion*

Heparin is mainly cleared from the circulation by liver and reticuloendothelial cells mediated uptake into extravascular space. Heparin undergoes biphasic clearance, a) rapid saturable clearance (zero order process due to binding to proteins, endothelial cells and macrophage) and b) slower first order elimination. The plasma half-life is dose-dependent and it ranges from 0.5 to 2 h.

## Specific Populations

### *Geriatric patients*

Patients over 60 years of age, following similar doses of heparin, may have higher plasma levels of heparin and longer activated partial thromboplastin times (aPTTs) compared with patients under 60 years of age [see *Use in Specific Populations (8.5)*].

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term studies in animals have been performed to evaluate carcinogenic potential of heparin. Also, no reproduction studies in animals have been performed concerning mutagenesis or impairment of fertility.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Heparin sodium injection, USP (porcine), **preservative free**, is available as follows:

<b>NDC Number</b>	<b>Concentration</b>	<b>Vial Fill Volume</b>	<b>Vial Type</b>	<b>Pack Size</b>
63739-900-26	1,000 USP Heparin Units/mL	2 mL	Single-Dose Vial	Carton of 25 vials

Use only if solution is clear and seal intact. Do not use if solution is discolored or contains a precipitate.

This container closure is not made with natural rubber latex.

Discard unused portion.



Heparin Sodium Injection, USP (porcine) contains **benzyl alcohol** and is available as follows:

<b>NDC Number</b>	<b>Concentration</b>	<b>Vial Fill Volume</b>	<b>Vial Type</b>	<b>Pack Size</b>
63739-901-28	5,000 USP Heparin Units/mL	10 mL	Multiple-Dose Vial	Carton of 25 vials

Use only if solution is clear and seal intact. Do not use if solution is discolored or contains a precipitate.

This container closure is not made with natural rubber latex.

Heparin Sodium Injection, USP (porcine) contains **parabens** and is available as follows:

<b>NDC Number</b>	<b>Concentration</b>	<b>Vial Fill Volume</b>	<b>Vial Type</b>	<b>Pack Size</b>
63739-920-25	1,000 USP Heparin Units/mL	1 mL	Multiple-Dose Vial	Carton of 25 vials
63739-931-28	1,000 USP Heparin Units/mL	10 mL	Multiple-Dose Vial	Carton of 25 vials
63739-942-29	1,000 USP Heparin Units/mL	30 mL	Multiple-Dose Vial	Carton of 25 vials
63739-953-25	5,000 USP Heparin Units/mL	1 mL	Multiple-Dose Vial	Carton of 25 vials
63739-964-25	10,000 USP Heparin Units/mL	1 mL	Multiple-Dose Vial	Carton of 25 vials
63739-975-27	10,000 USP Heparin Units/mL	5 mL	Multiple-Dose Vial	Carton of 25 vials
63739-986-25	20,000 USP Heparin Units/mL	1 mL	Multiple-Dose Vial	Carton of 25 vials

Use only if solution is clear and seal intact. Do not use if solution is discolored or contains a precipitate.

This container closure is not made with natural rubber latex.

#### **STORAGE:**

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

## **17 PATIENT COUNSELING INFORMATION**

### Hemorrhage

Inform patients that it may take them longer than usual to stop bleeding, that they may bruise and/or bleed more easily when they are treated with heparin, and that they should report any unusual bleeding or bruising to their physician. Hemorrhage can occur at

virtually any site in patients receiving heparin. Fatal hemorrhages have occurred [see *Warnings and Precautions (5.2)*].

#### Prior to Surgery

Advise patients to inform physicians and dentists that they are receiving heparin before any surgery is scheduled [see *Warnings and Precautions (5.2)*].

#### Heparin-Induced Thrombocytopenia

Inform patients of the risk of heparin-induced thrombocytopenia (HIT). HIT may progress to the development of venous and arterial thromboses, a condition known as heparin-induced thrombocytopenia and thrombosis (HITT). HIT and HITT can occur up to several weeks after the discontinuation of heparin therapy [see *Warnings and Precautions (5.3)*].

#### Hypersensitivity

Inform patients that generalized hypersensitivity reactions have been reported. Necrosis of the skin has been reported at the site of subcutaneous injection of heparin [see *Warnings and Precautions (5.8), Adverse Reactions (6.1)*].

#### Other Medications

Because of the risk of hemorrhage, advise patients to inform their physicians and dentists of all medications they are taking, including non-prescription medications, and before starting any new medication [see *Drug Interactions (7.1)*].

Manufactured by:

#### **Shenzhen Techdow Pharmaceutical Co., Ltd.**

19 Gaoxinzongyi Road, Nanshan District,  
Shenzhen, P.R. China, 518057

Distributed by:

#### **McKesson Corporation dba SKY Packaging**

Memphis, TN 38141.

Toll Free: 1-888-243-4363

Revised: 01/2022

#### **PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 1,000 USP Units/mL - 1 mL Container Label**

NDC 63739-920-11 Rx only

HEPARIN

SODIUM INJECTION, USP


1,000 USP Units/mL

For IV or SC use

Multiple Dose Vial

1 mL

NOT FOR LOCK FLUSH



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 1,000 USP Units/mL - 1 mL  
Container-Carton [25 Multiple Dose Vials]**

NDC 63739-920-25 Rx only

HEPARIN

SODIUM INJECTION, USP

1,000 USP Units/mL

1 mL


NOT FOR LOCK FLUSH

For Intravenous or

Subcutaneous use

25 Multiple-Dose Vials

Derived from Porcine Intestinal Mucosa



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 10,000 USP Units/10 mL (1,000 USP Units/mL) - 10 mL Container Label**

NDC 63739-931-14 Rx only

HEPARIN

SODIUM INJECTION, USP

10,000 USP Units/10 mL

(1,000 USP Units/mL)

For IV or SC use

10 mL Multiple Dose Vial

NOT FOR LOCK FLUSH

NDC 63739-931-14 Rx only  
**HEPARIN**  
SODIUM INJECTION, USP  
10,000<sup>USP</sup> Units/10 mL  
(1,000 USP Units/mL)  
**NOT FOR LOCK FLUSH**  
For IV or SC use  
10 mL Multiple Dose Vial

Derived from Porcine Intestinal Mucosa.  
Sterile, Nonpyrogenic.  
Store at 20° to 25°C (68° to 77°F).  
KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN  
Mfd. in CHINA for Mckesson Corporation  
dba SKY Packaging.

R11/2021

Lot  
Exp

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 10,000 USP Units/10 mL (1,000 USP Units/mL) - 10 mL Container-Carton [25 Multiple Dose Vials]**

NDC 63739-931-28 Rx only

HEPARIN

SODIUM INJECTION, USP

10,000 USP Units/10 mL

(1,000 USP Units/mL)

10 mL

NOT FOR LOCK FLUSH

For Intravenous or

Subcutaneous use

25 Multiple-Dose Vials

Derived from Porcine Intestinal Mucosa



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 30,000 USP Units/30 mL (1,000 USP Units/mL) - 30 mL Container Label**

NDC 63739-942-15 Rx only

HEPARIN

SODIUM INJECTION, USP

30,000 USP Units/30 mL

(1,000 USP Units/mL)

For IV or SC use

Multiple Dose Vial

30 mL

NOT FOR LOCK FLUSH

NDC 63739-942-15 Rx only

**HEPARIN**  
**SODIUM INJECTION, USP**

**30,000 USP Units/30 mL**  
**(1,000 USP Units/mL)**

**NOT FOR LOCK FLUSH**

**For IV or SC use**  
**Multiple Dose Vial**  
**30 mL**

WZ  
 0  
 1  
 2  
 3  
 4  
 5  
 6  
 7  
 8  
 9  
 0  
 1  
 2  
 3  
 4  
 5  
 6  
 7  
 8  
 9  
 0

Derived from Porcine Intestinal Mucosa  
 Sterile, Nonpyrogenic  
**Store at 20° to 25°C (68° to 77°F)**[See USP Controlled Room Temperature].  
**KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN**  
 Mfd. in CHINA for McKesson Corporation dba SKY Packaging

Lot: R12/2022  
 Exp:

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 30,000 USP Units/30 mL (1,000 USP Units/mL) - 30 mL Container-Carton [25 Multiple Dose Vials]**

NDC 63739-942-29 Rx only

HEPARIN

SODIUM INJECTION, USP

30,000 USP Units/30 mL

(1,000 USP Units/mL)

30 mL

NOT FOR LOCK FLUSH

For Intravenous or

Subcutaneous use

25 Multiple-Dose Vials

Derived from Porcine Intestinal Mucosa



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 5,000 USP Units/mL - 1 mL Container Label**

NDC 63739-953-11 Rx only

HEPARIN

SODIUM INJECTION, USP

5,000 USP Units/mL


For IV or SC use

Multiple Dose Vial

1 mL

NOT FOR LOCK FLUSH





**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 5,000 USP Units/mL - 1 mL  
Container-Carton [25 Multiple Dose Vials]**

NDC 63739-953-25 Rx only

HEPARIN

SODIUM INJECTION, USP

5,000 USP Units/mL

1 mL


NOT FOR LOCK FLUSH

For Intravenous or

Subcutaneous use

25 Multiple-Dose Vials

Derived from Porcine Intestinal Mucosa



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 10,000 USP Units/mL - 1 mL Container Label**

NDC 63739-964-11 Rx only

HEPARIN

SODIUM INJECTION, USP


10,000 USP Units/mL

For IV or SC use

Multiple Dose Vial

1 mL

NOT FOR LOCK FLUSH



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 10,000 USP Units/mL - 1 mL Container-Carton [25 Multiple Dose Vials]**

NDC 63739-964-25 Rx only

HEPARIN

SODIUM INJECTION, USP

10,000 USP Units/mL

1 mL

NOT FOR LOCK FLUSH

For Intravenous or

Subcutaneous use

25 Multiple-Dose Vials

Derived from Porcine Intestinal Mucosa



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 50,000 USP Units/5 mL (10,000 USP Units/mL) - 5 mL Container Label**

NDC 63739-975-13 Rx only

HEPARIN

SODIUM INJECTION, USP

50,000 USP Units/5 mL

(10,000 USP Units/mL)

For IV or SC use

Multiple Dose Vial

5 mL

NOT FOR LOCK FLUSH

NDC 63739-975-13 Rx only  
**HEPARIN**  
SODIUM INJECTION, USP  
**50,000** USP Units/5 mL  
(10,000 USP Units/mL)  
**NOT FOR LOCK FLUSH**

Derived from Porcine Intestinal Mucosa. Sterile, Nonpyrogenic.  
**Store at 20° to 25°C (68° to 77°F).** Mfd. in CHINA for  
McKesson Corporation dba SKY  
Packaging R12/2022

For IV or SC use  
Multiple Dose Vial  
5 mL

Lot  
Exp

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 50,000 USP Units/5 mL (10,000 USP Units/mL) - 5 mL Container-Carton [25 Multiple Dose Vials]**

NDC 63739-975-27 Rx only

HEPARIN

SODIUM INJECTION, USP

50,000 USP Units/5 mL

(10,000 USP Units/mL)

5 mL

NOT FOR LOCK FLUSH

For Intravenous or Subcutaneous use

25 Multiple-Dose Vials

Derived from Porcine Intestinal Mucosa



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 20,000 USP Units/mL - 1 mL Container Label**

NDC 63739-986-11 Rx only

HEPARIN

SODIUM INJECTION, USP


20,000 USP Units/mL

For IV or SC use

Multiple Dose Vial

1 mL

NOT FOR LOCK FLUSH



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 20,000 USP Units/mL - 1 mL  
Container-Carton [25 Multiple Dose Vials]**

NDC 63739-986-25 Rx only

HEPARIN

SODIUM INJECTION, USP

20,000 USP Units/mL

1 mL


NOT FOR LOCK FLUSH

For Intravenous or

Subcutaneous use

25 Multiple-Dose Vials

Derived from Porcine Intestinal Mucosa



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL- 2,000 USP Units/2 mL (1,000 USP Units/mL) - 2 mL Container Label**

NDC 63739-900-12 Rx only

HEPARIN

SODIUM INJECTION, USP

2,000 USP Units/2 mL

(1,000 USP Units/mL)


For IV or SC use

Preservative Free

Discard unused portion



2 mL Single Dose Vial  
NOT FOR LOCK FLUSH



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL- 2,000 USP Units/2 mL (1,000 USP Units/mL) - 2 mL Container-Carton [25 Single Dose Vials]**

NDC 63739-900-26 Rx only

HEPARIN

SODIUM INJECTION, USP

2,000 USP Units/2 mL

(1,000 USP Units/mL)

2 mL

NOT FOR LOCK FLUSH


For Intravenous or

Subcutaneous use

Preservative Free

25 Single-Dose Vials

Derived from Porcine Intestinal Mucosa



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL- 50,000 USP Units/10 mL (5,000 USP Units/mL) - 10 mL Container Label**

NDC 63739-901-14 Rx only

HEPARIN

SODIUM INJECTION, USP

50,000 USP Units/10 mL

(5,000 USP Units/mL)

For IV or SC use

Warning: Contains Benzyl Alcohol

10 mL Multiple Dose Vial

NOT FOR LOCK FLUSH

NDC 63739-901-14 Rx only  
**HEPARIN**  
SODIUM INJECTION, USP  
**50,000<sup>USP</sup> Units/10 mL**  
(5,000 USP Units/mL)  
**NOT FOR LOCK FLUSH**  
For IV or SC use  
Warning: Contains Benzyl Alcohol  
**10 mL** Multiple Dose Vial

WZ  
0373901148

Derived from Porcine Intestinal Mucosa.  
Sterile, Nonpyrogenic.  
Store at 20° to 25°C (68° to 77°F).[See  
USP Controlled Room Temperature].  
KEEP THIS AND ALL DRUGS OUT OF THE  
REACH OF CHILDREN  
Mfd. in CHINA for Mckesson Corporation  
dba SKY Packaging.

R05/2022

Lot  
Exp

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL- 50,000 USP Units/10 mL (5,000 USP Units/mL) - 10 mL Container-Carton [25 Multiple Dose Vials]**

NDC 63739-901-28 Rx only

HEPARIN

SODIUM INJECTION, USP

50,000 USP Units/10 mL

(5,000 USP Units/mL)

10 mL

NOT FOR LOCK FLUSH

Warning: Contains Benzyl Alcohol

For Intravenous or Subcutaneous use

25 Multiple-Dose Vials

Derived from Porcine Intestinal Mucosa



## HEPARIN SODIUM

heparin sodium injection, solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:63739-920
<b>Route of Administration</b>	INTRAVENOUS, SUBCUTANEOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HEPARIN SODIUM (UNII: ZZ45AB24CA) (HEPARIN - UNII:T2410KM04A)	HEPARIN	1000 [USP'U] in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	9 mg in 1 mL
METHYLPARABEN (UNII: A2I8C7HI9T)	1.5 mg in 1 mL
PROPYLPARABEN (UNII: Z8IX2SC1OH)	0.15 mg in 1 mL
WATER (UNII: 059QF0KO0R)	

<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63739-920-25	25 in 1 CARTON	06/12/2014	08/31/2024
1	NDC:63739-920-11	1 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202957	06/12/2014	

## HEPARIN SODIUM

heparin sodium injection, solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:63739-931
<b>Route of Administration</b>	SUBCUTANEOUS, INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>HEPARIN SODIUM</b> (UNII: ZZ45AB24CA) (HEPARIN - UNII:T2410KM04A)	HEPARIN	1000 [USP'U] in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	9 mg in 1 mL
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	1.5 mg in 1 mL
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	0.15 mg in 1 mL
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63739-931-28	25 in 1 CARTON	06/12/2014	05/31/2025
	NDC:63739-931-11	1 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a		

<b>1</b>	NDC:63739-931-14	10 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
ANDA	ANDA202957	06/12/2014		

## HEPARIN SODIUM

heparin sodium injection, solution

<b>Product Information</b>			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:63739-942
<b>Route of Administration</b>	INTRAVENOUS, SUBCUTANEOUS		

<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
HEPARIN SODIUM (UNII: ZZ45AB24CA) (HEPARIN - UNII:T2410KM04A)	HEPARIN	1000 [USP'U] in 1 mL	

<b>Inactive Ingredients</b>		
<b>Ingredient Name</b>	<b>Strength</b>	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	9 mg in 1 mL	
METHYLPARABEN (UNII: A2I8C7HI9T)	1.5 mg in 1 mL	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	0.15 mg in 1 mL	
WATER (UNII: 059QF0KO0R)		
HYDROCHLORIC ACID (UNII: QTT17582CB)		
SODIUM HYDROXIDE (UNII: 55X04QC32I)		

<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:63739-942-29	25 in 1 CARTON	06/12/2014	09/30/2020
1	NDC:63739-942-15	30 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

<b>Marketing Information</b>			
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA202957	06/12/2014	

# HEPARIN SODIUM

heparin sodium injection, solution

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:63739-953
<b>Route of Administration</b>	SUBCUTANEOUS, INTRAVENOUS		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HEPARIN SODIUM (UNII: ZZ45AB24CA) (HEPARIN - UNII:T2410KM04A)	HEPARIN	5000 [USP'U] in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	5 mg in 1 mL
METHYLPARABEN (UNII: A2I8C7HI9T)	1.5 mg in 1 mL
PROPYLPARABEN (UNII: Z8IX2SC1OH)	0.15 mg in 1 mL
WATER (UNII: 059QF0KO0R)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63739-953-25	25 in 1 CARTON	06/12/2014	04/30/2025
1	NDC:63739-953-11	1 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202957	06/12/2014	

# HEPARIN SODIUM

heparin sodium injection, solution

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:63739-964
<b>Route of Administration</b>	INTRAVENOUS, SUBCUTANEOUS		

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
HEPARIN SODIUM (UNII: ZZ45AB24CA) (HEPARIN - UNII:T2410KM04A)			HEPARIN	10000 [USP'U] in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
METHYLPARABEN (UNII: A2I8C7HI9T)	1.5 mg in 1 mL
PROPYLPARABEN (UNII: Z8IX2SC1OH)	0.15 mg in 1 mL
WATER (UNII: 059QF0KO0R)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63739-964-25	25 in 1 CARTON	06/12/2014	07/31/2024
1	NDC:63739-964-11	1 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA203198	06/12/2014	

HEPARIN SODIUM			
heparin sodium injection, solution			

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:63739-975
Route of Administration	SUBCUTANEOUS, INTRAVENOUS		

Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
HEPARIN SODIUM (UNII: ZZ45AB24CA) (HEPARIN - UNII:T2410KM04A)		HEPARIN	10000 [USP'U] in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
METHYLPARABEN (UNII: A2I8C7HI9T)	1.5 mg in 1 mL
PROPYLPARABEN (UNII: Z8IX2SC1OH)	0.15 mg in 1 mL
WATER (UNII: 059QF0KO0R)	



**HYDROCHLORIC ACID** (UNII: QTT17582CB)

**SODIUM HYDROXIDE** (UNII: 55X04QC32I)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63739-975-27	25 in 1 CARTON	06/12/2014	06/13/2014
1	NDC:63739-975-13	5 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA203198	06/12/2014	

## HEPARIN SODIUM

heparin sodium injection, solution

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:63739-986
Route of Administration	SUBCUTANEOUS, INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>HEPARIN SODIUM</b> (UNII: ZZ45AB24CA) (HEPARIN - UNII:T2410KM04A)	HEPARIN	20000 [USP'U] in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	1.5 mg in 1 mL
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	0.15 mg in 1 mL
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63739-986-25	25 in 1 CARTON	06/12/2014	08/31/2024
1	NDC:63739-986-11	1 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA203198	06/12/2014	

## HEPARIN SODIUM

heparin sodium injection, solution

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:63739-900
Route of Administration	INTRAVENOUS, SUBCUTANEOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HEPARIN SODIUM (UNII: ZZ45AB24CA) (HEPARIN - UNII:T2410KM04A)	HEPARIN	1000 [USP'U] in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	9 mg in 1 mL
WATER (UNII: 059QF0KO0R)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63739-900-26	25 in 1 CARTON	06/12/2014	04/30/2018
1	NDC:63739-900-12	2 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202732	06/12/2014	

## HEPARIN SODIUM

heparin sodium injection, solution

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:63739-901
<b>Route of Administration</b>	SUBCUTANEOUS, INTRAVENOUS		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>HEPARIN SODIUM</b> (UNII: ZZ45AB24CA) (HEPARIN - UNII:T2410KM04A)	HEPARIN	5000 [USP'U] in 1 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	15 mg in 1 mL
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	6 mg in 1 mL
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63739-901-28	25 in 1 CARTON	06/12/2014	12/31/2024
1	NDC:63739-901-14	10 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202733	06/12/2014	

**Labeler** - McKesson Corporation dba SKY Packaging (140529962)**Establishment**

Name	Address	ID/FEI	Business Operations
Shenzhen Techdow Pharmaceutical Co., Ltd.		527809171	analysis(63739-900, 63739-901, 63739-920, 63739-931, 63739-942, 63739-953, 63739-964, 63739-975, 63739-986) , manufacture(63739-900, 63739-901, 63739-920, 63739-931, 63739-942, 63739-953, 63739-964, 63739-975, 63739-986)

**Establishment**

Name	Address	ID/FEI	Business Operations
Shenzhen Hepalink Pharmaceutical Group Co., Ltd.		554436033	api manufacture(63739-901, 63739-920, 63739-931, 63739-942, 63739-953, 63739-964, 63739-900, 63739-975, 63739-986)

## Establishment

Name	Address	ID/FEI	Business Operations
Eugia Pharma Specialities Limited		650498244	analysis(63739-900, 63739-901, 63739-920, 63739-931, 63739-942, 63739-953, 63739-964, 63739-975, 63739-986) , manufacture(63739-900, 63739-901, 63739-920, 63739-931, 63739-942, 63739-953, 63739-964, 63739-975, 63739-986)

Revised: 1/2025

McKesson Corporation dba SKY Packaging