

METHOCARBAMOL- methocarbamol tablet, film coated
Asclemed USA, Inc.


Methocarbamol Tablets, USP

Rx only

DESCRIPTION

Methocarbamol tablets, USP, a carbamate derivative of guaifenesin, is a central nervous system (CNS) depressant with sedative and musculoskeletal relaxant properties.

The chemical name of methocarbamol is 3-(2-methoxyphenoxy)-1,2-propanediol 1-carbamate and has the empirical formula $C_{11}H_{15}NO_5$. Its molecular weight is 241.24. The structural formula is shown below.



Methocarbamol is a white bulky powder, sparingly soluble in water and chloroform, soluble in alcohol only with heating and insoluble in benzene and *n*-hexane.

Methocarbamol tablets USP, 500 mg are available as white in color, round, beveled edge, biconvex film-coated tablet containing 500 mg of methocarbamol USP for oral administration.

Methocarbamol tablets USP, 750 mg are available as white in color, biconvex capsule shaped film-coated tablet containing 750 mg of methocarbamol USP for oral administration.

Methocarbamol tablets USP, 500 mg and 750 mg contain the following inactive ingredients: corn starch, hypromellose, magnesium stearate, polyethylene glycol, povidone, sodium lauryl sulfate, sodium starch glycolate, talc and titanium dioxide.

FDA approved dissolution test specifications differ from USP for 500 mg.

CLINICAL PHARMACOLOGY

The mechanism of action of methocarbamol in humans has not been established, but may be due to general CNS depression. It has no direct action on the contractile mechanism of striated muscle, the motor end plate or the nerve fiber.

Pharmacokinetics

In healthy volunteers, the plasma clearance of methocarbamol ranges between 0.20 and 0.80 L/h/kg, the mean plasma elimination half-life ranges between 1 and 2 hours, and the plasma protein binding ranges between 46% and 50%.

Methocarbamol is metabolized via dealkylation and hydroxylation. Conjugation of methocarbamol also is likely. Essentially all methocarbamol metabolites are eliminated in the urine. Small amounts of unchanged methocarbamol also are excreted in the urine.

Special populations

Elderly

The mean ($\pm$ SD) elimination half-life of methocarbamol in elderly healthy volunteers (mean [$\pm$ SD] age, 69 [$\pm$ 4] years) was slightly prolonged compared to a younger (mean [$\pm$ SD] age, 53.3 [$\pm$ 8.8] years), healthy population (1.5 [$\pm$ 0.4] hours versus 1.1 [$\pm$ 0.27] hours, respectively). The fraction of bound methocarbamol was slightly decreased in the elderly versus younger volunteers (41% to 43% versus 46% to 50%, respectively).

Renally impaired

The clearance of methocarbamol in 8 renally-impaired patients on maintenance hemodialysis was reduced about 40% compared to 17 normal subjects, although the mean ($\pm$ SD) elimination half-life in these two groups was similar: 1.2 ($\pm$ 0.6) versus 1.1 ($\pm$ 0.3) hours, respectively.

Hepatically impaired

In 8 patients with cirrhosis secondary to alcohol abuse, the mean total clearance of methocarbamol was reduced approximately 70% compared to that obtained in 8 age- and weight-matched normal subjects. The mean ($\pm$ SD) elimination half-life in the cirrhotic patients and the normal subjects was 3.38 ($\pm$ 1.62) hours and 1.11 ($\pm$ 0.27) hours, respectively. The percent of methocarbamol bound to plasma proteins was decreased to approximately 40% to 45% compared to 46% to 50% in the normal subjects.

INDICATIONS AND USAGE

Methocarbamol tablets are indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute, painful musculoskeletal conditions. The mode of action of methocarbamol has not been clearly identified, but may be related to its sedative properties. Methocarbamol does not directly relax tense

skeletal muscles in man.

CONTRAINDICATIONS

Methocarbamol tablets are contraindicated in patients hypersensitive to methocarbamol or to any of the tablet components.

WARNINGS

Since methocarbamol may possess a general CNS depressant effect, patients receiving methocarbamol should be cautioned about combined effects with alcohol and other CNS depressants.

Safe use of methocarbamol has not been established with regard to possible adverse effects upon fetal development. There have been reports of fetal and congenital abnormalities following in utero exposure to methocarbamol. Therefore, methocarbamol should not be used in women who are or may become pregnant and particularly during early pregnancy unless in the judgment of the physician the potential benefits outweigh the possible hazards (see **PRECAUTIONS, Pregnancy**).

Use in Activities Requiring Mental Alertness

Methocarbamol may impair mental and/or physical abilities required for performance of hazardous tasks, such as operating machinery or driving a motor vehicle. Patients should be cautioned about operating machinery, including automobiles, until they are reasonably certain that methocarbamol therapy does not adversely affect their ability to engage in such activities.

PRECAUTIONS

Information for Patients

Patients should be cautioned that methocarbamol may cause drowsiness or dizziness, which may impair their ability to operate motor vehicles or machinery.

Because methocarbamol may possess a general CNS-depressant effect, patients should be cautioned about combined effects with alcohol and other CNS depressants.

Drug Interactions

See **WARNINGS** and **PRECAUTIONS** for interaction with CNS drugs and alcohol.

Methocarbamol may inhibit the effect of pyridostigmine bromide. Therefore, methocarbamol should be used with caution in patients with myasthenia gravis receiving anticholinesterase agents.

Drug/Laboratory Test Interactions

Methocarbamol may cause a color interference in certain screening tests for 5-hydroxyindoleacetic acid (5-HIAA) using nitrosonaphthol reagent and in screening tests for urinary vanillylmandelic acid (VMA) using the Gitlow method.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies to evaluate the carcinogenic potential of methocarbamol have not been performed. No studies have been conducted to assess the effect of methocarbamol on mutagenesis or its potential to impair fertility.

Pregnancy

Teratogenic Effects

Animal reproduction studies have not been conducted with methocarbamol. It is also not known whether methocarbamol can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Methocarbamol should be given to a pregnant woman only if clearly needed.

Safe use of methocarbamol has not been established with regard to possible adverse effects upon fetal development. There have been reports of fetal and congenital abnormalities following in utero exposure to methocarbamol. Therefore, methocarbamol should not be used in women who are or may become pregnant and particularly during early pregnancy unless in the judgment of the physician the potential benefits outweigh the possible hazards (see **WARNINGS**).

Nursing Mothers

Methocarbamol and/or its metabolites are excreted in the milk of dogs; however, it is not known whether methocarbamol or its metabolites are excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when methocarbamol is administered to a nursing woman.

Pediatric Use

Safety and effectiveness of methocarbamol in pediatric patients below the age of 16 have not been established.

ADVERSE REACTIONS

Adverse reactions reported coincident with the administration of methocarbamol include:

*Body as a whole:*Anaphylactic reaction, angioneurotic edema, fever, headache

*Cardiovascular system:*Bradycardia, flushing, hypotension, syncope, thrombophlebitis

*Digestive system:*Dyspepsia, jaundice (including cholestatic jaundice), nausea and vomiting

*Hemic and lymphatic system:*Leukopenia

*Immune system:*Hypersensitivity reactions

Nervous system: Amnesia, confusion, diplopia, dizziness or lightheadedness, drowsiness, insomnia, mild muscular incoordination, nystagmus, sedation, seizures (including grand mal), vertigo

*Skin and special senses:*Blurred vision, conjunctivitis, nasal congestion, metallic taste, pruritus, rash, urticaria

OVERDOSAGE

Limited information is available on the acute toxicity of methocarbamol. Overdose of methocarbamol is frequently in conjunction with alcohol or other CNS depressants and includes the following symptoms: nausea, drowsiness, blurred vision, hypotension, seizures, and coma.

In post-marketing experience, deaths have been reported with an overdose of methocarbamol alone or in the presence of other CNS depressants, alcohol or psychotropic drugs.

Treatment

Management of overdose includes symptomatic and supportive treatment. Supportive measures include maintenance of an adequate airway, monitoring urinary output and vital signs, and administration of intravenous fluids if necessary. The usefulness of hemodialysis in managing overdose is unknown.

DOSAGE AND ADMINISTRATION

Methocarbamol tablets, 500 mg - Adults:

Initial dosage: 3 tablets 4 times daily.

Maintenance dosage: 2 tablets 4 times daily.

Methocarbamol tablets: 750 mg - Adults:

Initial dosage: 2 tablets 4 times daily.

Maintenance dosage: 1 tablet q.4h. or 2 tablets 3 times daily.

Six grams a day are recommended for the first 48 to 72 hours of treatment. (For severe conditions 8 grams a day may be administered.) Thereafter, the dosage can usually be reduced to approximately 4 grams a day.

HOW SUPPLIED

Methocarbamol Tablets USP, 500 mg are white in color, round, beveled edge, biconvex film-coated tablet debossed with “500” above the score line on one side and “METT” on the other side. They are supplied as follows:

Bottles of 10 NDC 87063-040-10 (repackaged from NDC 59651-340-XX)

Bottles of 20 NDC 87063-040-20 (repackaged from NDC 59651-340-XX)

Bottles of 30 NDC 87063-040-30 (repackaged from NDC 59651-340-XX)

Bottles of 60 NDC 87063-040-60 (repackaged from NDC 59651-340-XX)

Bottles of 90 NDC 87063-040-90 (repackaged from NDC 59651-340-XX)

Bottles of 100 NDC 87063-040-01 (relabeled from NDC 59651-340-01)

Methocarbamol Tablets USP, 750 mg are white in color, biconvex capsule shaped film-coated tablet, debossed with “METT” on one side and “750” on the other side. They are supplied as follows:

Bottles of 10 NDC 87063-041-10 (repackaged from NDC 59651-341-XX)

Bottles of 20 NDC 87063-041-20 (repackaged from NDC 59651-341-XX)

Bottles of 30 NDC 87063-041-30 (repackaged from NDC 59651-341-XX)

Bottles of 60 NDC 87063-041-60 (repackaged from NDC 59651-341-XX)

Bottles of 90 NDC 87063-041-90 (repackaged from NDC 59651-341-XX)

Bottles of 100 NDC 87063-041-01 (relabeled from NDC 59651-341-01)

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Dispense in tight container.

For more information, call Aurobindo Pharma USA, Inc. at 1-866-850-2876.

Relabeled and Repackaged by:

Enovachem PHARMACEUTICALS

Torrance, CA 90501

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 500 mg

Packaged By:

Enovachem
PHARMACEUTICALS
379 Van Ness Ave.
Suite 1403-1406
Torrance, CA 90501

Methocarbamol Tablets, USP 500 mg

NDC: 87063-040-10
Qty: 10

Distributed By: Aurobindo Pharma USA, Inc.
Source NDC: 59651-340-XX
Description: white/round, film coated tablet/debossed 500, METT
Lot #: XXXXXXXX Exp:
Batch #: XXXXXXXX
Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

(01) 0 0387063 04010 6
(17)
(10) XXXXXXXX
(21)

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-10
S/N:
Qty: 10

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-10
S/N:
Qty: 10

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-10
S/N:
Qty: 10

Relabeled By:

Enovachem
PHARMACEUTICALS
379 Van Ness Ave.
Suite 1403-1406
Torrance, CA 90501

Methocarbamol Tablets, USP 500 mg

NDC: 87063-040-01
Qty: 100

Distributed By: Aurobindo Pharma USA, Inc.
Source NDC: 59651-340-01
Description: white/round, film coated tablet/debossed 500, METT
Lot #: XXXXXXXX Exp:
Batch #: XXXXXXXX
Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

(01) 0 0387063 04001 4
(17)
(10) XXXXXXXX
(21)

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-01
S/N:
Qty: 100

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-01
S/N:
Qty: 100

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-01
S/N:
Qty: 100

Packaged By:

Enovachem
PHARMACEUTICALS
379 Van Ness Ave.
Suite 1403-1406
Torrance, CA 90501

Methocarbamol Tablets, USP 500 mg

NDC: 87063-040-20
Qty: 20

Distributed By: Aurobindo Pharma USA, Inc.
Source NDC: 59651-340-XX
Description: white/round, film coated tablet/debossed 500, METT
Lot #: XXXXXXXX Exp:
Batch #: XXXXXXXX
Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

(01) 0 0387063 04020 5
(17)
(10) XXXXXXXX
(21)

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-20
S/N:
Qty: 20

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-20
S/N:
Qty: 20

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-20
S/N:
Qty: 20

Packaged By:

Enovachem
PHARMACEUTICALS
379 Van Ness Ave.
Suite 1403-1406
Torrance, CA 90501

Methocarbamol Tablets, USP 500 mg

NDC: 87063-040-30
Qty: 30

Distributed By: Aurobindo Pharma USA, Inc.
Source NDC: 59651-340-XX
Description: white/round, film coated tablet/debossed 500, METT
Lot #: XXXXXXXX Exp:
Batch #: XXXXXXXX
Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

(01) 0 0387063 04030 4
(17)
(10) XXXXXXXX
(21)

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-30
S/N:
Qty: 30

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-30
S/N:
Qty: 30

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-30
S/N:
Qty: 30

Packaged By:

Enovachem
PHARMACEUTICALS
379 Van Ness Ave.
Suite 1403-1406
Torrance, CA 90501

Methocarbamol Tablets, USP 500 mg

NDC: 87063-040-60
Qty: 60

Distributed By: Aurobindo Pharma USA, Inc.
Source NDC: 59651-340-XX
Description: white/round, film coated tablet/debossed 500, METT
Lot #: XXXXXXXX Exp:
Batch #: XXXXXXXX
Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

(01) 0 0387063 04060 1
(17)
(10) XXXXXXXX
(21)

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-60
S/N:
Qty: 60

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-60
S/N:
Qty: 60

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-60
S/N:
Qty: 60

Packaged By:

Enovachem
PHARMACEUTICALS
379 Van Ness Ave.
Suite 1403-1406
Torrance, CA 90501

Methocarbamol Tablets, USP 500 mg

NDC: 87063-040-90
Qty: 90

Distributed By: Aurobindo Pharma USA, Inc.
Source NDC: 59651-340-XX
Description: white/round, film coated tablet/debossed 500, METT
Lot #: XXXXXXXX Exp:
Batch #: XXXXXXXX
Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

(01) 0 0387063 04090 8
(17)
(10) XXXXXXXX
(21)

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-90
S/N:
Qty: 90

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-90
S/N:
Qty: 90

Methocarbamol Tablets, USP 500 mg
NDC: 87063-040-90
S/N:
Qty: 90

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 750 mg

Packaged By:
Enovachem
PHARMACEUTICALS
375 Van Ness Ave.
Suite 1403-1405
Torrance, CA 90501
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-10
Qty: 10
Distributed By: Aurobindo Pharma USA, Inc.
Source NDC: 59651-341-XX
Description: white capsule shaped, firm-coated tablet/debossed METT;750
Lot #: XXXXXXXXX Exp: (10) XXXXXXXXX
Batch #: XXXXXXXXX (21)
Drug Status: RX
CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01) 0 0367063 04110 3
(17)
(10) XXXXXXXXX
(21)

Packaged By:
Enovachem
PHARMACEUTICALS
375 Van Ness Ave.
Suite 1403-1405
Torrance, CA 90501
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-20
Qty: 20
Distributed By: Aurobindo Pharma USA, Inc.
Source NDC: 59651-341-XX
Description: white capsule shaped, firm-coated tablet/debossed METT;750
Lot #: XXXXXXXXX Exp: (10) XXXXXXXXX
Batch #: XXXXXXXXX (21)
Drug Status: RX
CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01) 0 0367063 04120 2
(17)
(10) XXXXXXXXX
(21)

Packaged By:
Enovachem
PHARMACEUTICALS
375 Van Ness Ave.
Suite 1403-1405
Torrance, CA 90501
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-30
Qty: 30
Distributed By: Aurobindo Pharma USA, Inc.
Source NDC: 59651-341-XX
Description: white capsule shaped, firm-coated tablet/debossed METT;750
Lot #: XXXXXXXXX Exp: (10) XXXXXXXXX
Batch #: XXXXXXXXX (21)
Drug Status: RX
CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01) 0 0367063 04130 1
(17)
(10) XXXXXXXXX
(21)

Packaged By:
Enovachem
PHARMACEUTICALS
375 Van Ness Ave.
Suite 1403-1405
Torrance, CA 90501
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-60
Qty: 60
Distributed By: Aurobindo Pharma USA, Inc.
Source NDC: 59651-341-XX
Description: white capsule shaped, firm-coated tablet/debossed METT;750
Lot #: XXXXXXXXX Exp: (10) XXXXXXXXX
Batch #: XXXXXXXXX (21)
Drug Status: RX
CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01) 0 0367063 04160 8
(17)
(10) XXXXXXXXX
(21)

Packaged By:
Enovachem
PHARMACEUTICALS
375 Van Ness Ave.
Suite 1403-1405
Torrance, CA 90501
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-90
Qty: 90
Distributed By: Aurobindo Pharma USA, Inc.
Source NDC: 59651-341-XX
Description: white capsule shaped, firm-coated tablet/debossed METT;750
Lot #: XXXXXXXXX Exp: (10) XXXXXXXXX
Batch #: XXXXXXXXX (21)
Drug Status: RX
CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01) 0 0367063 04190 5
(17)
(10) XXXXXXXXX
(21)

Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-10
S/N:
Qty: 10
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-10
S/N:
Qty: 10
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-10
S/N:
Qty: 10


Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-20
S/N:
Qty: 20
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-20
S/N:
Qty: 20
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-20
S/N:
Qty: 20

Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-30
S/N:
Qty: 30
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-30
S/N:
Qty: 30
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-30
S/N:
Qty: 30

Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-60
S/N:
Qty: 60
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-60
S/N:
Qty: 60
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-60
S/N:
Qty: 60

Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-90
S/N:
Qty: 90
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-90
S/N:
Qty: 90
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-90
S/N:
Qty: 90

Relabeled By:
Enovachem
PHARMACEUTICALS
375 Van Ness Ave.
Suite 1403-1405
Torrance, CA 90501
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-01
Qty: 100
Distributed By: Aurobindo Pharma USA, Inc.
Source NDC: 59651-341-01
Description: white capsule shaped, firm-coated tablet/debossed METT;750
Lot #: XXXXXXXXX Exp: (10) XXXXXXXXX
Batch #: XXXXXXXXX (21)
Drug Status: RX
CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01) 0 0367063 04101 1
(17)
(10) XXXXXXXXX
(21)

Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-01
S/N:
Qty: 100
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-01
S/N:
Qty: 100
Methocarbamol Tablets, USP 750 mg
NDC: 87063-041-01
S/N:
Qty: 100

METHOCARBAMOL

methocarbamol tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:87063-040(NDC:59651-340)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METHOCARBAMOL (UNII: 125OD7737X) (METHOCARBAMOL - UNII:125OD7737X)	METHOCARBAMOL	500 mg

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND (Biconvex)	Size	13mm
Flavor		Imprint Code	500;METT
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:87063-040-10	10 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2025	
2	NDC:87063-040-20	20 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2025	
3	NDC:87063-040-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2025	
4	NDC:87063-040-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2025	
5	NDC:87063-040-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2025	
6	NDC:87063-040-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA213967	02/09/2023	

METHOCARBAMOL

methocarbamol tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:87063-041(NDC:59651-341)
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
METHOCARBAMOL (UNII: 125OD7737X) (METHOCARBAMOL - UNII:125OD7737X)	METHOCARBAMOL	750 mg

Inactive Ingredients	
Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	white	Score	no score
Shape	CAPSULE (Biconvex)	Size	19mm
Flavor		Imprint Code	METT;750
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:87063-041-10	10 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2025	
2	NDC:87063-041-20	20 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2025	
3	NDC:87063-041-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2025	
4	NDC:87063-041-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2025	
5	NDC:87063-041-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2025	
6	NDC:87063-041-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2025	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA213967	08/12/2020	

Labeler - Asclemed USA, Inc. (059888437)

Establishment			
Name	Address	ID/FEI	Business Operations
ASCLEMED USA INC. DBA ENOVACHEM		059888437	relabel(87063-040, 87063-041) , repack(87063-040, 87063-041)

Revised: 11/2025

Asclemed USA, Inc.