

COLD AND FLU RELIEF HBP MAXIMUM STRENGTH MULTI-SYMPTOM- acetaminophen, chlorpheniramine maleate, dextromethorphan hbr tablet, film coated

Rite Aid Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rite Aid 44-461

Active ingredients (in each tablet)

Acetaminophen 500 mg

Chlorpheniramine maleate 2 mg

Dextromethorphan HBr 15 mg

Purpose

Pain reliever/fever reducer

Antihistamine

Cough suppressant

Uses

- temporarily relieves:
 - minor aches and pains
 - sneezing
 - runny nose
 - cough
 - headache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- cough that occurs with too much phlegm (mucus)
- glaucoma
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- avoid alcoholic beverages
- may cause marked drowsiness
- alcohol, sedatives, and tranquilizers may increase drowsiness
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over: 2 tablets every 6 hours while symptoms persist. Do not take more than 6 tablets in 24 hours.
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number
- protect from excessive moisture

Inactive ingredients

corn starch, crospovidone, D&C yellow #10 aluminum lake, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal display panel

RITE

AID®

PHARMACY

*Compare to the active ingredients in Coricidin® HBP Maximum Strength Flu

MAXIMUM STRENGTH

**multi-symptom
cold and flu
relief hbp**

acetaminophen 500 mg
chlorpheniramine maleate 2 mg
dextromethorphan HBr 15 mg

pain reliever/fever reducer, antihistamine
& cough suppressant

DECONGESTANT-FREE

***cold relief for people with
high blood pressure***

relieves:

body aches, pains and headache
fever
cough
runny nose

ACTUAL SIZE

20 TABLETS

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Maximum Strength Flu.

50844 REV1016D46109

DISTRIBUTED BY: RITE AID

**30 HUNTER LANE
CAMP HILL, PA 17011**

RITE AID® SATISFACTION GUARANTEE

IF YOU'RE NOT SATISFIED. WE'LL HAPPILY REFUND YOUR MONEY.

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Maximum Strength Flu. 50844 REV1016D46109

DISTRIBUTED BY: RITE AID
30 HUNTER LANE
CAMP HILL, PA 17011

RITE AID PHARMACY

Compare to the active ingredients in Coricidin® HBP Maximum Strength Flu

MAXIMUM STRENGTH multi-symptom cold and flu relief hbp

**acetaminophen 500 mg
chlorpheniramine maleate 2 mg
dextromethorphan HBr 15 mg**

pain reliever/fever reducer, antihistamine & cough suppressant

DECONGESTANT-FREE cold relief for people with high blood pressure

relieves:
body aches, pains and headache
fever
cough
runny nose

20 TABLETS

Drug Facts (continued)
Severe liver damage may occur if you take:
■ more than 4,000 mg in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ blisters ■ rash ■ skin reddening
If a skin reaction occurs, stop use and seek medical help right away.
Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if you have
■ difficulty in urination due to enlargement of the prostate gland ■ liver disease
■ cough that occurs with too much phlegm (mucus) ■ glaucoma
■ a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema
Ask a doctor or pharmacist before use if you are
■ taking the blood thinning drug warfarin
■ taking sedatives or tranquilizers
When using this product
■ avoid alcoholic beverages

Drug Facts (continued)
Severe liver damage may occur if you take:
■ may cause marked drowsiness
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ excitability may occur, especially in children
■ use caution when driving a motor vehicle or operating machinery
Stop use and ask a doctor if
■ pain or cough gets worse or lasts more than 7 days
■ new symptoms occur
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ cough comes back or occurs with rash or headache that lasts.
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ do not take more than directed
■ adults and children 12 years and over: 2 tablets every 6 hours while symptoms persist. Do not take more than 6 tablets in 24 hours.
■ children under 12 years: ask a doctor

Other information
■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ see end flap for expiration date and lot number
■ protect from excessive moisture

Drug Facts (continued)
Uses
■ temporarily relieves:
■ minor aches and pains
■ sneezing ■ runny nose ■ cough
■ temporarily reduces fever
Warnings
Liver warning: This product contains acetaminophen.

Drug Facts (continued)
Active ingredients (in each tablet)
Acetaminophen 500 mg Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg Antihistamine
Dextromethorphan HBr 15 mg Cough suppressant

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

ITEM 0304220

0 11822 04220 8

B-1702-461-09-R
REV1016D46109

No print/No varnish area
Lot no/Exp date

Rite Aid 44-461

COLD AND FLU RELIEF HBP MAXIMUM STRENGTH MULTI-SYMPTOM

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-4610
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	

FD&C RED NO. 40 (UNII: WZB9127XOA)
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)
MAGNESIUM STEARATE (UNII: 70097M6I30)
CELLULOSE, MICROCRYSTALLINE (UNII: OPIR32D61U)
CROSPVIDONE (UNII: 2S7830E561)
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)
POVIDONE (UNII: FZ989GH94E)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
STEARIC ACID (UNII: 4ELV7Z65AP)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)

Product Characteristics

Color	RED	Score	no score
Shape	OVAL	Size	18mm
Flavor		Imprint Code	44;461
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-4610-9	2 in 1 CARTON	06/01/2005	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	06/01/2005	

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(11822-4610)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(11822-4610)