

**PERIOMED- stannous fluoride rinse**  
**Solventum US OpCo LLC**

-----  
**Drug Facts**

**Active ingredient**

Stannous fluoride 0.63% w/w (0.19% w/v fluoride ion)

**Purpose**

anticavity, antigingivitis, antihypersensitivity, mouthrinse

**Uses**

- aids in the prevention of childhood dental decay
- helps prevent gingivitis
- helps control plaque bacteria
- builds increasing protection against painful tooth sensitivity

**Warnings**

- **When using this product** do not use for sensitivity longer than four weeks unless recommended by a dentist.
- **Stop use and ask a doctor if** sensitivity persists or worsens.
- Sensitive teeth may indicate a serious problem that may need prompt care.
- **Keep out of reach of children.** If more than used for rinsing is accidentally swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

adults and children 6 years of age and older:

- do not use before mixing with water
- use immediately after preparing the rinse
- use as directed by a dental professional
- use once a day after brushing your teeth with a toothpaste or as directed by your dentist
- remove cap and seal and replace with pump
- rotate spout counter clockwise to extend
- depress pump until medicine comes to the tip of the pump
- put 1/8 oz. (approx. 2 pump strokes) of concentrated PerioMed Rinse into the provided mixing vial
- add water to 1 oz. line on mixing vial
- tightly snap on the cap and shake well

- vigorously swish rinse between your teeth for 1 minute and then spit out
- do not swallow the rinse
- do not eat or drink for 30 minutes after rinsing
- instruct children in good rinsing habits (to minimize swallowing)
- supervise children as necessary until capable of using without supervision

children under 6 years of age:

- consult a dentist or doctor

### **Other information**

- For professional use only
- products containing stannous fluoride may produce surface staining of the teeth
- adequate toothbrushing may prevent these stains which are not harmful and may be removed by your dentist
- see your dentist regularly
- this product is not intended for home or unsupervised consumer use
- for your protection, do not use if seal under the cap is broken or missing

### **Inactive ingredients**

flavor, glycerin, methyl paraben, propyl paraben, sodium saccharin

### **Questions or comments?**

**1-800-634-2249**

### **Principal Display Panel - Box Label (Mint - 283.5 g)**

NDC 48878-3315-0

Mint

**3M ESPE**

**PerioMed™**

0.63% Stannous

Fluoride Concentrate

**Antimicrobial**

**Oral Rinse**

Non-Alcohol

Formula

IMPORTANT: Read


directions carefully

before using.

# Contents:

1 Bottle

NET WT. 10 oz (283.5g)



### Drug Facts

#### Active ingredient

Stannous Fluoride 0.63% w/w (0.19% w/v fluoride ion)

#### Purpose

antigingivitis, antihyperplasia, antihypersensitivity, mouthwash

#### Uses

- aids in the prevention of childhood dental decay
- helps prevent gingivitis
- helps control plaque bacteria
- builds increasing protection against painful tooth sensitivity

#### Warnings

**When using this product** do not use for sensitivity longer than four weeks unless recommended by a dentist

**Stop use and ask a doctor if** sensitivity persists or worsens.

• sensitive teeth may indicate a serious problem that may need prompt care

**Keep out of reach of children.** If more than used for rinsing is accidentally swallowed, get medical help or contact a Poison Control Center right away.

#### Directions

• do not use before rinsing with water

• use immediately after preparing the rinse

• use as directed by a dental professional

• use once a day after brushing your teeth with a toothpaste or as directed by your dentist

• remove cap and seal and replace with pump

• rotate spout counter clockwise to extend

• depress pump until medicine comes to the tip of the pump

• put 1/8 oz. (approx. 2 pump strokes) of concentrated PerioMed Rinse into the provided mixing vial

• add water to 1 oz. line on mixing vial

• tightly snap on the cap and shake well

• vigorously swish rinse between your teeth for 1 minute and then spit out

• do not swallow the rinse

• do not eat or drink for 30 minutes after rinsing

• instruct children in good rinsing habits (to minimize swallowing)

• supervise children as necessary until capable of using without supervision

• consult a dentist or doctor

#### Other information

• For professional use only

• products containing stannous fluoride may produce surface staining of the teeth

• adequate toothbrushing may prevent these stains which are not harmful and may be removed by your dentist

• use your dentist regularly

• this product is not intended for home or unsupervised consumer use

• for your protection, do not use if seal under the cap is broken or missing

**Inactive ingredients** flavor, glycerin, methyl paraben, propyl paraben, sodium saccharin

**Questions or comments?** 1-800-634-2249

3M, ESPE and PerioMed are trademarks of 3M or 3M Deutschland GmbH, © 3M 2016. All rights reserved.

Made in U.S.A. for 3M ESPE Dental Products 2510 Conway Avenue St. Paul, MN 55144-1000 USA

12105M 3M ESPE LD, No. 70-2014-0080-4

Barcode  
30605861000018  
12105M PerioMed 0.63% SNF 2 - Mint

PerioMed™  
0.63% Stannous Fluoride Concentrate  
Antimicrobial Oral Rinse

Mint

NDC 48878-111-0

Mint

Mint

3M ESPE  
PerioMed™  
0.63% Stannous Fluoride Concentrate

Antimicrobial Oral Rinse

Non-Alcohol Formula

IMPORTANT: Read directions carefully before using.

Treatment For:  
Tooth Decay  
Gingivitis  
Sensitivity

Contents:  
1 Bottle  
NET WT. 10oz (283.5g)

Principal Display Panel - Box Label (Mint - 78 g)

NDC 48878-3315-2

Mint

**3M ESPE**

**PerioMed™**

0.63% Stannous

Fluoride Concentrate

**Antimicrobial**

**Oral Rinse**

Non-Alcohol

Formula

IMPORTANT: Read  
directions carefully  
before using.

**Contents:**

**1 Bottle**

**NET WT. 2.75 oz (78g)**

34-8719-2946-8-A

Antimicrobial Oral Rinse  
Fluoride Concentrate  
0.63% Stannous  
PerioMed™  
3M ESPE

Mint

NDC 48878-3315-2

Mint

3M, ESPE and PerioMed are trademarks of 3M or 3M Deutschland GmbH. © 3M 2016. All rights reserved.

Made in U.S.A. for  
3M ESPE  
Dental Products  
2510 Conway Avenue  
St. Paul, MN 55144-1000 USA

REF 12105SM  
3M ESPE I.D. No.  
70-2014-0270-1

3M ESPE  
PerioMed™  
0.63% Stannous  
Fluoride Concentrate

Antimicrobial  
Oral Rinse

Non-Alcohol  
Formula

IMPORTANT: Read  
directions carefully  
before using.

Treatment For:  
Tooth Decay  
Gingivitis  
Sensitivity

Contents:  
1 Bottle  
NET WT. 2.75 oz (78g)

**Drug Facts**

**Active ingredient Purpose**  
Stannous fluoride.....antivitality,  
0.63% w/w antigingivitis,  
(0.19% w/v antihypersensitivity,  
fluoride ion) mouthrinse

**Uses** • aids in the prevention of childhood dental decay • helps prevent gingivitis  
• helps control plaque bacteria • builds increasing protection against painful tooth sensitivity

**Warnings**  
• when using this product do not use for sensitivity longer than four weeks unless recommended by a dentist  
• stop use and ask a dentist if sensitivity persists or worsens  
• sensitive teeth may indicate a serious problem that may need prompt care

**Keep out of reach of children.** If more than used for rinsing is accidentally swallowed, seek medical attention or contact a Poison Control Center immediately.


**Directions**  
• Adults and children 6 years of age and older: Do not use before mixing with water. Use once daily after brushing your teeth with a toothpaste or as directed by your dentist. Pour 1/8 oz. into cup provided. Add tap water to 1 oz. line on cup and mix. Vigorously swish rinse between your teeth for 1 minute and then spit out. Do not swallow the rinse. Do not eat or drink for 30 minutes after rinsing. Instruct children under 12 years of age in good rinsing habits (to minimize swallowing). Supervise children as necessary until capable of using without supervision.  
• Children under 6 years of age: Consult a dentist or doctor.

**Other information**  
• products containing stannous fluoride may produce surface staining of the teeth  
• adequate toothbrushing may prevent these stains which are not harmful and may be removed by your dentist  
• see your dentist regularly

**Inactive ingredients** flavor, glycerin, methyl paraben, propyl paraben, sodium saccharin

**Questions or comments?**  
1-800-634-2249

30605861060210  
12105SM PerioMed 0.63% SNF 2 - Mint



Tropical Fruit

**3M ESPE**

**PerioMed™**

0.63% Stannous

Fluoride Concentrate

**Antimicrobial**

**Oral Rinse**

Non-Alcohol


Formula

IMPORTANT: Read  
directions carefully  
before using.

**Contents:**

**1 Bottle**

**NET WT. 10 oz (283.5g)**



# PERIOMED

stannous fluoride rinse

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:48878-3315
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Stannous Fluoride (UNII: 3FTR44B32Q) (Fluoride Ion - UNII:Q80VPU408O)	Fluoride Ion	1.53 mg in 1 g

## Inactive Ingredients

Ingredient Name	Strength
Glycerin (UNII: PDC6A3C00X)	
Methylparaben (UNII: A2I8C7HI9T)	
Propylparaben (UNII: Z8IX2SC1OH)	
Saccharin (UNII: FST467XS7D)	

## Product Characteristics

Color		Score	
Shape		Size	
Flavor	MINT (MINT)	Imprint Code	
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:48878-3315-0	1 in 1 BOX	07/01/1967	
1		283.5 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:48878-3315-2	1 in 1 BOX	02/01/2016	
2		85 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M021	07/01/1967	

## PERIOMED

stannous fluoride rinse

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:48878-3317
Route of Administration	ORAL		



**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>Stannous Fluoride</b> (UNII: 3FTR44B32Q) (Fluoride Ion - UNII:Q80VPU408O)	Fluoride Ion	1.53 mg in 1 g

**Inactive Ingredients**

Ingredient Name	Strength
<b>Glycerin</b> (UNII: PDC6A3C0OX)	
<b>Methylparaben</b> (UNII: A2I8C7HI9T)	
<b>Propylparaben</b> (UNII: Z8IX2SC1OH)	
<b>Saccharin</b> (UNII: FST467XS7D)	

**Product Characteristics**

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	FRUIT (FRUIT)	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:48878-3317-0	1 in 1 BOX	07/01/1967	
1		283.5 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M021	07/01/1967	

**Labeler** - Solventum US OpCo LLC (801390852)

Revised: 4/2024

Solventum US OpCo LLC