

## **PRO-SAN ACTIVATOR- pro-san activator solution IBA**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----

## **PRO-SAN SANITIZING SANITIZING PRE-MILKING AND POST-MILKING TEAT DIP A HIGHLY EFFECTIVE GERMICIDAL NON-IODINE FORMULATION CONTAINING 10% GLYCERIN**

### **ACTIVE INGREDIENT**

Lactic Acid.....2.6%

### **INERT INGREDIENTS**

Glycerin, Surfactants, Viscosity Builders and Food Grade Dyes

### **INDICATIONS AND USAGE**

#### **GENERAL RECOMMENDATIONS**

To make a ready-to-use solution of Pro-San teat dip, add equal volumes of Pro-San Activator and Pro-San Base into a clean dip cup/container and mix until the color is uniform throughout. Do not dilute. Do not mix with any other products. Mix only the amount needed for a 24 hour period. Discard any unused teat dip. Unused teat dip may be diluted with water and safely flushed down the drain.

#### **RECOMMENDED PROCEDURES FOR TEAT SANITATION**

##### **A. PRE-MILKING (PRE-DIPPING)**

1. Remove visible soils from the teats
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip or spray the cow's teats full length with READY-TO-USE PRO-SAN TEAT DIP.
4. Wait at least 15-30 seconds.
5. Remove all teat dip by thoroughly drying the teats, teat orifices and udder with a clean, single service paper towel(s) to avoid contamination of milk.
6. Attach milking unit.

##### **B. POST-MILKING (POST-DIPPING)**

Immediately after milking, dip or spray all teats full length with READY-TO-USE PRO-SAN TEAT DIP. Ensure good coverage of all sides of the teats and teat orifices. Allow to dry. Do not wipe. Do not turn cows out in freezing weather until teat dip is completely dry. r until teat dip is completely dry.

### **WARNING**

Do not dilute this product.  
Do not return teat dip left in the teat dipper to storage container.

Ensure that dipped teats are dry before exposing the animals to weather conditions which may cause damage to the teats.

Store at room temperature. Always store away from direct sunlight.

Protect this product from freezing. However, freezing will not affect PRO-SAN ACTIVATOR provided that the contents are thawed completely and agitated thoroughly before mixing base and activator together.

**CAUTION: KEEP OUT OF REACH OF CHILDREN.  
AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.**

## **FIRST AID PROCEDURES**

**EYE CONTACT:** Rinse cautiously with fresh, tepid water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing for 15 minutes. Get immediate medical advice/attention. **SKIN CONTACT:** For RTU products wash any sensitive areas. **INGESTION:** Do not induce vomiting. Never give anything by mouth to an unconscious person. Drink large amounts of water. If vomiting occurs, keep airways open. Keep head lower than the hips to prevent aspiration into the lungs. Get immediate medical advice/attention.

**ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION  
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300**

## **PRO-SAN ACTIVATOR**

NDC 29983-0099-4



# PRO-SAN



## SANITIZING PRE-MILKING AND POST-MILKING TEAT DIP A Highly Effective Germicidal Non-Iodine Formulation Containing 10% Glycerin

### ACTIVE INGREDIENT:

Lactic Acid 2.6%

### INERT INGREDIENTS:

Glycerin, Surfactants, Viscosity Builders and Food Grade Dyes

### GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

### CAUTION:

**KEEP OUT OF REACH OF CHILDREN.  
AVOID CONTACT WITH FOOD.  
DO NOT TAKE INTERNALLY.  
NOT FOR HUMAN USE.**

### FIRST AID PROCEDURES


**EYE CONTACT:** RINSE CAUTIOUSLY WITH FRESH, TEPID WATER FOR SEVERAL MINUTES. REMOVE CONTACT LENSES IF PRESENT AND EASY TO DO. CONTINUE RINSING FOR 15 MINUTES. GET IMMEDIATE MEDICAL ATTENTION.

**SKIN CONTACT:** FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

**INGESTION:** DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS, KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS. GET IMMEDIATE MEDICAL ATTENTION.

**ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION  
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300**

## ACTIVATOR FOR USE ONLY WITH PRO-SAN BASE



**MIXING DIRECTIONS:** TO MAKE A READY-TO-USE SOLUTION OF PRO-SAN TEAT DIP, ADD EQUAL VOLUMES OF PRO-SAN ACTIVATOR AND PRO-SAN BASE INTO A CLEAN DIP CUP/CONTAINER AND MIX UNTIL THE COLOR IS UNIFORM THROUGHOUT. DO NOT DILUTE. DO NOT MIX WITH ANY OTHER PRODUCTS. MIX ONLY THE AMOUNT NEEDED FOR A 24 HOUR PERIOD. DISCARD ANY UNUSED TEAT DIP. UNUSED TEAT DIP MAY BE DILUTED WITH WATER AND SAFELY FLUSHED DOWN THE DRAIN.

### RECOMMENDED PROCEDURES FOR TEAT SANITATION

#### A. PRE-MILKING (PRE-DIPPING)

1. Remove visible soots from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip or spray the cow's teats full length with **READY-TO-USE PRO-SAN TEAT DIP**.
4. Wait at least 15-30 seconds.
5. Remove all teat dip by thoroughly drying the teats, teat orifices and udder with a clean, single service paper towel(s) to avoid contamination of milk.
6. Attach milking unit.

#### B. POST-MILKING (POST-DIPPING)

Immediately after milking, dip or spray all teats full length with **READY-TO-USE PRO-SAN TEAT DIP**. Ensure good coverage of all sides of the teats and teat orifices. Allow to dry. Do not wipe. Do not turn cows out in freezing weather until teat dip is completely dry.

### WARNING

DO NOT DILUTE THIS PRODUCT.  
DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.  
ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.  
STORE AT ROOM TEMPERATURE. ALWAYS STORE AWAY FROM DIRECT SUNLIGHT.  
PROTECT THIS PRODUCT FROM FREEZING. HOWEVER, FREEZING WILL NOT AFFECT PRO-SAN PROVIDED THAT THE CONTENTS ARE THAWED COMPLETELY AND AGITATED THOROUGHLY BEFORE MIXING BASE AND ACTIVATOR TOGETHER.

## A PART OF THE COMPLETE IBA SANITATION PROGRAM

EXP. DATE:

LOT NO.:

NET CONTENTS: 5 GALLONS

IBA STOCK # 162556

122E

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 103 GILMORE DRIVE, SUTTON, MA 01590

NDC 29983-0099-4

**NPE FREE** **PRO-SAN** **NPE FREE**

**SANITIZING PRE-MILKING AND POST-MILKING TEAT DIP**  
A Highly Effective Germicidal Non-Iodine Formulation Containing 10% Glycerin

**ACTIVE INGREDIENT:**  
Lactic Acid 2.6%

**INERT INGREDIENTS:**  
Glycerin, Surfactants, Viscosity Builders and Food Grade Dyes

**GENERAL RECOMMENDATIONS**

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

**CAUTION:**  
KEEP OUT OF REACH OF CHILDREN.  
AVOID CONTACT WITH FOOD.  
DO NOT TAKE INTERNALLY.  
NOT FOR HUMAN USE.

**FIRST AID PROCEDURES**


**EYE CONTACT:** RINSE CAUTIOUSLY WITH FRESH, TEPID WATER FOR SEVERAL MINUTES. REMOVE CONTACT LENSES IF PRESENT AND EASY TO DO. CONTINUE RINSING FOR 15 MINUTES. GET IMMEDIATE MEDICAL ADVICE/ATTENTION.

**SKIN CONTACT:** FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

**INGESTION:** DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS, KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS. GET IMMEDIATE MEDICAL ADVICE/ATTENTION.

**ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION FOR MEDICAL EMERGENCIES CALL 1-800-424-9300**

**ACTIVATOR**  
FOR USE ONLY WITH  
PRO-SAN  
BASE



**MIXING DIRECTIONS:** TO MAKE A READY-TO-USE SOLUTION OF PRO-SAN TEAT DIP, ADD EQUAL VOLUMES OF PRO-SAN ACTIVATOR AND PRO-SAN BASE INTO A CLEAN DIP CUP/CONTAINER AND MIX UNTIL THE COLOR IS UNIFORM THROUGHOUT. DO NOT DILUTE. DO NOT MIX WITH ANY OTHER PRODUCTS. MIX ONLY THE AMOUNT NEEDED FOR A 24 HOUR PERIOD. DISCARD ANY UNUSED TEAT DIP. UNUSED TEAT DIP MAY BE DILUTED WITH WATER AND SAFELY FLUSHED DOWN THE DRAIN.

**RECOMMENDED PROCEDURES FOR TEAT SANITATION**

**A. PRE-MILKING (PRE-DIPPING)**

1. Remove visible soots from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip or spray the cow's teats full length with **READY-TO-USE PRO-SAN TEAT DIP**.
4. Wait at least 15-30 seconds.
5. Remove all teat dip by thoroughly drying the teats, teat orifices and udder with a clean, single service paper towel(s) to avoid contamination of milk.
6. Attach milking unit.

**B. POST-MILKING (POST-DIPPING)**  
Immediately after milking, dip or spray all teats full length with **READY-TO-USE PRO-SAN TEAT DIP**. Ensure good coverage of all sides of the teats and teat orifices. Allow to dry. Do not wipe. Do not turn cows out in freezing weather until teat dip is completely dry.

**WARNING**  
DO NOT DILUTE THIS PRODUCT.  
DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.  
ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.  
STORE AT ROOM TEMPERATURE. ALWAYS STORE AWAY FROM DIRECT SUNLIGHT.  
PROTECT THIS PRODUCT FROM FREEZING. HOWEVER, FREEZING WILL NOT AFFECT PRO-SAN PROVIDED THAT THE CONTENTS ARE THAWED COMPLETELY AND AGITATED THOROUGHLY BEFORE MIXING BASE AND ACTIVATOR TOGETHER.

**A PART OF THE COMPLETE IBA SANITATION PROGRAM**

EXP. DATE:                      LOT NO.:                      NET CONTENTS: 5 GALLONS                      IBA STOCK # 162556                      122E

**MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 103 GILMORE DRIVE, SUTTON, MA 01590**

## PRO-SAN ACTIVATOR

pro-san activator solution

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:29983-0099
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LACTIC ACID</b> (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)	LACTIC ACID	26 g in 1 L

### Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>UREA</b> (UNII: 8W8T17847W)	

## Product Characteristics

<b>Color</b>	orange	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-0099-4	18.9 L in 1 DRUM		
2	NDC:29983-0099-6	56.7 L in 1 DRUM		
3	NDC:29983-0099-8	207.9 L in 1 DRUM		
4	NDC:29983-0099-9	1039.5 L in 1 DRUM		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		05/18/2022	

**Labeler** - IBA (019494160)

Revised: 12/2024

IBA