

## **UTI PAIN RELIEF- phenazopyridine hydrochloride tablet**

### **Cadence Health**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----

### **Drug Facts**

#### **Active Ingredient (in each tablet)**

Phenazopyridine Hydrochloride 99.5 mg

#### **Purpose**

Urinary tract analgesic

#### **Uses**

Fast relief from urinary pain, burning, urgency and frequency associated with urinary tract infections. Treatment should not exceed 2 days; see Directions

#### **Warnings**

**Do not exceed recommended dosage**

#### **Do Not Use:**

if you have Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency unless approved by your physician.

#### **Ask a doctor before use if you have:**

- kidney disease
- allergies to foods, preservatives or dyes
- had a hypersensitive reaction to phenazopyridine

#### **When using this product:**

- stomach upset may occur, taking this product with or after meals may reduce stomach upset
- your urine will become reddish-orange in color. This is not harmful, but care should be taken to avoid staining clothing or other items.

#### **Stop use and ask a doctor if:**

- your symptoms last for more than 2 days

- you suspect you are having an adverse reaction to the medication
- long-term administration of phenazopyridine hydrochloride has induced neoplasia in rats (large intestine) and mice (liver). Although no association between phenazopyridine hydrochloride and human neoplasia has been reported, adequate epidemiological studies along these lines have not been conducted.

### **If pregnant or breastfeeding**

ask a health professional before use.

### **Keep out of reach of children**

In case of an overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

### ***Directions***

#### **Adults and children 12 years and over:**

- Take 2 tablets 3 times daily with a full glass of water, with or after meals as needed.
- **Do not use for more than 2 days (12 tablets) without consulting a doctor.**

#### **Children under 12 years:**

- consult a doctor.

### ***Other information***

- this product may stain contact lenses
- this product can interfere with laboratory tests including urine, glucose (sugar), and ketones tests
- store at room temperature between 15-30°C (59-86°F) in a dry place and protect from light
- Tamper evident: tablets sealed in blisters. Do not use if blister foil or seal is open and damaged

### ***Inactive Ingredients***

corn starch, croscarmellose sodium, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, silicon dioxide, sodium starch glycolate, talc, triacetin.

### **MANUFACTURED FOR**

Cadence OTC, Oakland, CA 94612

### **Questions or comments:**

Call 1-877-862-7455 or visit us at [cadenceotc.com](http://cadenceotc.com)

# PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

Cadence<sup>OTC</sup>

## UTI PAIN RELIEF

Maximum Strength, fast relief  
from UTI pain and burning

99.5 MG PHENAZOPYRIDINE HYDROCHLORIDE

2 day supply  
12 tablets

**WARNING:**This product can expose you to phenazopyridine hydrochloride, which is known to the state of California to cause cancer. For more information visit [www.P65Warnings.ca.gov](http://www.P65Warnings.ca.gov)

CSSMAXCAD Rev 12/25



**Cadence<sup>OTC</sup>**  
**UTI**  
**PAIN RELIEF**

Maximum strength, fast relief  
from UTI pain and burning

99.5 MG PHENAZOPYRIDINE HYDROCHLORIDE

2 day supply  
12 tablets

**Drug Facts**

Active ingredient (in each tablet)	Purpose
Phenazopyridine Hydrochloride 99.5 mg	Urinary tract analgesic

**Uses** Fast relief from urinary pain, burning, urgency and frequency associated with urinary tract infections. Treatment should not exceed 2 days; see Directions

**Warnings**  
Do not exceed recommended dosage  
Do not use: if you have Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency unless approved by your physician.  
Ask a doctor before use if you have: • kidney disease • allergies to foods, preservatives or dyes • had a hypersensitive reaction to phenazopyridine  
When using this product: • stomach upset may occur, taking this product with or after meals may reduce stomach upset • your urine will become reddish-orange in color. This is not harmful, but care should be taken to avoid staining clothing or other items.  
Stop use and ask a doctor if: • your symptoms last for more than 2 days • you suspect you are having an adverse reaction to the medication  
• long-term administration of phenazopyridine hydrochloride has induced neoplasia in rats (large intestine) and mice (liver). Although no association between phenazopyridine hydrochloride and human neoplasia has been reported, adequate epidemiological studies along these lines have not been conducted.  
If pregnant or breastfeeding, ask a health professional before use.  
Keep out of reach of children. In case of an overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

**Directions** Adults and children 12 years and over: • Take 2 tablets 3 times daily with a full glass of water, with or after meals as needed. • Do not use for more than 2 days (12 tablets) without consulting a doctor.  
Children under 12 years: • consult a doctor.

**Other information** • this product may stain contact lenses • this product can interfere with laboratory tests including urine, glucose (sugar), and ketones tests • store at room temperature between 15-30°C (59-86°F) in a dry place and protect from light • Tamper evident: tablets sealed in blisters. Do not use if blister foil or seal is open and damaged

**Inactive ingredients** corn starch, croscarmellose sodium, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, silicon dioxide, sodium starch glycolate, taic, triacetin.

**MANUFACTURED FOR**  
Cadence OTC, Oakland, CA 94612  
Questions or comments: Call 1-877-862-7455  
or visit us at [cadenceotc.com](http://cadenceotc.com)

**WARNING:** This product can expose you to phenazopyridine hydrochloride, which is known to the state of California to cause cancer. For more information visit [www.P65Warnings.ca.gov](http://www.P65Warnings.ca.gov)

CSSMAXCAD Rev 12/25

## UTI PAIN RELIEF

phenazopyridine hydrochloride tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:73436-2441
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>PHENAZOPYRIDINE HYDROCHLORIDE</b> (UNII: 0EWG668W17) (PHENAZOPYRIDINE - UNII:K2J09EMJ52)	PHENAZOPYRIDINE HYDROCHLORIDE	99.5 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>STARCH, CORN</b> (UNII: O8232NY35J)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>LACTOSE, UNSPECIFIED FORM</b> (UNII: J2B2A4N98G)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A</b> (UNII: H8AV0SQX4D)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	

## Product Characteristics

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	p99
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:73436-2441-2	1 in 1 CARTON	04/15/2026	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
unapproved drug other		04/06/2026	

---

**Labeler** - Cadence Health (116830263)

Revised: 4/2026

Cadence Health