

**FAMOTIDINE- famotidine tablet**  
**Chain Drug Marketing Association INC**

-----  
**Dr.Reddy's Laboratories Limited**

**Active ingredient (in each tablet)**

Famotidine USP, 20 mg

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## **Directions**

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
  - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
  - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

## **Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture and light

## **Inactive ingredients**

colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, talc and titanium dioxide

## **Questions or comments?**

call **1-888-375-3784**


## **Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Famotidine 20 mg

Container Label

NDC 83324-116-25



## Maximum Strength Famotidine Tablets USP, 20 mg Acid Reducer

**Just One Tablet!**  
Prevents & Relieves Heartburn  
Due to Acid Indigestion

**25 Tablets**

**MADE IN INDIA**  
REV 05/2024

LOT  
EXP

**150 100739**

Distributed by C.D.M.A., Inc.  
43157 W 9 Mile Rd  
Novi, MI 48375  
www.qualitychoice.com  
Questions: 800-939-5362

**AT S F A C T I O N**  
**GUARANTEED**

**TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING**

**Drug Facts**  
**Active ingredient (in each tablet)** Famotidine USP, 20 mg  
**Purpose** Acid reducer. **Uses** ■ relieves heartburn associated with acid indigestion and sour stomach  
■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages  
**Warnings** **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers  
**Do not use** ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.  
■ with other acid reducers  
**Ask a doctor before use if you have** ■ had heartburn over 2 months. This may be a sign of a more serious condition.  
■ heartburn with lightheadedness, sweating, or dizziness  
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
■ frequent chest pain  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss  
■ nausea or vomiting  
■ stomach pain  
■ kidney disease

**Drug Facts (continued)**  
**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs.  
**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).  
**Directions** ■ adults and children 12 years and over:  
■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.  
■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn  
■ do not use more than 2 tablets in 24 hours  
■ children under 12 years: ask a doctor  
**Other information** ■ read the directions and warnings before use  
■ keep the carton, it contains important information. ■ store at 20°-25°C (68°-77°F)  
■ protect from moisture  
**Inactive ingredients** colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, talc and titanium dioxide  
**Questions?** call 1-888-375-3784

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

**JUST ONE TABLET** prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods or beverages

**TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING**

\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Maximum Strength Peppid AC<sup>®</sup>. Peppid AC<sup>®</sup> is a registered trademark of Johnson & Johnson.

Famotidine 20 mg Container Carton Label

NDC 83324-116-25

Maximum Strength  
**Famotidine**  
Tablets USP, 20 mg  
Acid Reducer

1 BOTTLE INSIDE

NDC 83324-116-25

Compare to the active ingredient in Maximum Strength Peppid AC<sup>®</sup>

Maximum Strength  
**Famotidine**  
Tablets USP, 20 mg  
Acid Reducer

Just One Tablet!  
Prevents & Relieves Heartburn  
Due to Acid Indigestion

**25 Tablets**

LOT  
EXP

6 35515-95891 3

actual size

Distributed by C.D.M.A., Inc.  
43157 W 9 Mile Rd  
Novi, MI 48375  
www.qualitychoice.com  
Questions: 800-939-5362

**AT S F A C T I O N**  
**GUARANTEED**

**MADE IN INDIA** REV 06/2024

**Drug Facts**  
**Active ingredient (in each tablet)** Famotidine USP, 20 mg  
**Purpose** Acid reducer

**Uses**  
■ relieves heartburn associated with acid indigestion and sour stomach  
■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**  
**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers  
**Do not use** ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.  
■ with other acid reducers  
**Ask a doctor before use if you have** ■ had heartburn over 2 months. This may be a sign of a more serious condition.  
■ heartburn with lightheadedness, sweating, or dizziness  
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
■ frequent chest pain  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss  
■ nausea or vomiting  
■ stomach pain  
■ kidney disease

**Drug Facts (continued)**  
**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs.  
**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).  
**Directions** ■ adults and children 12 years and over:  
■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.  
■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn  
■ do not use more than 2 tablets in 24 hours  
■ children under 12 years: ask a doctor  
**Other information** ■ read the directions and warnings before use  
■ keep the carton, it contains important information. ■ store at 20°-25°C (68°-77°F)  
■ protect from moisture  
**Inactive ingredients** colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, talc and titanium dioxide  
**Questions?** call 1-888-375-3784

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

**JUST ONE TABLET** prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods or beverages

**TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING**

\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Maximum Strength Peppid AC<sup>®</sup>. Peppid AC<sup>®</sup> is a registered trademark of Johnson & Johnson.

## FAMOTIDINE

famotidine tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83324-116(NDC:55111-396)
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>FAMOTIDINE</b> (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>Polyethylene Glycol, Unspecified</b> (UNII: 3WJQ0SDW1A)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	L1
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83324-116-25	1 in 1 CARTON	11/01/2024	
1		25 in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077367	12/01/2020	

**Labeler** - Chain Drug Marketing Association INC (011920774)