

NEURORESCUEX HH333- scutellaria baicalensis root, coptis japonica root, phellodendron chinensis bark, gardenia jasminoides fruit, rheum palmatum root capsule
Panacura Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

84074-001_NeuroRescueX HH333

■ For adults, take 2 capsules, once a day, ■ before or between meals

Scutellaria Baicalensis Root (240 mg in 400 mg)
Coptis Japonica Root (240 mg in 400 mg)
Phellodendron Chinensis Bark (240 mg in 400 mg)
Gardenia Jasminoides Fruit (240 mg in 400 mg)
Rheum Palmatum Root (60 mg in 400 mg)

prevention of cerebrovascular diseases, cardiovascular disease

Do not use if you have

- galactose intolerance
- lapp lactase deficiency
- glucose-galactose malabsorption (this medicine contains lactose)

Ask a doctor before use if you are

a patient with liver dysfunction (liver enzyme value (AST, ALT) levels more than 3 times normal)

a patient who is being treated by a doctor or taking other medications.

an elderly person (in general, elderly people should take with care, such as reducing their dosage, as their physiological function has deteriorated.)

a child (the safety of children's use is not established. (Less experience in use))

- prevention of cerebrovascular diseases, cardiovascular disease
- improvement of vascular endothelial cell function
- adjunctive use to control hypertension, hyperlipidemia
- weight loss support

Read the directions and warnings before use.

KEEP THE PRODUCT INFORMATION. IT CONTAINS IMPORTANT INFORMATION.

Stop use and ask a doctor, pharmacist or health professionals if

□ There is no improvement of symptoms after taking it for a few days

□ you have following symptoms after use

■ Skin trouble: ■ rash, ■ itch, ■ bloodshot, and etc.

■ Digestive system disorder: ■ anorexia, ■stomach discomfort, ■nausea, ■vomiting, and etc.

■ Liver disorder: ■ jaundice, ■ fatigue, and etc.

■ Interstitial lung disease: ■ difficult to breathe, ■ fever, and etc. accompanied by ■ cough.


■ Urinary system disorder: ■ dysuria

■ The autonomic system disorder: ■ insomnia, ■ excessive sweating, ■ fast pulse,

■ frequent urination, ■ palpitations, ■ feeling helplessness, ■ excitement, and etc.

Keep out of reach of the children.

Corn Starch, Animal originated ingredients: Lactose Hydrate/ Magnesium Stearate



PanaCura
NeuroRescueX
(HH333)
Capsule

경희청혈단
NeuroRescueX
(HH333) Capsule

60 Capsules
10 caps. x 6 blis.

Prevention of recurrence of ischemic cerebral infarction,
prevention of cerebrovascular diseases,
cardiovascular disease

Drug Facts

Active ingredients :
Scutellaria Baicalensis Root (240 mg in 400 mg)
→ Prevention of recurrence of ischemic cerebral infarction
Coptis japonica Root (240 mg in 400 mg)
→ Prevention of recurrence of ischemic cerebral infarction
Pellodendron Chinensis Bark (240 mg in 400 mg)
→ prevention of cerebrovascular diseases, cardiovascular disease
Gardenia Jasminoides Fruit (240 mg in 400 mg)
Rheum Palmatum Root (80 mg in 400 mg)

Uses :
• prevention of cerebrovascular diseases, cardiovascular disease
• improvement of vascular endothelial cell function
• adjunctive use to control hypertension, hyperlipidemia
• weight loss support

Warnings :
Read the directions and warnings before use.
KEEP THE PRODUCT INFORMATION. IT CONTAINS IMPORTANT INFORMATION.

Do not use if you have :
• galactose intolerance
• hepatic lactase deficiency
• glucose-galactose malabsorption (this medicine contains lactose)

Ask a doctor before use if you are :
• a patient with liver dysfunction (liver enzyme value (AST, ALT) levels more than 3 times normal)
• a patient who is being treated by a doctor or taking other medications.
• an elderly person (in general, elderly people should take with care, such as reducing their dosage, as their physiological function has deteriorated.)
• a child (the safety of children's use is not established. (Less experience in use))

경희청혈단
NeuroRescueX
(HH333) Capsule

60 Capsules
10 caps. x 6 blis.

Prevention of recurrence of ischemic cerebral infarction,
prevention of cerebrovascular diseases, cardiovascular disease

Lot No. :
Exp. Date :

Drug Facts(continued)

Stop use and ask a doctor, pharmacist or health professionals if :
• There is no improvement of symptoms after taking it for a few days
• you have following symptoms after use
• Skin trouble: rash, itch, bloodshot, and etc.
• Digestive system disorder: anorexia, stomach discomfort, nausea, vomiting, and etc.
• Liver disorder: jaundice, fatigue, and etc.
• Urinary system disorder: dysuria
• The autonomic system disorder: insomnia, excessive sweating, fast pulse, frequent urination, palpitations, feeling helplessness, excitement, and etc.

Other Warnings:
do not take more than directed
if taken with other herbal medicines, be careful of overlapping of the crude medicines.
In principle, you should not to take it continuously for a long time, but if it is necessary, consult with a doctor, herb doctor, dentist, pharmacist, or other health care professionals.
If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children:
If swallowed, get medical help or contact a poison Control Center (1-800-222-1222) right away.

Directions:
• For adults, take 2 capsules, once a day, before or between meals.

Other Information:
Store in a dry and cool place at 65-77°F (20-25°C) away from the direct sunlight. To avoid misuse, do not change container.



Inactive ingredients:
Corn Starch, Magnesium Stearate, Animal originated ingredients: Lactose hydrate

Questions or Comments?:
Contact us at +82-2-561-2171 or panacura@naver.com

This product has not been evaluated by Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

Manufactured for PanaCura Inc.
32, Soyanggang-ro, Chuncheon-si, Gangwon-do 24232, Republic of Korea
Manufactured by KyungJin Pharm.co., Ltd
6 Itaeng-ro 40beon-gil, Yulmyeon, Icheon, Gyeonggi 17424, Republic of Korea

PanaCura
Made in Korea



NEURORESCUE X
HH3 33

Drug Facts

Action Inefficients:

Active ingredients:
Sourashitsa (acaculonic Root) [240 mg in 400 mg]
Coptis Japonica Root [240 mg in 400 mg]
Phellodendron Chinensis Bark [240 mg in 400 mg]

 proportion of cardiovascular disease
Gardenia Jasminoides Fruit [240 mg in 400 mg]
 disease, cardiovascular disease
Shengmai (Palmatum Root) [60 mg in 400 mg]

Ungar:

- prevention of cerebrovascular diseases, cardiovascular disease
- improvement of vascular endothelial cell function
- adjunctive use to control hypertension, hyperlipidemia
- weight loss support

Warnings

Read the directions and warnings before use.
KEEP THE PRODUCT INFORMATION. IT CONTAINS
IMPORTANT INFORMATION.

Do not use if you have:

Do not use you have:

- galactose intolerance
- lappactase deficiency
- glucose-galactose-malabsorption (this medicine contains lactose)

Ask a doctor before use if you are :

a patient with liver dysfunction (liver enzyme value (AST, ALT) levels more than 3 times normal)
a patient who is being treated by a doctor or taking other medications.
an elderly person (in general, elderly people should take with care, such as reducing their dosage, as their physiological function has deteriorated)
a child (the safety of children's use is not established. Less experience in use)

Drug Facts (continued)

Stop use and ask a doctor, pharmacist or health professional if:

- There is no improvement of symptoms after taking it for 4 days
- You have difficulty swallowing or other oral problems
- You have a rash, hives, itching, or swelling of the face
- You experience dizziness, headache, blurred vision, or ringing in the ears
- You experience stomach pain, nausea, stomach discomfort, indigestion, vomiting, or stools that are black or bloody
- You experience fatigue and weakness
- You experience difficulty breathing, fever, and/or accompanying cough
- You experience urinary system disorder or flu-like symptoms
- You experience system disorder, insomnia, excessive sweating, fast pulse, frequent urination, constipation, feeling helpless, numbness, and/or dizziness

Other Warnings:
Do not take more

do not take more than directed.
If taken with other herbal medicines, be careful of overlapping of the same medicines.
In principle, you should not take it continuously for a long time, but if it is necessary, consult with a doctor, herb doctor, dentist, pharmacist, or other health care professionals.
If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a poison control center (1-800-222-1222) right away.

Directions :

For adults, take 2 capsules, once a day, before or between meals.

Other information :

Store in a dry and cool place at 68-77°F (20-25°C) away from the direct sunlight. To avoid misuse, do not change container.

Inertive Ingredients:

Corn Starch, Animal originated Ingredients,
 Lactose Hydrolyzate/Magnesium Stearate

Questions or Comments

Contact us at 800-2-667-2772

This product has not been evaluated by Food and Drug Administration.
This product is not intended to diagnose, cure, treat, or prevent any disease.

This product is not intended to diagnose, treat, cure, or prevent any disease.


Seller by Panacura Inc.
IT-109689-00-CT/Church of Christ - do NOT call 800 or www.panic.com

Manufactured by PyungDa Pharma Co., Ltd.

Elisavinda Adeniran-gb, Yabiyew, Ibadan, Oyo State, Nigeria

 ParasCura Media in Korea

67-30-104



NEURORESCUEX HH333

scutellaria baicalensis root, coptis japonica root, phellodendron chinensis bark, gardenia jasminoides fruit, rheum palmatum root capsule

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84074-001
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SCUTELLARIA BAICALENSIS ROOT (UNII: 7J95K7ID2S) (SCUTELLARIA BAICALENSIS ROOT - UNII:7J95K7ID2S)	SCUTELLARIA BAICALENSIS ROOT	240 mg
COPTIS JAPONICA ROOT (UNII: JH0SWT16D1) (COPTIS JAPONICA ROOT - UNII:JH0SWT16D1)	COPTIS JAPONICA ROOT	240 mg
PHELLODENDRON CHINENSIS BARK (UNII: 2866QMZ434) (PHELLODENDRON CHINENSIS BARK - UNII:2866QMZ434)	PHELLODENDRON CHINENSIS BARK	240 mg
GARDENIA JASMINOIDES FRUIT (UNII: 7CTH8MD549) (GARDENIA JASMINOIDES FRUIT - UNII:7CTH8MD549)	GARDENIA JASMINOIDES FRUIT	240 mg
RHEUM PALMATUM ROOT (UNII: G025DAL7CE) (RHEUM PALMATUM ROOT - UNII:G025DAL7CE)	RHEUM PALMATUM ROOT	60 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	

Product Characteristics

Color	yellow	Score	no score
Shape	CAPSULE	Size	21mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84074-001-03	2 in 1 CARTON	12/08/2025	
1	NDC:84074-001-02	6 in 1 CARTON		
1	NDC:84074-001-01	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:84074-001-05	3 in 1 CARTON	12/08/2025	
2	NDC:84074-001-04	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		05/05/2024	

Labeler - Panacura Inc. (695390655)

Registrant - Panacura Inc. (695390655)

Establishment

Name	Address	ID/FEI	Business Operations
Kyungjin Pharm.co.,Ltd		695542024	manufacture(84074-001)