

**WALGREEN CHILDRENS COLD PLUS COUGH PLUS SORE THROAT-
acetaminophen and dextromethorphan hydrobromide suspension
WALGREENS CO.**

Walgreens Children's Cold + Cough + Sore Throat Relief

Drug Facts

**Active ingredients (in each 5 Purposes
mL)**

Acetaminophen 160 mg	Pain reliever/fever reducer
Dextromethorphan HBr 5 mg	Cough suppressant

Uses

- temporarily relieves the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - cough
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert:acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning:if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain

drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

- liver disease
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product

- **do not exceed recommended dose (see overdose warning)**

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours, while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)
under 36	under 4 years	do not use
36-47	4 to 5 years	5 mL
48-95	6 to 11 years	10 mL

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- **each 5 mL contains:** sodium 3 mg
- store at room temperature

Inactive ingredients

carboxymethylcellulose sodium, anhydrous citric acid, D&C red no. 33, disodium edetate, FD&C red no. 40, flavors, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-866-467-2748

PRINCIPAL DISPLAY PANEL

Compare to the active ingredients in Children's Tylenol® Cold + Cough + Sore Throat*

NDC 0363-6504-04

Children's

Cold + Cough

+ Sore Throat Relief

Acetaminophen, Pain Reliever-Fever Reducer

Dextromethorphan HBr, Cough Suppressant

Oral Suspension

Ages 4-11 Years

- **Fever**
- **Sore Throat**
- **Cough**

Ibuprofen free, aspirin free,

No added alcohol

Bubblegum Flavor

Naturally and Artificially Flavored

4 FL OZ (120 mL)

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc, distributor of Children's Tylenol® Cold + Cough + Sore Throat.

TAMPER-EVIDENT: DO NOT USE IF PRINTED INNER SEAL UNDER CAPS IS TORN

OR MISSING.



WALGREEN CHILDRENS COLD PLUS COUGH PLUS SORE THROAT
 acetaminophen and dextromethorphan hydrobromide suspension

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-6504
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
WATER (UNII: 059QF0KOOR)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	pink	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6504-04	1 in 1 CARTON	02/24/2023	
1		120 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/24/2023	

Labeler - WALGREENS CO. (008965063)