

CAREALL ACETAMINOPHEN EXTRA STRENGTH- acetaminophen tablet
New World Imports, Inc

Drug Facts

Acetaminophen 500 mg

Pain reliever / Fever reducer

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs of symptoms.

Temporary relieves minor aches and pains due to: the common cold, headache, toothache, muscular aches, backache, minor pains of arthritis, and menstrual and premenstrual cramps

Temporarily reduces fever.

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include skin reddening, blisters, rash.

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if:

- Pain gets worse or lasts for more than 10 days
- Fever gets worse or lasts for more than 3 days
- New symptoms occur
- Redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Do not take more than directed (see overdose warnings)

Adults and children 12 years of age and older: Take 2 caplets every 4 to 6 hours while symptoms last.


Do not take more than 6 caplets in 24 hours, unless directed by a doctor.

Do not use more than 10 days unless directed by a doctor.

Children under 12 years of age: ask a doctor


povidone, pregelatinized starch, sodium starch glycolate, stearic acid

NDC 51824-101-01
DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING



For Adults
Acetaminophen
Pain Reliever / Fever Reducer

†Compare to the active ingredient of Extra Strength Tylenol®



100 CAPLETS
500 MG EACH

Drug Facts

Active ingredient (in each caplet) Purpose
Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - minor pain of arthritis
 - backache
 - toothache
 - muscular aches
 - menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- rash
- blisters


If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

PEEL BACK HERE FOR MORE DRUG FACTS

Distributed by: CareALL Products
160 Athens Way
Nashville, TN 37228
www.careallproducts.com



0 85317 00486 6

†This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark Tylenol®.

STOP PEELING

Drug Facts (continued)

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

<ul style="list-style-type: none"> ■ do not take more than directed (see overdose warning) 	<ul style="list-style-type: none"> ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor
-----------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

adults and children 12 years and over

children under 12 years

- ask a doctor

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients povidone, pregelatinized starch, sodium starch glycolate, stearic acid

Questions or comments?
Call 1-844-705-4384 Mon - Fri 8:30 AM - 5 PM EST

CAREALL ACETAMINOPHEN EXTRA STRENGTH

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51824-101
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Acetaminophen (UNII: 36209ITL9D) (Acetaminophen - UNII:36209ITL9D)	Acetaminophen	500 mg

Inactive Ingredients

Ingredient Name	Strength
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
Stearic Acid (UNII: 4ELV7Z65AP)	
Povidone (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white (White)	Score	no score
Shape	OVAL (Oval)	Size	17mm
Flavor		Imprint Code	M2A4;57344
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51824-101-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/23/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/23/2024	

Labeler - New World Imports, Inc (075372276)

Revised: 8/2024

New World Imports, Inc