

**MOTION SICKNESS RELIEF- meclizine hcl tablet  
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredient (in each tablet)**

Meclizine HCl 25 mg

**Purpose**

Antiemetic

**Uses**

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.

**Warnings**

**Do not use**

for children under 12 years of age unless directed by a doctor.

**Ask a doctor before use if you have**

- breathing problems such as emphysema or chronic bronchitis
- glaucoma
- trouble urination due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers.

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- to prevent motion sickness, the first dose should be taken 1/2 to 1 hour before starting activity.
- adults and children 12 years of age and over: take 1 to 2 tablets once daily or as directed by a doctor

- do not exceed 2 tablets in 24 hours

**Other information**

- store between 20-25°C (68-77°F)

**Inactive ingredients**

anhydrous lactose, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminium lake, magnesium stearate, microcrystalline cellulose, sodium starch glycolate

**Principal Display Panel****Compare to the active ingredient of Dramamine® Less Drowsy Formula\***

Less Drowsy Formula

Motion Sickness Relief

Meclizine HCl 25 mg / Antiemetic

Long Lasting

- Up to 24 hours of protection
- Prevents nausea, dizziness & vomiting

25 mg

Tablets

\*This product is not manufactured or distributed by Prestige Brands, Inc., distributor of Dramamine® Less Drowsy Formula

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGN OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

DISTRIBUTED BY DOLGENCORP, LLC

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

**Product Label**



Compare to the active ingredient of Dramamine® Less Drowsy Formula\*

Less Drowsy Formula

# Motion Sickness Relief

Meclizine HCl 25 mg/Antiemetic  
Long Lasting

- Up to 24 hours of protection
- Prevents nausea, dizziness & vomiting



**25**  
mg each



**8 Tablets**

Actual Tablet Size

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot No.:  
Exp. Date:

\*This product is not manufactured or distributed by Prestige Brands, Inc., distributor of Dramamine® Less Drowsy Formula.

DISTRIBUTED BY DOLGENCORP, LLC  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072

PLD-D50J  
FC002774

**100%**  
Satisfaction  
Guaranteed!  
(888) 309-9030



<b>Purpose</b>	Medicine HCl 25 mg.....Antiemetic
<b>Uses</b>	For the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.
<b>Warnings</b>	Do not use for children under 12 years of age unless directed by a doctor. Ask a doctor before use if you have ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ trouble urinating due to an enlarged prostate gland Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. When using this product ■ avoid alcoholic drinks ■ drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
<b>Directions</b>	■ to prevent motion sickness, the first dose should be taken 1/2 to 1 hour before starting activity ■ adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor ■ do not exceed 2 tablets in 24 hours
<b>Other information</b>	■ store between 20-25°C (68-77°F)
<b>Inactive ingredients</b>	stearate, microcrystalline cellulose, sodium starch glycolate anhydrous lactose, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, magnesium
<b>Questions or comments?</b>	Call 1-888-309-9030

## DOLLAR GENERAL Motion Sickness Relief

**MOTION SICKNESS RELIEF**

meclizine hcl tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:559 10 -620
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDRO CHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

**Inactive Ingredients**

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
ALUMINUM OXIDE (UNII: LM26O6933)	

**Product Characteristics**

<b>Color</b>	YELLOW	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	TCL086
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:559 10 -620-08	1 in 1 CARTON	10/03/2012	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part336	10/03/2012	

Revised: 12/2019

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)