

TUSSIN NIGHTTIME COUGH- dextromethorphan hydrobromide, doxylamine succinate solution
Rite Aid Corporation

Rite Aid Corporation TUSSIN NIGHTTIME COUGH Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr, USP 30 mg

Doxylamine succinate, USP 12.5 mg

Purposes

Cough suppressant

Antihistamine

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- controls the impulse to cough to help you sleep

Warnings

Do not use

- to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- do not take more than 4 doses in any 24-hour period
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 6 hours
children under 12 years	do not use

Other information

- **each 20 mL contains:** sodium 11 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, benzoic acid, benzyl alcohol, carboxymethylcellulose sodium,

FD&C blue #1, FD&C red #40, flavor, glycerin, menthol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to the active ingredients in Robitussin[®] Maximum Strength Nighttime Cough DM

FREE FROM

GLUTEN FREE

ALCOHOL FREE

MAXIMUM STRENGTH

TUSSIN NIGHTTIME COUGH

COUGH SUPPRESSANT

DEXTROMETHORPHAN HBr

ANTIHISTAMINE

DOXYLAMINE SUCCINATE

ADULT

Relieves cough, itchy throat, runny nose

For ages 12 & over

DM MAX

BERRY MENTHOL FLAVOR

4 FL OZ (118 mL)



TUSSIN NIGHTTIME COUGH

dextromethorphan hydrobromide, doxylamine succinate solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-6005
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 20 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
BENZOIC ACID (UNII: 8SKN0B0MIM)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	RED	Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-6005-0	1 in 1 CARTON	03/16/2023	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/16/2023	

Revised: 11/2024

Rite Aid Corporation