

**NICOTINE MINI- nicotine polacrilex lozenge**  
**HC Jones, Inc.**

-----  
**Drug Facts**

**Active ingredient (in each lozenge)**

Nicotine polacrilex USP, 4 mg

**Purpose**

Stop smoking aid

**Use**

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**

**If you are pregnant or breast-feeding,**

**only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

## Keep out of reach of children and pets.

Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### Directions (4 mg lozenge)

- **if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine lozenge
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

- **nicotine lozenge is a medicine and must be used a certain way to get the best results**
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. **Do not chew or swallow lozenge.**
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- **do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.**
- It is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

### Other information

- each lozenge contain: sodium, 5 mg
- store at 20° to 25°C (68° to 77°F)
- keep bottle tightly closed and protect from light

### Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, cherry flavor, hydroxypropyl cellulose, magnesium stearate, mannitol, potassium bicarbonate, povidone, sodium alginate, sodium carbonate, sucralose, xanthan gum

### ***Questions or comments?***

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### **Principal Display Panel (4 mg lozenge)**

\*COMPARE TO THE ACTIVE INGREDIENT IN NICORETTE® MINI LOZENGE

### **MINI NICOTINE LOZENGES**

NICOTINE POLACRILEX 4 mg

### **STOP SMOKING AID**

Cherry flavor

### **FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.**

If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use the 2 mg Nicotine Polacrilex Lozenge.

### **TO INCREASE YOUR SUCCESS IN QUITTING:**

- 1.** You must be motivated to quit.
- 2. Use Enough** - Use **at least 9** Nicotine mini lozenges per day during the first six weeks.
- 3. Use Long Enough** - Use Nicotine mini lozenges for the full 12 weeks.
- 4. Use With a Support Program** as directed in the enclosed User's Guide.

Gluten Free

Retain outer carton for full product uses, directions and warnings.

- **not for sale to those under 18 years of age**
- **proof of age required**
- **not for sale in vending machines or from any source where proof of age cannot be verified**

\*This product is not affiliated with, manufactured by or produced by the makers or owners of Nicorette® Lozenge.

**TAMPER EVIDENT FEATURE: DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.**

**RETAIN OUTER CARTON FOR FULL PRODUCT USES, DIRECTIONS AND WARNINGS.**

**Distributed by: HC Jones, Inc.**

**424 Broadway Suite 401, New York 10013**

Package Label (4 mg lozenge)

Drug Facts

Active ingredient (in each lozenge)  
nicotine polacrilex USP, 4 mg

Purpose  
Stop smoking aid

Use

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breastfeeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

If you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.

before using this product, read the enclosed User's Guide for complete directions and other important information

begin using the lozenge on your quit day

If you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge

If you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

nicotine lozenge is a medicine and must be used a certain way to get the best results

place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge.

you may feel a warm or tingling sensation

Drug Facts (continued)

occasionally move the lozenge from one side of your mouth to the other until completely dissolved

do not eat or drink 15 minutes before using or while the lozenge is in your mouth

to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks

do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects

do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.

it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

Other information

each lozenge contains: sodium, 5 mg

store at 20° to 25° C (68° to 77°)

keep bottle tightly closed and protect from light

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, cherry flavor, hydroxypropyl cellulose, magnesium stearate, mannitol, potassium bicarbonate, povidone, sodium alginate, sodium carbonate, sucralose, xanthan gum

Questions?

Call 1-877-733-3835 Monday-Friday 9AM-5PM EST

\*This product is not affiliated with, manufactured by, or produced by the makers or owners of Nicorette® Lozenge.

not for sale to those under 18 years of age

proof of age required

not for sale in vending machines or from any source where proof of age cannot be verified

JONES

MINI NICOTINE LOZENGES

NICOTINE POLACRILEX 4 MG STOP SMOKING AID

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette **more than 30 minutes** after waking up, use nicotine polacrilex 2 mg lozenge.

4 MG

81 LOZENGES  
4 mg each (3 bottles of 27)

CHERRY FLAVOR

ACTUAL SIZE

\*COMPARE TO THE ACTIVE INGREDIENT IN NICORETTE® MINI LOZENGE

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use Enough - Use at least 9 nicotine mini lozenges per day during the first six weeks.
3. Use Long Enough - Use nicotine mini lozenges for the full 12 weeks.
4. Use With a Support Program as directed in the enclosed User's Guide.

SCAN FOR JONES APP

Distributed by: J.C. Jones, Inc.  
424 Broadway Suite 401, New York, NY 10013

See inside for more details or visit [www.quitwithjones.com](http://www.quitwithjones.com)

TAMPER EVIDENT FEATURE: DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

RETAIN OUTER CARTON FOR FULL PRODUCT USES, DIRECTIONS AND WARNINGS.

PLD-A696A F0X00368

3 59726 97981 8

Lot No:

Exp. Date:

JONES Mini Nicotine Lozenges Cherry Flavor

NICOTINE MINI

nicotine polacrilex lozenge

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83253-979
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name		Basis of Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)		NICOTINE
		4 mg

Inactive Ingredients	
Ingredient Name	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
MANNITOL (UNII: 3OWL53L36A)	

<b>SODIUM ALGINATE</b> (UNII: C269C4G2ZQ)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POTASSIUM BICARBONATE</b> (UNII: HM5Z15LEBN)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>SODIUM CARBONATE</b> (UNII: 45P3261C7T)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Product Characteristics

<b>Color</b>	white (Off white)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	10mm
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	A415
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83253-979-81	3 in 1 BOX	03/31/2025	
1		27 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:83253-979-27	27 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2025	
3	NDC:83253-979-72	1 in 1 BOX	03/31/2025	
3		27 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210711	03/31/2025	

**Labeler** - HC Jones, Inc. (118845364)

Revised: 7/2025

HC Jones, Inc.