

**THERAFLU FLU RELIEF MAX STRENGTH DAYTIME NIGHTTIME COMBO PACK-
acetaminophen, dextromethorphan hbr, and acetaminophen,
chlorpheniramine maleate, dextromethorphan hbr
Haleon US Holdings LLC**

Drug Facts

Theraflu Flu Relief Max Strength* Daytime Powder

Active ingredients (in each packet)

Acetaminophen 1000 mg

Dextromethorphan HBr 30 mg

Purposes

Pain reliever/Fever reducer

Cough suppressant

Uses

- temporarily relieves these symptoms due to a common cold or flu:
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation
 - minor sore throat pain
- temporarily reduces fever

Warnings

Liver warning:This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert:Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning:If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson’s disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- take every 6 hours, while symptoms persist. Do not take more than 3 packets in 24 hours unless directed by a doctor.

1. Age	1. Dose
1. adults and children 12 years of age and over	1. one packet
1. children under 12 years of age	1. do not use

-
- dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
 - if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

- **each packet contains:**potassium 10 mg, sodium 20 mg
- **phenylketonurics:**contains phenylalanine 19.6 mg per packet
- store at a controlled room temperature at 20 - 25°C (68-77°F)

Inactive ingredients

acesulfame potassium, anhydrous citric acid, aspartame, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, maltodextrin, natural and artificial flavors, silicon dioxide, sodium citrate, soy lecithin, sucrose, tribasic calcium phosphate

Questions or comments?

1-855-297-3031

Theraflu Flu Relief Max Strength* Nighttime Powder

Drug Facts

Active ingredients (in each packet)

Acetaminophen 1000 mg

Chlorpheniramine maleate 4 mg

Dextromethorphan HBr 30 mg

Purposes

Pain reliever/Fever reducer

Antihistamine

Cough suppressant

Uses

- temporarily relieves these symptoms due to a common cold or flu:
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation

- minor sore throat pain
- runny nose
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- avoid alcoholic drinks
- marked drowsiness may occur

- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- take every 6 hours, while symptoms persist. Do not take more than 3 packets in 24 hours unless directed by a doctor.

1. Age	1. Dose
1. adults and children 12 years of age and over	1. one packet
1. children under 12 years of age	1. do not use

- dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating.
Do not overheat

Other information

- **each packet contains:**potassium 5 mg, sodium 22 mg
- **phenylketonurics:**contains phenylalanine 12.9 mg per packet
- store at a controlled room temperature at 20 - 25°C (68-77°F)

Inactive ingredients

acesulfame potassium, anhydrous citric acid, aspartame, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, maltodextrin, natural and artificial flavors, silicon dioxide, sodium citrate, soy lecithin, sucrose, tribasic calcium phosphate

Questions or comments?

1-855-297-3031

Additional Information

**READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE.
KEEP CARTON FOR REFERENCE. DO NOT DISCARD.**

**DO NOT TAKE THE DAYTIME AND NIGHTTIME PRODUCTS AT THE SAME TIME.
DO NOT TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.**

**DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 6 HOURS
AFTER THE LAST DOSE OF THE DAYTIME PRODUCT UNLESS DIRECTED BY
YOUR DOCTOR.**

**DO NOT TAKE BOTH PRODUCTS AT THE SAME TIME
OR TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.**

PARENTS: Learn about teen medicine abuse

www.StopMedicineAbuse.org

***Maximum Strength per 6 hour dose**

**TAMPER-EVIDENT INNER UNIT. DO NOT USE IF SEALED THERAFLU PACKET IS
TORN OR BROKEN.**

1-855-297-3031

Distributed by: Haleon, Warren, NJ 07059

Trademarks are owned by or licensed to the Haleon group of companies.

©2025 Haleon group of companies or its licensor.

Principal Display Panel

MULTI-SYMP TOM FLU RELIEF

HALEON

THERAFLU

FLU RELIEFMAX STRENGTH*

6 x DAYTIME

Acetaminophen Pain Reliever/Fever Reducer

Dextromethorphan HBr Cough Suppressant

6 x NIGHTTIME

AcetaminophenPain Reliever/Fever Reducer

Chlorpheniramine MaleateAntihistamine

Dextromethorphan HBrCough Suppressant

Hot liquid therapythat relieves:

/ Fever

/ Body ache

/ Headache

/ Sore throat pain

/ Cough

/ Runny nose (Nighttime only)

6 DAYTIME PACKETS

6 NIGHTTIME PACKETS

12 TOTAL PACKETS

Honey Lemon Flavor

USE ONLY AS DIRECTED

62000000214400 Front Carton

MULTI-SYMPTOM FLU RELIEF

HALEON

THERAFLU

**FLU RELIEF
MAX STRENGTH***

6 x DAYTIME

Acetaminophen Pain Reliever / Fever Reducer
Dextromethorphan HBr Cough Suppressant

6 x NIGHTTIME

Acetaminophen Pain Reliever / Fever Reducer
Chlorpheniramine Maleate Antihistamine
Dextromethorphan HBr Cough Suppressant

Hot liquid therapy that relieves:

- / Fever / Body ache / Headache
- / Sore throat pain / Cough
- / Runny nose (Nighttime only)

**6 DAYTIME PACKETS
6 NIGHTTIME PACKETS
12 TOTAL PACKETS**

**Honey Lemon
Flavor**

USE ONLY AS DIRECTED



THERAFLU FLU RELIEF MAX STRENGTH DAYTIME NIGHTTIME COMBO PACK

acetaminophen, dextromethorphan hbr, and acetaminophen, chlorpheniramine maleate,
dextromethorphan hbr kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0067-7923
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0067-7923-02	1 in 1 CARTON; Type 1: Convenience Kit of Co-Package	06/15/2022	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 CARTON	6
Part 2	1 CARTON	6

Part 1 of 2

THERAFLU FLU RELIEF MAX STRENGTH DAYTIME

acetaminophen, dextromethorphan hbr powder

Product Information

Item Code (Source)	NDC:0067-7921
---------------------------	---------------

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	1000 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
ASPARTAME (UNII: Z0H242BBR1)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	

SOYBEAN LECITHIN (UNII: 1DI56QDM62)

SUCROSE (UNII: C151H8M554)

TRIBASIC CALCIUM PHOSPHATE (UNII: 91D9GV0Z28)

Product Characteristics

Color	white (to off white, yellow, beige, and brown color)	Score	
Shape		Size	
Flavor	HONEY (Lemon)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0067-7921-02	6 in 1 CARTON; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/15/2022	

Part 2 of 2

THERAFLU FLU RELIEF MAX STRENGTH NIGHTTIME

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr powder

Product Information

Item Code (Source)	NDC:0067-7922
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	1000 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	

ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
ASPARTAME (UNII: Z0H242BBR1)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SOYBEAN LECITHIN (UNII: 1DI56QDM62)	
SUCROSE (UNII: C151H8M554)	
TRIBASIC CALCIUM PHOSPHATE (UNII: 91D9GV0Z28)	

Product Characteristics

Color	white (white to off-white, yellow, beige, and brown color)	Score	
Shape		Size	
Flavor	HONEY (lemon)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0067-7922-02	6 in 1 CARTON; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/15/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/15/2022	

Labeler - Haleon US Holdings LLC (079944263)