

TOPCARE ASPIRIN- aspirin tablet, chewable
Topco Associates LLC

Topco Associates LLC. Aspirin 81 mg Drug Facts

Active ingredient (in each chewable tablet)

Aspirin 81 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, D&C red #27 aluminum lake, dextrose excipient, FD&C red #40 aluminum lake, flavors, saccharin sodium

Questions or comments?

1-888-423-0139

Principal Display Panel

TopCare® health

COMPARE TO BAYER® CHEWABLE ASPIRIN ACTIVE INGREDIENT

LOW DOSE

ASPIRIN REGIMEN

Talk to your doctor or other healthcare provider before using this product for your heart.


Aspirin 81 mg

Chewable • PAIN RELIEVER (NSAID)

36 TABLETS

actual size

CHERRY FLAVOR



Drug Facts

Active ingredient (in each chewable tablet)
Aspirin 81 mg (NSAID)*

Purpose
Pain reliever
*nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

27468 88 C-10



DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts (continued)

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer for at least 7 days after fentanyl only or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 3 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present

Drug Facts (continued)

■ new symptoms occur

■ ringing in the ears or a loss of hearing occurs

■ these could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

■ store at 20-25°C (68-77°F)

Inactive ingredients contain starch, D&C red #27 aluminum lake, deacrylose excipient, FD&C red #40 aluminum lake, flavors, saccharin sodium

Questions or comments?
1-888-423-0139



GLUTEN FREE

Scan here for more information or call 1-888-423-0139



© TOPCO PERMUTAS RESTAURANTS LLC
ELK GROVE VILLAGE, IL 60107
www.bpc.aebhand.com

COMPARE TO BAYER® CHEWABLE ASPIRIN ACTIVE INGREDIENT



LOW DOSE ASPIRIN REGIMEN

Talk to your doctor or other healthcare provider before using this product for your heart.

Aspirin 81 mg


Chewable • PAIN RELIEVER (NSAID)

36 TABLETS

actual size



CHERRY FLAVOR



NDC 36800-274-68

TOPCARE ASPIRIN

aspirin tablet, chewable

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:36800-274 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | ASPIRIN | 81 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX) | |
| FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T) | |

Product Characteristics

| | | | |
|-----------------|--------|---------------------|----------|
| Color | PINK | Score | no score |
| Shape | ROUND | Size | 8mm |
| Flavor | CHERRY | Imprint Code | L274 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:36800-274-68 | 1 in 1 CARTON | 03/29/2005 | |
| 1 | | 36 in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M013 | 03/29/2005 | |

Labeler - Topco Associates LLC (006935977)

Revised: 8/2024

Topco Associates LLC