

NIGHT TIME MULTI-SYMPTOM COLD AND FLU RELIEF- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled
Discount Drug Mart

DISCOUNT drug mart Night Time Multi-Symptom Cold & Flu Relief

Drug Facts

Active Ingredients (In each softgel)

Acetaminophen 325 mg
Dextromethorphan HBr 15 mg
Doxylamine succinate 6.25 mg

Purpose

Acetaminophen	Pain reliever/Fever reducer
Dextromethorphan HBr	Cough suppressant
Doxylamine succinate	Antihistamine

UsesTemporarily relieves common cold and flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- minor aches and pain
- fever
- runny nose and sneezing

Warnings

Liver warning:This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 8 Liquid Caps in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert:Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- to make a child sleep

Ask a doctor before use if you have • liver disease • glaucoma

- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product, do not use more than directed.

- excitability may occur, especially in children,
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present • new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast feeding,ask a health professional before use.

Keep out of reach of children.In case of accidental overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

Directions

- take only as directed (see Warning)
- do not exceed 4 doses per 24 hours

Adults and Children 12 yrs & over	2 softgels with water every 6 hours
Children 4 to under 12 yrs	ask a doctor
Children under 4 yrs	do not use

• when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- store at room temperature 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredientsD&C Yellow # 10, FD&C Blue # 1, Gelatin, glycerin, Polyethylene glycol, Povidone, Propylene Glycol, Purified water, Sorbitol sorbitan solution, Titanium dioxide.

Questions? Call Weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

Package Labeling: DISCOUNT drug mart Night Time Multi-Symptom Cold & Flu Relief

12
1201240
3/22

Parents:
Learn about teen medicine abuse
www.stopmedicabuse.org

DISCOUNT drug mart FOOD MARKET
AN EMPLOYEE OWNERSHIP COMPANY

Cold & Flu Relief MULTI-SYMPTOM

NightTime

NDC-53943-603-03

DISCOUNT drug mart FOOD MARKET
AN EMPLOYEE OWNERSHIP COMPANY

Cold & Flu Relief MULTI-SYMPTOM

NightTime

Acetaminophen, Dextromethorphan HBr,
Doxylamine succinate
Headache, Fever, Sore Throat, Minor Aches & Pains
Sneezing, Runny Nose
Cough

Compare to the active ingredients in Vicks® NyQuil™ Cold & Flu LiquidCaps™

24 Softgels

MAY CONTAIN ANTI-THEFT DEVICE

DISTRIBUTED BY:
DRUG MART – FOOD FARE
211 COMMERCE DRIVE
MEDINA, OH 44256
www.discount-drugmart.com

SUBSTITUTION GUARANTEE
IF DISGATED, RETURN UNLACED PASTOR AND PACKAGE TO STORE
WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REQUEST
FOR DISGATION FACTURE, NAME, ADDRESS AND EMPTY PACKAGE TO:
DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

BK-131

Drug Facts	
Active ingredients (in each softgel)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine
Uses	temporarily relieves common cold/flu symptoms: cough due to minor throat & bronchial irritation sore throat
Warnings	<ul style="list-style-type: none"> liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: <ul style="list-style-type: none"> more than 8 Liquid Caps in 24 hrs, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks daily while using this product Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> skin reddening blisters rash
Other information	<ul style="list-style-type: none"> runny nose & sneezing

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product, do not use more than directed.

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives & tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

Drug Facts (continued)

- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.
In case of accidental overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children, even if you do not notice any signs or symptoms.

Directions

- take only as directed (see Warnings)
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgel with water every 6 hours
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- Store at room temperature 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients D&C Yellow # 10, FD&C Blue # 1, Gelatin, Glycerin, Polyethylene glycol, Povidone, Propylene glycol, Purified water, Sorbitol sorbitan solution, Titanium dioxide

Questions? Call weekdays from 9:30 a.m. to 4:30 p.m. EST at 1-877-798-5944

*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Vicks® NyQuil™ LiquidCaps™.

drug mart

FOOD MARKET

**Cold &
Flu Relief**

MULTI-SYMPTOM

NIGHT TIME

***Compare to the active ingredients in Vicks [®] NyQuil[™] Cold & Flu LiquiCaps[™]**

24 Softgels

DISTRIBUTED BY:

DRUG MART - FOOD FAIR

211 COMMERCE DRIVE

MEDINA, OH 44256

www.discount-drugmart.com

This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Vicks [®] DayQuil [™] LiquiCaps [™]

TAMPER EVIDENT: DO NOT USE IF

PACKAGE IS OPENED OR BLISTER

UNIT IS TORN, BROKEN OR SHOWS

ANY SIGNS OF TAMPERING

IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION , NAME, ADDRESS AND EMPTY PACKAGE TO:
DISCOUNT DRUG MART, 211 COMMERCE DRIVE MEDINNA, OHIO 44256

NIGHT TIME MULTI-SYMPTOM COLD AND FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-603
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg

DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
---	----------------------	---------

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	603
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-603-03	2 in 1 CARTON	11/08/2023	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/08/2023	

Labeler - Discount Drug Mart (047741335)