# WELBY OMEPRAZOLE- omeprazole tablet, delayed release Aldi Inc

-----

Aldi Inc. Omeprazole Drug Facts

## Active ingredient (in each tablet)

Omeprazole 20 mg

## **Purpose**

Acid reducer

## Use

- treats frequent heartburn (occurs <u>2 or more</u> days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

## **Warnings**

**Allergy alert:** Do not use if you are allergic to omeprazole

## Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

## Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

## Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

## Stop use and ask a doctor if:

• your heartburn continues or worsens

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## **Directions**

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

## **14-Day Course of Treatment**

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- · do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets

## Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

## Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

## Inactive ingredients

carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

## **Questions or comments?**

1-800-719-9260

## **Principal Display Panel**

Compare to Prilosec OTC®

See current Drug Facts

Omeprazole

delayed release tablets 20 mg

acid reducer

Treats Frequent Heartburn!

24HR

actual size

14 TABLETS

One 14-day course of treatment

May take 1 to 4 days for full effect





Compare to Prilosec OTC®\*

See current Drug Facts



# Omeprazole

delayed release tablets 20 mg

acid reducer

Treats Frequent Heartburn!





**14** TABLETS

One 14-day course of treatment May take 1 to 4 days for full effect

## Drug Facts

Active ingredient (in each tablet)

Purpose

91574 1D C4

■treats frequent hearbum (occurs 2 ormore days aweek) ■not intended for immediate relief of hearbum; this drug may take 1 to 4 days for full effect

All ergy alert: Donot us eif you are allergic toome prazole

## Do not use if you have:

- ■trouble or pain swallowingfood, vorning withblood, or bloody or black stools
- ■heartburn with light headedness, sweating ord izz in ess
- ■c hest painor shoulder pain with shortre as of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■frequentchest pain

These may be signs of a serious condition. See your doc br.

## Ask a doctor before use if you have:

- ■had heartburn over 3 months. This may be a sign of a more serious condition.
- ■frequentwheezing, part cularly with heartburn ■unexplained weight loss
- ■nausea or vomiting
- s tomach pain

Ask a doctor orp harmacist beforeuse if you are taking a prescription drug. Acid reducers may interact with certain prescript on drugs.

## Stop u se and ask a doctor if:

- your heartburns ontinues or worsers
- ■you need to take this product for more than 14 days
- ■you need to take more than 1 courseof treatment every 4 months
- ■v ou oet diarrhea

## Drug Facts (continued)

If pregnant orbreast-feeding, ask a healthprofessional before use. Keep out of reach of children . In case ofoverdose, get medical help or contact a Pois on Control Center right away (1-800-222-1222).

## Directions

- ■foradults18 years of ageand older
- ■this product is to be used onceaday (ever y 24hours), every day for 14 days ■itmay take 1 to 4 days for full effect, some peoplegetcomplete relief of symptoms within 24 hours

## 14-DayCourse of Treatment

- s wallow 1 tabletwith aglass of water before eating in the morning
- ■take every day for 14 days
- ■do not take more than 1 table t aday
- ■do not use for more than 14 days unless directed by your doctor
- ■s wallow whole. Donotchew or crush tablets.

### Repeated 14-Day Courses (if needed)

- ■y ou may repeat a14-day course every 4 morths
- ■do not take formore than 14d aysor moreoften than every 4 month s un less directed by a doctor
- ■children under 18 years of age: as ka doc br. He artburn inchildren may sometimes becaused by as erious condition.

- ■read the directions andwarnings before use
- keep the carton. It contains important in 6 m ation.
- ■s bre at 20-25°C(68-77° F) and protect from moisture

Inactive ingredients cama uba wax, ferric oxidered, ferric oxide yellow, hypromellose, hypromellose a cetates us cinate, lac bise monohydrate, moncethanolamine, propylene glyc ol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium steary lium arate, talc, itarium dioxide, triethyl citrate

product is not manufactured or distribute octer & Gamble , distributor of Prilose c OT fety Feature - Do not use if printed blet blister unit is open ortorn.

## **WELBY OMEPRAZOLE**

omeprazole tablet, delayed release

## **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:64024-915

Route of Administration ORAL

## **Active Ingredient/Active Moiety**

Ingredient Name Basis of Strength Strength

OMEPRAZOLE (UNII: KG60484QX9) (OMEPRAZOLE - UNII: KG60484QX9) OMEPRAZOLE 20 mg

## **Inactive Ingredients**

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)

LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)
MONOETHANOLAMINE (UNII: 5KV86114PT)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
SODIUM LAURYL SULFATE (UNII: 368GB5141J)

FERRIC OXIDE YELLOW (UNII: EX43802MRT)

**SODIUM STEARATE** (UNII: QU7E2XA9TG)

**SODIUM STEARYL FUMARATE** (UNII: 7CV7WJK4UI)

TALC (UNII: 7SEV7J4R1U)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

## **Product Characteristics**

Color	BROWN	Score	no score
Shape	OVAL	Size	12mm
Flavor		Imprint Code	20
Contains			

## **Packaging**

7	# Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-------------	---------------------	-------------------------	-----------------------

1	NDC:64024-915- 74	14 in 1 CARTON	10/04/2017		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product			
Marketing Information					
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NI	DA	NDA022032	10/04/2017		

## **Labeler -** Aldi Inc (944259522)

Revised: 11/2022 Aldi Inc