

DAYTIME LIQUID CAPS- acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

Strategic Sourcing Services, LLC (Health Mart)

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Health Mart 44-659

***Active ingredients
(in each liquid-filled capsule)***

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - nasal congestion
 - headache
 - fever
 - minor aches and pains
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a

doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease
- thyroid disease
- diabetes
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- heart disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- persistent cough lasts for more than one week, tends to recur, or is accompanied by fever, rash, or persistent headache

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- do not take more than 8 capsules per 24 hours
- adults and children 12 years and over: take 2 capsules with water every 4 hours
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

- protect from heat, humidity and light
- see end flap for expiration date and lot number

Inactive ingredients

edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, lecithin, mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol

Questions or comments?

1-800-426-9391

Principal Display Panel

NDC 62011-0355-1

Health Mart®

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu LiquiCaps®*

MULTI-SYMPTOM

Daytime Liquid Caps

Acetaminophen - Pain Reliever /Fever Reducer

Dextromethorphan HBr - Cough Suppressant

Phenylephrine HCl - Nasal Decongestant

Multi-Symptom Cold & Flu Relief

Headache, Fever, Sore Throat, Nasal Congestion, Cough

Non-Drowsy • Alcohol-Free • Antihistamine-Free

24 Liquid Caps

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Vicks® DayQuil® Cold & Flu LiquiCaps®.

Product of China

Packaged and Quality Assured in the USA



MULTI-SYMP TOM

Daytime Liquid Caps

NDC 62011-0355-1



Health Mart.

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu LiquiCaps®*

MULTI-SYMP TOM

Daytime Liquid Caps

Acetaminophen - Pain Reliever/Fever Reducer

Dextromethorphan HBr - Cough Suppressant

Phenylephrine HCl - Nasal Decongestant

Multi-Symptom Cold & Flu Relief

Headache, Fever, Sore Throat, Nasal Congestion, Cough

Non-Drowsy • Alcohol-Free • Antihistamine-Free



Actual Size


24 Liquid Caps

MCKESSON
Another Quality Product
Distributed by McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee
healthmart.com

*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Vicks® DayQuil® Cold & Flu LiquiCaps®.
Product of China
Packaged and Quality Assured in the USA

OMI T I M O N

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING



8

Drug Facts (continued)

Other Information
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

■ see end flap for expiration date and lot number

■ protect from heat, humidity and light

Drug Facts (continued)

Inactive ingredients edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, lecithin, mineral oil, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitol

Questions or comments? 1-800-426-9391

50844 ORG021665908

Drug Facts

Active ingredients
Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Other ingredients
Nasal decongestant
Cough suppressant
Pain reliever/fever reducer

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients

Uses
■ temporarily relieves common cold and flu symptoms:
■ sore throat ■ nasal congestion
■ headache ■ fever ■ minor aches and pains
■ cough due to minor throat and bronchial irritation

Directions
■ do not take more than 8 capsules per 24 hours
■ adults and children 12 years and over: take 2 capsules with water every 4 hours
■ children under 12 years: ask a doctor

Warnings
When using this product do not exceed recommended dosage.
Stop use and ask a doctor if:
■ nervousness, dizziness, or sleeplessness occur
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ persistent cough lasts for more than one week, or persistent headache accompanied by fever, rash, or persistent headache
■ If pregnant or breast-feeding, ask a health professional before use. These could be signs of a serious condition.
Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
■ do not take more than directed
■ do not take more than 8 capsules per 24 hours
■ adults and children 12 years and over: take 2 capsules with water every 4 hours
■ children under 12 years: ask a doctor

Drug Facts (continued)
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ Ask a doctor before use if you have liver disease ■ thyroid disease ■ diabetes ■ emphysema ■ heart disease ■ high blood pressure
■ persistent or chronic cough such as occurs with smoking, asthma, or difficulty in urination due to enlargement of the prostate gland
■ cough that occurs with too much phlegm (mucus)
■ difficulty in urination due to enlargement of the prostate gland
■ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
When using this product do not exceed recommended dosage.

Drug Facts (continued)
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ Ask a doctor before use if you have liver disease ■ thyroid disease ■ diabetes ■ emphysema ■ heart disease ■ high blood pressure
■ persistent or chronic cough such as occurs with smoking, asthma, or difficulty in urination due to enlargement of the prostate gland
■ cough that occurs with too much phlegm (mucus)
■ difficulty in urination due to enlargement of the prostate gland
■ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
When using this product do not exceed recommended dosage.
Stop use and ask a doctor if:
■ nervousness, dizziness, or sleeplessness occur
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ persistent cough lasts for more than one week, or persistent headache accompanied by fever, rash, or persistent headache
■ If pregnant or breast-feeding, ask a health professional before use. These could be signs of a serious condition.
Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
■ do not take more than directed
■ do not take more than 8 capsules per 24 hours
■ adults and children 12 years and over: take 2 capsules with water every 4 hours
■ children under 12 years: ask a doctor

B-1241-659-08-R
ORG021665908

DAYTIME LIQUID CAPS

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62011-0355
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
MINERAL OIL (UNII: T5L8T28FGP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SORBITOL (UNII: 506T60A25R)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	

Product Characteristics

Color	ORANGE (clear)	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	659
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62011-0355-1	2 in 1 CARTON	03/01/2015	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	03/01/2015	

Labeler - Strategic Sourcing Services, LLC (Health Mart) (116956644)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(62011-0355)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(62011-0355)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(62011-0355)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(62011-0355)

Revised: 1/2020

Strategic Sourcing Services, LLC (Health Mart)