

PHENOBARBITAL- phenobarbital tablet
QUAGEN PHARMACEUTICALS LLC


Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

PHENOBARBITAL TABLETS, USP C-IV
Rx only

WARNING: MAY BE HABIT-FORMING

DESCRIPTION

Phenobarbital is a barbituric acid derivative for oral administration and occurs as a white, odorless, slightly bitter powder that is soluble in chloroform, freely soluble in alcohol or ether, and slightly soluble in water. Its saturated solution has a pH of about 5.6. Chemically, it is 5-ethyl-5-phenylbarbituric acid with the molecular formula $C_{12}H_{12}N_2O_3$ (232.24). The structural formula is as follows:



Each Phenobarbital Tablet, USP contains 15 mg, 30 mg, 60 mg or 100 mg of phenobarbital, USP.

Inactive ingredients include: colloidal silicon dioxide, lactose monohydrate, microcrystalline cellulose, sodium starch glycolate and magnesium stearate.

CLINICAL PHARMACOLOGY

Phenobarbital, a long-acting barbiturate, is a central nervous system depressant. In ordinary doses, the drug acts as a sedative and anticonvulsant. Its onset of action occurs within 30 minutes, and the duration of action ranges from 5 to 6 hours. It is detoxified in the liver.

INDICATIONS AND USAGE

Phenobarbital Tablets, USP are indicated for use as a sedative or anticonvulsant.

CONTRAINDICATIONS

Phenobarbital is contraindicated in patients who are hypersensitive to barbiturates. In such patients, severe hepatic damage can occur from ordinary doses and is usually associated with dermatitis and involvement of parenchymatous organs. A personal or familial history of acute intermittent porphyria represents one of the few absolute contraindications to the use of barbiturates. Phenobarbital is also contraindicated in patients with marked impairment of liver function, or respiratory disease in which dyspnea or obstruction is evident. It should not be administered to persons with known previous addiction to the sedative/hypnotic group, since ordinary doses may be ineffectual and may contribute to further addiction.

WARNINGS

In small doses, the barbiturates may increase the reaction to painful stimuli. Taken by themselves, the barbiturates cannot be relied upon to relieve pain or even to produce sedation or sleep in the presence of severe pain.

PRECAUTIONS

General Precautions

Barbiturates induce liver microsomal enzyme activity. This accelerates the biotransformation of various drugs and is probably part of the mechanism of the tolerance encountered with barbiturates.

Phenobarbital, therefore, should be used with caution in patients with decreased liver function. This drug should also be administered cautiously to patients with a history of drug dependence or abuse (see **DRUG ABUSE AND DEPENDENCE**).

Phenobarbital may decrease the potency of coumarin anticoagulants; therefore, patients receiving such concomitant therapy should have more frequent prothrombin determinations. As with other sedatives and hypnotics, elderly or debilitated patients may react to barbiturates with marked excitement, depression, or confusion.

The systemic effects of exogenous hydrocortisone and endogenous hydrocortisone (cortisol) may be diminished by phenobarbital. Thus, this product should be administered with caution to patients with borderline hypoadrenal function, regardless of whether it is of pituitary or of primary adrenal origin.

Information for Patients

Phenobarbital may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks, such as driving a car or operating machinery. The patient should be cautioned accordingly.

Drug Interactions

Phenobarbital in combination with alcohol, tranquilizers, and other central nervous system depressants has additive depressant effects, and the patient should be so advised. Patients taking this drug should be warned not to exceed the dosage recommended by their physician. Toxic effects and fatalities have occurred following overdoses of phenobarbital alone and in combination with other central nervous system depressants. Caution should be exercised in prescribing unnecessarily large amounts of phenobarbital for patients who have a history of emotional disturbances or suicidal ideation or who have misused alcohol and other CNS drugs (see **OVERDOSAGE**).

Usage in Pregnancy:

Pregnancy Category B - Reproduction studies have been performed in animals and have revealed no evidence of impaired fertility or harm to the fetus due to phenobarbital. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers

Caution should be exercised when phenobarbital is administered to a nursing woman.

ADVERSE REACTIONS

The following adverse reactions have been reported:

CNS Depression:

Residual sedation or "hangover", drowsiness, lethargy, and vertigo. Emotional disturbances and phobias may be accentuated. In some persons, barbiturates such as phenobarbital repeatedly produce excitement rather than depression, and the patient may appear to be inebriated. Like other non-analgesic hypnotic drugs, barbiturates, such as phenobarbital, when given in the presence of pain, may cause restlessness, excitement, and even delirium. Rarely, the use of barbiturates results in localized or diffused myalgic, neuralgic, or arthritic pain, especially in psychoneurotic patients with insomnia. The pain may appear in paroxysms, is most intense in the early morning hours, and is most frequently located in the region of the neck, shoulder girdle, and upper limbs. Symptoms may last for days after the drug is discontinued.

Respiratory/Circulatory:

Respiratory depression, apnea, circulatory collapse.

Allergic:

Acquired hypersensitivity to barbiturates consists chiefly in allergic reactions that occur especially in persons who tend to have asthma, urticaria, angioedema, and similar conditions. Hypersensitivity reactions in this category include localized swelling, particularly of the eyelids, cheeks, or lips, and erythematous dermatitis. Rarely, exfoliative dermatitis (e.g., Stevens-Johnson syndrome and toxic epidermal necrolysis) may be caused by phenobarbital and can prove fatal. The skin eruption may be associated with fever, delirium, and marked degenerative changes in the liver and other parenchymatous organs. In a few cases, megaloblastic anemia has been associated with

the chronic use of phenobarbital.

Other:

Nausea and vomiting; headache.

To report SUSPECTED ADVERSE REACTIONS, contact Rising Pharma Holdings, Inc. at 1-844-874-7464, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG ABUSE AND DEPENDENCE

Controlled Substance - Phenobarbital is a Schedule IV drug.

Dependence

Prolonged, uninterrupted use of barbiturates (particularly the short-acting drugs), even in therapeutic doses, may result in psychic and physical dependence. Withdrawal symptoms due to physical dependence following chronic use of large doses of barbiturates may include delirium, convulsions, and death.

OVERDOSAGE

The signs and symptoms of barbiturate poisoning are referable especially to the central nervous system and the cardiovascular system. Moderate intoxication resembles alcoholic inebriation. In severe intoxication, the patient is comatose, the level of reflex activity conforming in a general way to the intensity of the central depression. The deep reflexes may persist for some time despite coexistent coma. The Babinski sign is often positive. The EEG may be of the “burst-suppression” type, with brief periods of electrical silence. The pupils may be constricted and react to light, but late in the course of barbiturate poisoning they may show hypoxic paralytic dilatation. Respiration is affected early. Breathing may be either slow or rapid and shallow; Cheyne-Stokes rhythm may be present. Respiratory minute volume is diminished, and hypoxia and respiratory acidosis may develop. The blood pressure falls, owing partly to depression of medullary vasomotor centers; partly to a direct action of the drug on the myocardium, sympathetic ganglia, and vascular smooth muscle; partly to hypoxia.

The patient thus develops a typical shock syndrome, with a weak and rapid pulse, cold and clammy skin, and a rise in the hematocrit. Respiratory complications (atelectasis, pulmonary edema, and bronchopneumonia) and renal failure are much dreaded and not infrequent concomitant of severe barbiturate poisoning. There is usually hypothermia, sometimes with temperatures as low as 32°C.

Treatment:

General management should consist of symptomatic and supportive therapy, including gastric lavage, administration of intravenous fluids, and maintenance of blood pressure, body temperature and adequate respiratory exchange. Dialysis will increase the rate of removal of barbiturates from the body fluids. Antibiotics may be required to control pulmonary complications.

DOSAGE AND ADMINISTRATION

Oral Sedative Dose, Adults - 30 to 120 mg daily in 2 or 3 divided doses. Children - 6 mg/kg of body weight daily in 3 divided doses.

Oral Hypnotic Dose, Adults - 100 to 320 mg.

Oral Anticonvulsant Dose, Adults - 50 to 100 mg 2 or 3 times daily.

Children - 15 to 50 mg 2 or 3 times daily.

HOW SUPPLIED

Phenobarbital Tablets, USP 15 mg: White to off-white round, biconvex tablets debossed "T35" on one side and plain on the other side.

Available in bottles of 100 tablets, NDC 70752-165-10; 500 tablets, 70752-165-14; and 1000 tablets, NDC 70752-165-11.

Phenobarbital Tablets, USP 30 mg: White to off-white round, biconvex tablets debossed "T" above and "36" below the bisect on one side and plain on the other side.

Available in bottles of 100 tablets, NDC 70752-166-10; 500 tablets, NDC 70752-166-14; and 1000 tablets, NDC 70752-166-11.

Phenobarbital Tablets, USP 60 mg: White to off-white round, biconvex tablets debossed "T" above and "37" below the bisect on one side and plain on the other side.

Available in bottles of 100 tablets, NDC 70752-167-10; 500 tablets, NDC 70752-167-14; and 1000 tablets, NDC 70752-167-11.

Phenobarbital Tablets, USP 100 mg: White to off-white round, biconvex tablets debossed "T" above and "38" below the bisect on one side and plain on the other side.

Available in bottles of 100 tablets, NDC 70752-168-10; 500 tablets, NDC 70752-168-14; and 1000 tablets, NDC 70752-168-11.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

Manufactured by:

Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006

52024
Rev. 07/20

PRINCIPAL DISPLAY PANEL

NDC 70752-165-10
Phenobarbital Tablets, USP CIV
15 mg
Rx Only

100 Tablets

Rising
PHARMACEUTICALS

NDC 16571-670-01

Phenobarbital*
Tablets, USP CIV

15 mg

*WARNING: May be habit forming.

100 Tablets **Rx Only**

Each tablet contains:
Phenobarbital, USP..... 15 mg
***WARNING: May be habit forming.**
USUAL DOSAGE ORAL: See package insert for full prescribing information.
Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.
Storage: Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.
Distributed by:
Rising Pharma Holdings, Inc. 50056
East Brunswick, NJ 08816 Rev. 06/20
Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006

NON VARNISH AREA
1.5" X 1.00"

3 16571-67001 7

NDC 70752-165-14
Phenobarbital Tablets, USP CIV
15 mg
Rx Only
500 Tablets

Rising
PHARMACEUTICALS

NDC 16571-670-50

Phenobarbital*
Tablets, USP CIV

15 mg

*WARNING: May be habit forming.

500 Tablets **Rx Only**

Each tablet contains:
Phenobarbital, USP..... 15 mg
***WARNING: May be habit forming.**
USUAL DOSAGE ORAL: See package insert for full prescribing information.
Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.
Storage: Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.
Distributed by:
Rising Pharma Holdings, Inc. 50057
East Brunswick, NJ 08816 Rev. 06/20
Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006


NON VARNISH AREA
1.75" X 1.00"

3 16571-67050 5

NDC 70752-165-11
Phenobarbital Tablets, USP CIV
15 mg
Rx Only
1000 Tablets

Rising
PHARMACEUTICALS

NDC 16571-670-10

Phenobarbital*
Tablets, USP 

15 mg

*WARNING: May be habit forming.

1000 Tablets **Rx Only**

Each tablet contains:
Phenobarbital, USP.....15 mg

***WARNING: May be habit forming.**


USUAL DOSAGE ORAL: See package insert for full prescribing information.

Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.

Storage: Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.

Distributed by:
Rising Pharma Holdings, Inc. 50058
East Brunswick, NJ 08816 Rev. 06/20

Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006


N 16571-67010 9

2.75" X 1.00"
NON VARNISH AREA

NDC 70752-166-10
Phenobarbital Tablets, USP CIV
30 mg
Rx Only
100 Tablets

Rising
PHARMACEUTICALS

NDC 16571-672-01

Phenobarbital*
Tablets, USP 

30 mg

*WARNING: May be habit forming.

100 Tablets **Rx Only**

Each tablet contains:
Phenobarbital, USP.....30 mg

***WARNING: May be habit forming.**

USUAL DOSAGE ORAL: See package insert for full prescribing information.

Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.

Storage: Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.

Distributed by:
Rising Pharma Holdings, Inc. 50062
East Brunswick, NJ 08816 Rev. 06/20

Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006


N 16571-67201 1

1.5" X 1.00"
NON VARNISH AREA

NDC 70752-166-14
Phenobarbital Tablets, USP CIV
30 mg
Rx Only
500 Tablets

Rising
PHARMACEUTICALS

NDC 16571-672-50

Phenobarbital*
Tablets, USP 

30 mg

*WARNING: May be habit forming.

500 Tablets **Rx Only**

Each tablet contains:
Phenobarbital, USP.....30 mg

***WARNING: May be habit forming.**

USUAL DOSAGE ORAL: See package insert for full prescribing information.

Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.

Storage: Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.

Distributed by:
Rising Pharma Holdings, Inc. 50063
East Brunswick, NJ 08816 Rev. 06/20

Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006



N 3 16571-67250 9

2.00" X 1.00"
NON VARNISH AREA

NDC 70752-166-11
Phenobarbital Tablets, USP CIV
30 mg
Rx Only
1000 Tablets

Rising
PHARMACEUTICALS

NDC 16571-672-10

Phenobarbital*
Tablets, USP 

30 mg

*WARNING: May be habit forming.

1000 Tablets **Rx Only**

Each tablet contains:
Phenobarbital, USP.....30 mg

***WARNING: May be habit forming.**


USUAL DOSAGE ORAL: See package insert for full prescribing information.

Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.

Storage: Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.

Distributed by:
Rising Pharma Holdings, Inc. 50064
East Brunswick, NJ 08816 Rev. 06/20

Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006


N 3 16571-67210 3

2.75" X 1.00"
NON VARNISH AREA

NDC 70752-167-10
Phenobarbital Tablets, USP CIV
60 mg
Rx Only
100 Tablets

Rising
PHARMACEUTICALS

NDC 16571-674-01

Phenobarbital*
Tablets, USP 

60 mg

*WARNING: May be habit forming.

100 Tablets **Rx Only**

Each tablet contains:
Phenobarbital, USP.....60 mg
***WARNING: May be habit forming.**
USUAL DOSAGE ORAL: See package insert for full prescribing information.
Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.
Storage: Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.
Distributed by:
Rising Pharma Holdings, Inc.
East Brunswick, NJ 08816 50068
Rev. 06/20
Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006



N 16571-67401 5

1.75" X 1.00"
NON VARNISH AREA

NDC 70752-167-14
Phenobarbital Tablets, USP CIV
60 mg
Rx Only
500 Tablets

Rising
PHARMACEUTICALS

NDC 16571-674-50


Phenobarbital*
Tablets, USP 

60 mg

*WARNING: May be habit forming.

500 Tablets **Rx Only**

Each tablet contains:
Phenobarbital, USP.....60 mg
***WARNING: May be habit forming.**
USUAL DOSAGE ORAL: See package insert for full prescribing information.
Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.
Storage: Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.
Distributed by:
Rising Pharma Holdings, Inc.
East Brunswick, NJ 08816 50069
Rev. 06/20
Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006



N 16571-67450 3

2.75" X 1.00"
NON VARNISH AREA

NDC 70752-167-11
Phenobarbital Tablets, USP CIV
60 mg
Rx Only
1000 Tablets

Rising
PHARMACEUTICALS

NDC 16571-674-10

Phenobarbital*
Tablets, USP 

60 mg

*WARNING: May be habit forming.

1000 Tablets **Rx Only**

Each tablet contains:
Phenobarbital, USP.....60 mg

***WARNING: May be habit forming.**

USUAL DOSAGE ORAL: See package insert for full prescribing information.


Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.

Storage: Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.

Distributed by:
Rising Pharma Holdings, Inc.
East Brunswick, NJ 08816 50070

Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006

Rev. 06/20



3 16571-67410 7

3.00" X 1.00"
NON VARNISH AREA

NDC 70752-168-10
Phenobarbital Tablets, USP CIV
100 mg
Rx Only
100 Tablets

Rising
PHARMACEUTICALS

NDC 16571-669-01

Phenobarbital*
Tablets, USP 

100 mg

*WARNING: May be habit forming.

100 Tablets **Rx Only**

Each tablet contains:
Phenobarbital, USP.....100 mg

***WARNING: May be habit forming.**

USUAL DOSAGE ORAL: See package insert for full prescribing information.

Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.

Storage: Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.

Distributed by:
Rising Pharma Holdings, Inc.
East Brunswick, NJ 08816 50077

Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006


Rev. 06/20



3 16571-66901 1

1.75" X 1.00"
NON VARNISH AREA

NDC 70752-168-14
Phenobarbital Tablets, USP CIV
100 mg
Rx Only
500 Tablets



NDC 16571-669-50

Phenobarbital*

Tablets, USP CIV

100 mg

*WARNING: May be habit forming.

500 Tablets

Rx Only

Each tablet contains:
Phenobarbital, USP.....100 mg

***WARNING: May be habit forming.**

USUAL DOSAGE ORAL: See package insert for full prescribing information.


Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.

Storage: Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.

Distributed by:
Rising Pharma Holdings, Inc.
East Brunswick, NJ 08816

Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006


50078
Rev. 06/20



16571-66950

3.00" X 1.00"
NON VARNISH AREA

NDC 70752-168-11
Phenobarbital Tablets, USP CIV
100 mg
Rx Only
1000 Tablets



NDC 16571-669-10

Phenobarbital*

Tablets, USP CIV

100 mg

*WARNING: May be habit forming.

1000 Tablets

Rx Only

Each tablet contains:
Phenobarbital, USP.....100 mg

***WARNING: May be habit forming.**

USUAL DOSAGE ORAL: See package insert for full prescribing information.


Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.

Storage: Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.

Distributed by:
Rising Pharma Holdings, Inc.
East Brunswick, NJ 08816

Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006

50079
Rev. 06/20



16571-66910

3.00" X 1.00"
NON VARNISH AREA

PHENOBARBITAL

phenobarbital tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70752-165
Route of Administration	ORAL	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENOBARBITAL (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)	PHENOBARBITAL	15 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white (White to off-white)	Score	no score
Shape	ROUND (Round, Biconvex)	Size	6mm
Flavor		Imprint Code	T;35
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70752-165-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	
2	NDC:70752-165-14	500 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	
3	NDC:70752-165-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Unapproved drug other		09/30/2020	

PHENOBARBITAL

phenobarbital tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70752-166
Route of Administration	ORAL	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

PHENOBARBITAL (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)	PHENOBARBITAL	30 mg		
Inactive Ingredients				
Ingredient Name		Strength		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
Product Characteristics				
Color	white (White to off-white)	Score	2 pieces	
Shape	ROUND (Round, Biconvex)	Size	8mm	
Flavor		Imprint Code	T;36	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70752-166-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	
2	NDC:70752-166-14	500 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	
3	NDC:70752-166-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
Unapproved drug other		09/30/2020		

PHENOBARBITAL			
phenobarbital tablet			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70752-167
Route of Administration	ORAL	DEA Schedule	CIV
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
PHENOBARBITAL (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)		PHENOBARBITAL	60 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white (White to off-white)	Score	2 pieces
Shape	ROUND (Round, Biconvex)	Size	10mm
Flavor		Imprint Code	T;37
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70752-167-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	
2	NDC:70752-167-14	500 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	
3	NDC:70752-167-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Unapproved drug other		09/30/2020	

PHENOBARBITAL

phenobarbital tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70752-168
Route of Administration	ORAL	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENOBARBITAL (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)	PHENOBARBITAL	100 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white (White to off-white)	Score	2 pieces
Shape	ROUND (Round, Biconvex)	Size	12mm
Flavor		Imprint Code	T;38
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70752-168-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	
2	NDC:70752-168-14	500 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	
3	NDC:70752-168-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Unapproved drug other		09/30/2020	

Labeler - QUAGEN PHARMACEUTICALS LLC (073645339)

Registrant - QUAGEN PHARMACEUTICALS LLC (073645339)

Establishment

Name	Address	ID/FEI	Business Operations
QUAGEN PHARMACEUTICALS LLC		080281331	MANUFACTURE(70752-165, 70752-166, 70752-167, 70752-168) , PACK(70752-165, 70752-166, 70752-167, 70752-168)

Revised: 2/2023

QUAGEN PHARMACEUTICALS LLC