

**EFFERVESCENT ORANGE COLD RELIEF- aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent**  
**Meijer Distribution Inc.**

-----  
**Effervescent Cold Relief - Orange**

**Active Ingredients**

**(in each effervescent tablet)**

Aspirin 325 mg **(NSAID)\***

Chlorpheniramine maleate 2 mg

Phenylephrine bitartrate 7.8 mg

\*Nonsteroidal anti-inflammatory drug

**Purpose**

Aspirin 325 mg **(NSAID)\***. . . . .Pain reliever/fever reducer

Chlorpheniramine maleate 2 mg . . . . .Antihistamine

Phenylephrine bitartrate 7.8 mg . . . . .Nasal decongestant

\*Nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves these symptoms due to the common cold:
- sneezing ■ nasal congestion ■ sore throat ■ headache
- minor aches and pains ■ runny nose ■ sinus congestion and pressure
- temporarily reduces fever

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using

this product ■ take more or for a longer time than directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ if you are allergic to aspirin ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients.

### **Ask a doctor before use if**

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

### **Ask a doctor before use if you have**

**Ask a doctor before use if you have** ■ asthma ■ glaucoma ■ diabetes ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers ■ presently taking a prescription drug ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

### **When using this product**

#### **When using this product ■ do not exceed recommended dosage**

- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding

- feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- nervousness, dizziness or sleeplessness occurs

### **If pregnant or breast-feeding**

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

**Keep out of reach of children.**

### **In case of overdose**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4 hours
- do not take more than 8 tablets in 24 hours
- *children under 12 years: ask a doctor*

### **Other information**

**each tablet contains:** sodium 464 mg

- phenylketonurics: contains phenylalanine 9 mg per tablet
- store at room temperature (59°-86°F)

**Inactive ingredients**

acesulfame potassium, anhydrous citric acid, aspartame, docusate sodium, FD&C red #40, FD&C yellow #6, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

**Questions or comments?****Questions or comments? Call 1-888-228-6937****PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

PN: 87524 MJ C1  
REV: 04/24



PH 3820970  
BOX 1744  
PACKET 10  
QUANTITY 24  
DISTRIBUTOR  
DIST. BY MEJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49514  
This product is not manufactured or distributed by Bayer, owner of the registered trademark Alka-Seltzer®.

**DO NOT USE IF PRINTED**  
**PACKETS ARE TORN OR PUNCTURED**  
**Call 1-888-228-6697**  
**(Questions or comments?)**  
bicarbonate  
sodium benzoate, sodium  
#6, flavors, mannitol, povidone,  
sodium, FD&C red #40, FD&C yellow  
citric acid, aspartame, docusate  
ascorbic acid, anhydrous  
(99-86-7)  
store at room temperature  
phenylephrine 9 mg per tablet  
phenylephrine: contains  
each tablet contains: sodium 464 mg  
**Other Information**  
**Drug Facts (continued)**


meijer  
**Cold Relief**  
Effervescent  
ORANGE  
Fast Relief of:  
Nasal Congestion  
Runny Nose  
Sneezing  
Headache  
Sinus Pain & Pressure  
COMPARE TO  
ALKA-SELTZER PLUS® COLD  
ORANGE ZEST  
ACTIVE INGREDIENTS\*  
NDC 41250-123-24  
meijer  
**Cold Relief**  
Effervescent  
ORANGE  
Fast Relief of:  
Nasal Congestion  
Runny Nose  
Sneezing  
Headache  
Sinus Pain & Pressure  
COMPARE TO  
ALKA-SELTZER PLUS® COLD  
ORANGE ZEST  
ACTIVE INGREDIENTS\*  
24 TABLETS  
24 TABLETS

**Drug Facts (continued)**  
an allergic reaction occurs. Seek medical help right away.  
pain or nasal congestion gets worse or lasts more than 7 days  
never gets worse or lasts more than 3 days  
new symptoms occur  
parosmia or swelling is present  
ringing in the ears or a loss of hearing occurs  
dizziness or sleeplessness occurs  
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.  
**Directions**  
do not use more than directed (see overdose warning)  
adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4 hours  
do not take more than 8 tablets in 24 hours  
child 6 to 11 years: ask a doctor  
child under 12 years: ask a doctor



**Drug Facts**  
**Active ingredients**  
**(in each effervescent tablet)**      **Purpose**  
Aspirin 325 mg (NSAID)\* ..... Pain reliever/fever reducer  
Chlorpheniramine maleate 2 mg ..... Antihistamine  
Phenylephrine bitartrate 7.8 mg ..... Nasal decongestant  
\*Nonsteroidal anti-inflammatory drug  
**Uses**  
temporarily relieves these symptoms due to the common cold:  
sneezing ■ nasal congestion ■ sore throat  
headache ■ minor aches and pains ■ runny nose  
sinus congestion and pressure  
temporarily reduces fever  
**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: hives ■ facial swelling ■ asthma (wheezing) ■ shock  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or

**Drug Facts (continued)**  
for a longer time than directed  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ if you are allergic to aspirin ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients.  
**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic  
**Ask a doctor before use if you have** ■ asthma ■ glaucoma ■ diabetes ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet  
**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers ■ presently taking a prescription drug ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis  
**When using this product** ■ do not exceed recommended dosage ■ you may get drowsy ■ avoid alcoholic drinks ■ excitability may occur, especially in children ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery  
**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding  
feel faint ■ vomit blood ■ have bloody or black stools  
have stomach pain that does not get better



# EFFERVESCENT ORANGE COLD RELIEF

aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-123
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PHENYLEPHRINE BITARTRATE</b> (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>DOCUSATE SODIUM</b> (UNII: F05Q2T2JA0)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	25mm
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-123-20	10 in 1 CARTON	05/26/2013	12/31/2026
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:41250-123-24	12 in 1 CARTON	06/06/2024	
3		2 in 1 POUCH; Type 0: Not a Combination		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/26/2013	

**Labeler** - Meijer Distribution Inc. (006959555)

Revised: 12/2024

Meijer Distribution Inc.