

**HABITROL NICOTINE POLACRILEX GUM 2 MG COATED MINT- nicotine  
polacrilex gum, chewing  
Dr. Reddy's Laboratories Inc.**

-----  
**Nicotine Polacrilex Gum Mint 2 mg**

***Drug Facts***

***Active ingredient (in each chewing piece) - 2 mg***

Nicotine polacrilex (equal 2 mg (nicotine))

***Purpose***

Stop smoking aid

***Use***

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

***Warnings***

**If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.**

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center(1-800-22-1222) right away.

**Directions - 2 mg**

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum according to the following 12 week schedule:

<b>Weeks 1 to 6</b>	<b>Weeks 7 to 9</b>	<b>Weeks 10 to 12</b>
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider

**Other information - 2 mg**

- **each piece contains:** calcium 115 mg, sodium 8 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

### ***Inactive ingredients - 2 mg***

acacia, acesulfame potassium, carnauba wax, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, titanium dioxide, xylitol.

### ***Questions or comments?***

Call toll-free **1-800-585-8682** weekdays (8:00 am - 8:00 pm EST)

### **Principal display panel- 2 mg**

\*Compare to the active ingredient of Nicorette® Gum

Habitrol

Nicotine Polacrilex Gum USP,

2 mg (nicotine)

STOP SMOKING AID

Sugar-free

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE **MORE THAN 30 MINUTES** AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg

### **TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** - Chew **at least 9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

### **BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.**

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

\*This product is not manufactured or distributed by Haleon group of companies, distributor of Nicorette® Gum.

DISTRIBUTED BY:

Dr. Reddy's Laboratories, Inc.

Princeton, NJ 08540

## Package label



# HABITROL NICOTINE POLACRILEX GUM 2 MG COATED MINT

nicotine polacrilex gum, chewing

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:75907-302
Route of Administration	BUCCAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	2 mg

Inactive Ingredients	
Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
MAGNESIUM OXIDE (UNII: 3A3U0G171G)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XYLITOL (UNII: VCQ006KQ1E)	

Product Characteristics			
Color	white	Score	no score
Shape	SQUARE	Size	14mm
Flavor	MINT	Imprint Code	
Contains			

PLD-46178 F009496  
06/2025  
DISTRIBUTED BY:  
Dr. Healdy's Laboratories, Inc.  
Princeton, NJ 08540

BLISTER PACKAGED FOR YOUR PROTECTION, DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

not for sale to those under 18 years of age  
a proof of age required  
source when proof of age cannot be verified

**Drug Facts**

**Warnings**

Do not use if you are pregnant or breastfeeding. Nicotine can increase your risk of heart disease, stroke, or high blood pressure. Nicotine can increase your risk of heart disease, stroke, or high blood pressure. Nicotine can increase your risk of heart disease, stroke, or high blood pressure.

**Use**

Use 2 mg nicotine gum (read to 2 mg nicotine) as directed.

**Other Information**

Keep this and all other medicines out of the reach of children. This medicine is not for use in children.

**Habitrol**  
Nicotine Polacrilex Gum  
USP, 2 mg (nicotine)  
Gum  
2 mg  
Sugar-free  
STOP SMOKING AID

NDC 75907-302-80  
\*Compare to the active ingredient in Nicorette® Gum

## HABITROL Nicotine Polacrilex Coated Mint Gum

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75907-302-80	8 in 1 CARTON	06/12/2025	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:75907-302-15	15 in 1 CARTON	06/12/2025	
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:75907-302-10	10 in 1 BLISTER PACK; Type 0: Not a Combination Product	06/12/2025	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078699	06/12/2025	

**Labeler** - Dr. Reddy's Laboratories Inc. (802315887)

Revised: 12/2025

Dr. Reddy's Laboratories Inc.