

MINOXIDIL FOR WOMEN- minoxidil aerosol, foam
Taro Pharmaceuticals U.S.A., Inc.

Minoxidil Topical Aerosol, 5% (For Women)

Drug Facts

Active ingredient

Minoxidil 5% w/w (without propellant)

Purpose

Hair regrowth treatment for women

Use

to regrow hair on the top of the scalp (see pictures on side of this carton)

Warnings

For external use only.

Extremely Flammable

Avoid fire, flame, or smoking during and immediately following application.

Do not use if

- your degree of hair loss is different than that shown on side of this carton because this product may not work for you
- you have no family history of hair loss
- your hair loss is sudden and/or patchy
- your hair loss is associated with childbirth
- you do not know the reason for your hair loss
- you are under 18 years of age. Do not use on babies and children.
- your scalp is red, inflamed, infected, irritated, or painful
- you use other medicines on the scalp

Ask a doctor before use if you have heart disease

When using this product

- do not use more than directed
- do not apply on other parts of the body
- avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water.
- some people have experienced changes in hair color and/or texture
- it takes time to regrow hair. Results may occur at 3 months with **once-daily** use. For

some women, you may need to use this product **once a day** for at least 6 months before you see results.

- the amount of hair regrowth is different for each person. This product will not work for all women.

Stop use and ask a doctor if

- chest pain, rapid heartbeat, faintness, or dizziness occurs
- sudden, unexplained weight gain occurs
- your hands or feet swell
- scalp irritation or redness occurs
- unwanted facial hair growth occurs
- you do not see hair regrowth in 6 months

May be harmful if used when pregnant or breast-feeding.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- see enclosed leaflet for complete directions on how to use
- apply half a capful **once daily** directly to the scalp in the hair loss area
- massage into scalp with fingers, then wash hands well
- using more or more often will not improve results
- continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again

Other information

- see hair loss pictures on side of this carton
- before use, read all information on carton and enclosed leaflet
- keep the carton. It contains important information.
- store at controlled room temperature 20° to 25°C (68° to 77°F)
- contents under pressure. Do not puncture or incinerate container. Do not expose to heat or store at temperatures above 120°F (49°C).

Inactive ingredients

butylated hydroxytoluene, cetyl alcohol, citric acid anhydrous, dehydrated alcohol, isobutane, lactic acid, n-butane, polysorbate 60, propane, purified water, stearyl alcohol

Questions?

- Call **1-866-923-4914**

Distributed by:

Taro Pharmaceuticals U.S.A., Inc.

Hawthorne, NY 10532

PRINCIPAL DISPLAY PANEL - 60 g Can Package

*Compare to the active
ingredient of
Women's Rogaine®

NDC 51672-2129-4

Minoxidil
Topical Aerosol,
5% (For Women)

HAIR REGROWTH
TREATMENT

REACTIVATES
HAIR FOLLICLES
TO STIMULATE
REGROWTH

CLINICALLY
PROVEN TO
REGROW HAIR

Once-a-Day
Foam for
Women

UNSCENTED

#1 DERMATOLOGIST
RECOMMENDED
ACTIVE INGREDIENT†

Two Month
Supply
One 60 g
(2.11 oz) Can

NDC 51672-2129-4

Minoxidil Topical Aerosol, 5% (For Women)

HAIR REGROWTH
TREATMENT



Once-a-Day
Foam for
Women

131
5200621-0223-04

Drug Facts

Active ingredient Purpose
Minoxidil 5% w/w (without propellant).....
.....Hair regrowth treatment for women

Use to regrow hair on the top of the scalp (see pictures on side of this carton)

Warnings

For external use only.

Extremely Flammable: Avoid fire, flame, or smoking during and immediately following application.

Do not use if

- your degree of hair loss is different than that shown on side of this carton because this product may not work for you
- you have no family history of hair loss
- your hair loss is sudden and/or patchy
- your hair loss is associated with childbirth
- you do not know the reason for your hair loss
- you are under 18 years of age. Do not use on babies and children.
- your scalp is red, inflamed, infected, irritated, or painful
- you use other medicines on the scalp

Ask a doctor before use if you have heart disease

When using this product

- do not use more than directed
- do not apply on other parts of the body
- avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water.
- some people have experienced changes in hair color and/or texture
- it takes time to regrow hair. Results may occur at 3 months with once-daily use. For some women, you may need to use this product once a day for at least 6 months before you see results.
- the amount of hair regrowth is different for each person. This product will not work for all women.

Stop use and ask a doctor if

- chest pain, rapid heartbeat, faintness, or dizziness occurs
- sudden, unexplained weight gain occurs
- your hands or feet swell
- scalp irritation or redness occurs
- unwanted facial hair growth occurs
- you do not see hair regrowth in 6 months

May be harmful if used when pregnant or breast-feeding.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)



Drug Facts (continued)

Directions

- see enclosed leaflet for complete directions on how to use
- apply half a capful once daily directly to the scalp in the hair loss area
- massage into scalp with fingers, then wash hands well
- using more or more often will not improve results
- continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again

Other information

- see hair loss pictures on side of this carton
- before use, read all information on carton and enclosed leaflet
- keep the carton. It contains important information.
- store at controlled room temperature 20° to 25°C (68° to 77°F)
- contents under pressure. Do not puncture or incinerate container. Do not expose to heat or store at temperatures above 120°F (49°C).

Inactive ingredients

butylated hydroxytoluene, cetyl alcohol, citric acid anhydrous, dehydrated alcohol, isobutane, lactic acid, n-butane, polysorbate 60, propane, purified water, stearyl alcohol

Questions?

- Call 1-866-923-4914

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Women's Rogaine®

Minoxidil Topical Aerosol, 5% (For Women) is for women who have a general thinning of hair on the top of the scalp (as shown below) and has been shown to regrow hair in women with the following degrees of thinning hair or hair loss



If women have more hair loss than shown above, Minoxidil Topical Aerosol, 5% (For Women) may not work.

THIS PACKAGE CONTAINS:

One 60 g (2.11 oz) can of Minoxidil Topical Aerosol, 5% (For Women) (two month supply)

*Compare to the active ingredient of Women's Rogaine®

NDC 51672-2129-4

Minoxidil Topical Aerosol, 5% (For Women)

HAIR REGROWTH
TREATMENT

REACTIVATES
HAIR FOLLICLES
TO STIMULATE
REGROWTH

CLINICALLY
PROVEN TO
REGROW HAIR



Once-a-Day
Foam for
Women

UNSCENTED

#1 DERMATOLOGIST
RECOMMENDED
ACTIVE INGREDIENT*

Two Month
Supply
One 60 g
(2.11 oz) Can

DIRECTIONS FOR USE

See information leaflet with complete directions on how to use and obtain best results.



1: PREP HANDS WITH COLD WATER

Rinse your fingers in cold water and dry them. Note: foam will melt on contact with warm surfaces.



2: OPEN CHILD RESISTANT CAP

Be sure to align arrow on the cap with arrow on white ring. Tilt cap back and pull off.



3: DISPENSE VERTICALLY ONTO A COLD SURFACE

Hold the can straight upside down. If you hold the can at an angle, foam may not dispense properly. Press nozzle to dispense half a capful of foam onto a cold surface (e.g. dish, or cold hands).



4: APPLY TO SCALP, NOT HAIR

Part your hair to expose hair loss area. Massage foam into scalp, not hair. Repeat until all hair loss areas have been covered.



5: CLOSE CAP AND WASH HANDS

Snap cap back into place. Be sure arrows do not line up, so cap remains child resistant. Wash hands and any surface thoroughly after use.

Read and keep package and leaflet for full product information.



Made in Israel
Distributed by:
Taro Pharmaceuticals U.S.A., Inc.
Hawthorne, NY 10532
TARO is a registered trademark of
Taro Pharmaceuticals U.S.A., Inc.

*In the U.S.
Source: AlphaImpactRx

MINOXIDIL FOR WOMEN

minoxidil aerosol, foam

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51672-2129
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Minoxidil (UNII: 5965120SH1) (Minoxidil - UNII:5965120SH1)	Minoxidil	50 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
butylated hydroxytoluene (UNII: 1P9D0Z171K)	
cetyl alcohol (UNII: 936JST6JCN)	
anhydrous citric acid (UNII: XF417D3PSL)	
alcohol (UNII: 3K9958V90M)	
isobutane (UNII: BXR49TP611)	
lactic acid, unspecified form (UNII: 33X04XA5AT)	
butane (UNII: 6LV4FOR43R)	
polysorbate 60 (UNII: CAL22UVI4M)	
propane (UNII: T75W9911L6)	
water (UNII: 059QF0KO0R)	
stearyl alcohol (UNII: 2KR89I4H1Y)	

Product Characteristics

Color	WHITE (white to off-white foam)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51672-2129-4	1 in 1 PACKAGE	04/22/2019	
1		60 g in 1 CAN; Type 0: Not a Combination Product		
2	NDC:51672-	2 in 1 PACKAGE	04/22/2019	

1	2129-8	2 III 1 PACKAGE	04/22/2019	
2		60 g in 1 CAN; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA209074		04/22/2019	

Labeler - Taro Pharmaceuticals U.S.A., Inc. (145186370)

Establishment			
Name	Address	ID/FEI	Business Operations
Taro Pharmaceutical Industries Ltd.		600072078	MANUFACTURE(51672-2129)

Revised: 2/2023

Taro Pharmaceuticals U.S.A., Inc.