

LEVOCETIRIZINE DIHYDROCHLORIDE- levocetirizine dihydrochloride tablet, coated
Walmart Inc.

Drug Facts

Active ingredient (in each tablet)

Levocetirizine dihydrochloride USP, 5 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings

Do not use

- if you have kidney disease
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have

- ever had trouble urinating or emptying your bladder

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask doctor if

- you have trouble urinating or emptying your bladder
- an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

- if breast-feeding: not recommended

- if pregnant: ask a health professional before use

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

| | |
|--|--|
| adults 65 years of age and older | <ul style="list-style-type: none"> • ask a doctor |
| adults and children 12-64 years of age | <ul style="list-style-type: none"> • take 1 tablet (5 mg) once daily in the evening • do not take more than 1 tablet (5 mg) in 24 hours • 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms |
| children 6-11 years of age | <ul style="list-style-type: none"> • take 1/2 tablet (2.5 mg) once daily in the evening • do not take more than 1/2 tablet (2.5 mg) in 24 hours |
| children under 6 years of age | <ul style="list-style-type: none"> • do not use |
| consumers with kidney disease | <ul style="list-style-type: none"> • do not use |

Other information

- store between 20° and 25°C (68° and 77°F)
- safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients

colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

Questions or comments?

Call **1-888-375-3784**

Carton Label

Carton Label:

equate™

Allergy Relief

Levocetirizine
Dihydrochloride
Tablets USP, 5 mg

Antihistamine

ORIGINAL PRESCRIPTION STRENGTH



equate™

- Clinically proven relief in 60 minutes
- Clinically proven relief for 24 hours



Actual Size

equate™

NDC 49305-492-35



Allergy Relief

Levocetirizine
Dihydrochloride
Tablets USP, 5 mg

Antihistamine

ORIGINAL PRESCRIPTION STRENGTH

- 24 Hour Relief of:
- Sneezing
 - Runny nose
 - Itchy nose or throat
 - Itchy, watery eyes



Actual Size

5 mg

35 TABLETS

Drug Facts

Active ingredient (in each tablet) Purpose
Levocetirizine dihydrochloride USP 5 mg.....Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other respiratory allergies:
 ■ runny nose ■ itchy, watery eyes
 ■ sneezing ■ itching of the nose or throat

Warnings
Do not use ■ if you have kidney disease
 ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have
 ■ ever had trouble urinating or emptying your bladder

When using this product ■ drowsiness may occur
 ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if
 ■ you have trouble urinating or emptying your bladder
 ■ an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:
 ■ if breast-feeding, not recommended
 ■ if pregnant: ask a health professional before use
Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

| | |
|--|---|
| adults 65 years of age and older | ask a doctor |
| adults and children 12-64 years of age | take 1 tablet (5 mg) once daily in the evening do not take more than 1 tablet (5 mg) in 24 hours |
| children 6-11 years of age | take 1/2 tablet (2.5 mg) once daily in the evening do not take more than 1/2 tablet (2.5 mg) in 24 hours |
| children under 6 years of age | do not use |
| consumers with kidney disease | do not use |

Drug Facts (continued)

Other information

- store between 20° and 25°C (68° and 77°F)
- safety sealed; do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

Questions or comments? 1-888-287-1915

Read directions and warnings before use. Keep this carton, it has important information.



Revised: 12/24

Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments or to report an undesired reaction or side effect, please call 1-888-287-1915.

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716
PRODUCT OF INDIA
 *This product is not manufactured or distributed by Chattem, Inc. (part of the Sanofi Group), distributor of Xyzal® Allergy 24HR Tablets. Xyzal® is a registered trademark of UCB Biopharma SRL.



LOT

EXP



Container Label

equate™ NDC 49035-492-35

Allergy Relief
Levocetirizine Dihydrochloride Tablets USP, 5 mg Antihistamine
ORIGINAL PRESCRIPTION STRENGTH

24 HOUR

5 mg 35 TABLETS

Active ingredient (in each tablet) Purpose
Levocetirizine dihydrochloride USP, 5 mg.....Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings Do not use ■ if you have kidney disease ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine **Ask a doctor before use if you have** ■ ever had trouble urinating or emptying your bladder

When using this product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if ■ you have trouble urinating or

(Continued On Back Of Label)

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915.

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716 67363
REV 09/22
150090207

LOT
EXP
PEEL HERE

emptying your bladder ■ an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding:** ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions **adults 65 years of age and older** ■ ask a doctor **adults and children 12-64 years of age** ■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours ■ 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms **children 6-11 years of age** ■ take 1/2 tablet (2.5 mg) once daily in the evening ■ do not take more than 1/2 tablet (2.5 mg) in 24 hours **children under 6 years of age** ■ do not use **consumers with kidney disease** ■ do not use

Other information ■ store between 20° and 25°C (68° and 77°F) ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

Questions and comments? call 1-888-287-1915

LEVOCETIRIZINE DIHYDROCHLORIDE

levocetirizine dihydrochloride tablet, coated

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|------------------------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:49035-492(NDC:43598-735) |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|--------------------------------|----------|
| levocetirizine dihydrochloride (UNII: SOD6A38AGA) (levocetirizine - UNII:6U5EA9RT2O) | levocetirizine dihydrochloride | 5 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) | |
| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| HYPROMELLOSES (UNII: 3NXW29V3WO) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | white | Score | 2 pieces |
| Shape | OVAL | Size | 9mm |
| Flavor | | Imprint Code | L |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:49035-492-35 | 1 in 1 BOTTLE | 07/26/2019 | |
| 1 | | 35 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 2 | NDC:49035-492-79 | 2 in 1 BLISTER PACK | 07/26/2019 | |
| 2 | | 5 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |
| 3 | NDC:49035-492-10 | 1 in 1 CARTON | 01/15/2026 | |
| 3 | | 10 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 4 | NDC:49035-492-12 | 1 in 1 CARTON | 01/15/2026 | |
| 4 | | 120 in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA210375 | 07/26/2019 | |

Labeler - Walmart Inc. (051957769)

Revised: 1/2026

Walmart Inc.