

**COLD AND ALLERGY D- pseudoephedrine hcl, triprolidine hcl tablet, film coated**  
**Walgreen Company**

-----  
**Walgreens 44-287**

***Active ingredients (in each tablet)***

Pseudoephedrine HCl 60 mg  
Triprolidine HCl 2.5 mg

***Purpose***

Nasal decongestant  
Antihistamine

***Uses***

- temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:
  - runny nose
  - itchy, watery eyes
  - nasal congestion
  - sneezing
  - itching of the nose or throat
- temporarily relieves these symptoms due to the common cold:
  - runny nose
  - sneezing
  - nasal congestion

***Warnings***

**Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- high blood pressure
- heart disease
- thyroid disease
- diabetes
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

## **Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers.

## **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness

## **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with fever

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

## ***Directions***

- **do not take more than directed**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours. Do not take more than 4 tablets in 24 hours.
- children under 12 years: do not use

## ***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

## ***Inactive ingredients***

hypromellose, magnesium stearate, polyethylene glycol, silicified microcrystalline cellulose, titanium dioxide

## ***Questions or comments?***

**1-800-426-9391**

## **Principal display panel**

NDC 0363-0287-22

**Walgreens**

• WALGREENS •  
PHARMACIST RECOMMENDED†

**Cold & Allergy D**

PSEUDOEPHEDRINE HCl / NASAL DECONGESTANT  
TRIPROLIDINE HCl / ANTIHISTAMINE

Maximum Strength

- Relieves runny nose, nasal congestion,  
sneezing & itchy, watery eyes

**48**

TABLETS

ONE TABLET PER DOSE

ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

†Our pharmacists recommend  
the Walgreens brand.

50844 ORG112328722

DISTRIBUTED BY:

**WALGREEN CO.,  
DEERFIELD, IL 60015**

**walgreens.com** ©2022 Walgreen Co.

W30RG1022-F  
REV-0124



DISTRIBUTED BY:  
WALGREENS CO.,  
DEERFIELD, IL 60015  
walgreens.com ©2022 Walgreen Co.

Questions or comments? 1-800-426-9391

**Inactive ingredients**  
hypromellose, magnesium stearate, polyethylene glycol, stifted microcrystalline cellulose, titanium dioxide

**Other information**  
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED  
■ OR BLISTER IS TORN OR BROKEN  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number

**Directions**  
■ do not take more than directed  
■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours. Do not take more than 4 tablets in 24 hours.  
■ children under 12 years: do not use

**Stop use and ask a doctor if**  
■ nervousness, dizziness, or sleeplessness occur  
■ symptoms do not improve within 7 days or occur with fever  
■ If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts (continued)**  
■ drowsiness may occur ■ avoid alcoholic beverages  
■ excitability may occur, especially in children  
■ use caution when driving a motor vehicle or operating machinery  
■ alcohol, sedatives, and tranquilizers may increase drowsiness  
Stop use and ask a doctor if

When using this product  
■ do not exceed recommended dosage  
or tranquilizers.

Ask a doctor or pharmacist before use if you are taking sedatives  
■ difficulty in urination due to enlargement of the prostate gland  
■ a breathing problem such as emphysema or chronic bronchitis

Ask a doctor before use if you have  
■ high blood pressure  
■ heart disease ■ thyroid disease ■ diabetes ■ glaucoma  
■ MAOI, ask a doctor or pharmacist before taking this product.  
MAOI drug; if you do not know if your prescription drug contains an MAOI drug, or Parkinson's disease), or for 2 weeks after stopping the inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the

**Warnings**  
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug; if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
■ temporarily relieves these symptoms due to hay fever  
■ temporarily relieves these symptoms due to the common cold.

**Uses**  
■ temporarily relieves these symptoms due to the common cold.  
■ runny nose ■ sneezing ■ nasal congestion  
■ temporarily relieves these symptoms due to the common cold.  
■ runny nose ■ sneezing ■ nasal congestion  
■ temporarily relieves these symptoms due to hay fever  
■ temporarily relieves these symptoms due to the common cold.

**Purpose**  
Pseudoephedrine HCl 60 mg... Nasal decongestant  
Triprolidine HCl 2.5 mg... Antihistamine

**Drug Facts**  
COMPLETE PRODUCT INFORMATION  
KEEP OUTER PACKAGE FOR

Our pharmacists recommend the Walgreens brand.

50844 ORG112328722

No print area  
Lot no/Exp date



TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

B-2201-287A-22-HR  
ORG112328722

NDC 0363-0287-22

Walgreens



# Cold & Allergy D

PSEUDOEPHEDRINE HCl / NASAL DECONGESTANT  
TRIPROLIDINE HCl / ANTIHISTAMINE

Maximum Strength

- Relieves runny nose, nasal congestion, sneezing & itchy, watery eyes

48

TABLETS

ONE TABLET PER DOSE

ACTUAL SIZE

Walgreens 44-287A

## COLD AND ALLERGY D

pseudoephedrine hcl, triprolidine hcl tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-0287
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	60 mg
<b>TRIPROLIDINE HYDROCHLORIDE</b> (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	2.5 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	287
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0363-0287-22	2 in 1 CARTON	11/29/2022	
1		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M012	11/29/2022	

**Labeler** - Walgreen Company (008965063)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		038154464	pack(0363-0287)

## **Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867837	manufacture(0363-0287) , pack(0363-0287)

## **Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867894	manufacture(0363-0287)

## **Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		117025878	manufacture(0363-0287)

Revised: 2/2026

Walgreen Company