

WALGREENS INSTANT TOOTHACHE AND GUM RELIEF TRIPLE MEDICATED- benzocaine, menthol, zinc chloride gel

Walgreens

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Walgreens Instant Toothache & Gum Relief Gel Triple Med

Drug Facts

Active ingredients

Benzocaine 20%

Menthol 0.5%

Zinc Chloride 0.15%

Purpose

Oral pain reliever

Oral pain reliever

Astringent

Use

- for the temporary relief of oral pain due to minor irritation or injury of the mouth and gums

Warnings

Methemoglobinemia warning: use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics

Do not use

- more than directed
- for more than 7 days unless directed by a dentist or doctor
- for teething
- in children under 2 years of age

Stop use and ask a doctor if

- swelling, rash or fever develops
- irritation, pain or redness persists or worsens
- symptoms do not improve in 7 days
- allergic reaction occurs

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- cut open tip of tube on score mark
- adults and children 2 years of age and older: apply a small amount of the product to the cavity and around the gum surrounding the teeth. Use up to 4 times daily or as directed by a dentist or doctor.
- children between 2 and 12 years of age should be supervised in the use of this product
- children under 2 years of age: do not use

Other information

- do not use if tip is cut prior to opening
- this preparation is intended for use in cases of toothache, only as a temporary expedient until a dentist can be consulted
- do not use continuously
- this formula will stay in place for extended duration of relief
- avoid using toothpaste or drinking soft drinks or fruit juices for at least one hour after applying

Inactive ingredients blue 1, cellulose gum, gelatin, methyl salicylate, mineral oil, pectin, petrolatum, polyethylene glycol, sodium saccharin

Questions or comments? 1-800-925-4733

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com

Walgreens

Instant Toothache & Gum Relief

ORAL PAIN RELIEVER / ASTRINGENT

SEVERE **GEL TRIPLE MEDICATED**

- 20% maximum strength benzocaine for to relieve oral pain
- Astringent to relieve gum irritation
- Menthol to soothe

NET WT 0.25 OZ (7.0 g)

Walgreens

Compare to Orajel™ Severe
Toothache & Gum Relief Plus Triple
Medicated Gel active ingredients!

Instant Toothache & Gum Relief

ORAL PAIN RELIEVER / ASTRINGENT

SEVERE GEL TRIPLE MEDICATED

- 20% maximum strength benzocaine to relieve oral pain
- Astringent to relieve gum irritation
- Menthol to soothe

Fast-acting oral pain relief

Walgreens

Instant Toothache & Gum Relief

ORAL PAIN RELIEVER / ASTRINGENT NET WT 0.25 OZ (7 g)

Questions or comments? 1-800-925-4733

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2018 Walgreen Co.

ITEM 757551 W00000-0000-0



Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey
This product is not manufactured or distributed by Church & Dwight Co., Inc., makers of Orajel™.

Drug Facts

Active ingredients

Benzocaine 20%
Menthol 0.5%
Zinc Oxide 0.15%
Astringent

Purpose

Oral pain reliever
Oral pain reliever
Astringent

Warnings

Medicinal benzocaine warning: use of this product may cause methemoglobinemia, a serious condition that may be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used the product before. Stop use and seek immediate medical attention if you or a child in your care develops:
• pale, gray, or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy

Allergy info: do not use this product if you have a history of allergy to local anesthetics such as procaine, butabupivacaine, or other "caine" anesthetics.

Do not use: • more than directed • for more than 7 days unless told to do so by a dentist or doctor • for bleeding • in children under 2 years of age

Stop use and ask a doctor if: • swelling, rash or fever develops • irritation, pain or redness persists or worsens • symptoms do not improve in 7 days • allergic reaction occurs

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • Cut open tip of tube on score mark • Adults and children 2 years of age and older: apply a small amount of product to the cavity and around the gum surrounding the tooth. Use up to 4 times daily or as directed by a dentist or doctor • Children between 2 and 12 years of age should be supervised in the use of this product. • Children under 2 years of age: do not use.

Other information • Do not use if tip is cut prior to opening • This preparation is intended for use in cases of toothache, only as a temporary expedient until a dentist can be consulted • Do not use continuously • Use for 10 minutes only • Do not use if you are allergic to any of the ingredients • Do not use if you are pregnant or breastfeeding • Do not use if you are taking any other medicine • Do not use if you are taking any other medicine • Do not use if you are taking any other medicine

Inactive ingredients Benzocaine, zinc oxide, menthol, methyl salicylate, mineral oil, polyethylene glycol, sodium ascorbate, polyethylene glycol, sodium ascorbate.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

TAMPER EVIDENT: DO NOT USE IF TUBE TIP IS CUT PRIOR TO USE

WALGREENS INSTANT TOOTHACHE AND GUM RELIEF TRIPLE MEDICATED

benzocaine, menthol, zinc chloride gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-5351
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ZINC CHLORIDE (UNII: 86Q357L16B) (ZINC CATION - UNII:13S1S8SF37)	ZINC CATION	0.15 g in 100 g
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	0.5 g in 100 g
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	20 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
PETROLATUM (UNII: 4T6H12BN9U)	
MINERAL OIL (UNII: T5L8T28FGP)	
PECTIN (UNII: 89NA02M4RX)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
METHYL SALICYLATE (UNII: LAV5U5022Y)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	blue	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-5351-09	1 in 1 CARTON	03/06/2019	
1		7 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	03/06/2019	

Labeler - Walgreens (008965063)

Registrant - Lornamead Inc. (080046418)

Establishment

Name	Address	ID/FEI	Business Operations
Lornamead Inc.		080046418	manufacture(0363-5351)

Revised: 1/2023

Walgreens