

FRITZ AQUATICS MARDEL MARACYN TWO- minocycline powder

Fritz Industries


Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

FRITZ - Maracyn Two


ACTIVE INGREDIENT: Each packet contains 10 mg Minocycline activity.

TREATMENT
LOTT
FIN & TAIL ROT
POPEYE
DROOPY


For expert advice on the use of this product scan the QR code




Common Symptoms of Aeromonas and Flavobacterium infection.




Septicemia: Bleeding or red streaks on the body; ulcers may also be present.




Fin & Tail Rot: Split, ragged and deteriorating fin end/or tail.



Droopy: Swollen, labored body, protruding scales and/or clamped fins.



Pop-eye: Protruding eyes, may be cloudy or hazy.



Gill Disease: Swollen, discolored gills, gasping for air and decreased activity.

INDICATIONS: Use of a hospital tank is recommended. No water changes, pH or temperature adjustments necessary if readings are in ideal range (Freshwater pH 6.4 - 7.6 / Saltwater 8.0 - 8.4, temperature 68° to 86° F). Remove all chemical filtration (carbon, ion exchange resins) before and during use. Aeration and water flow should be maintained. In marine aquariums turn off protein skimmer during treatment; resume use after completion of treatment. Do not perform water changes during treatment if possible. Antibiotics can negatively affect beneficial bacteria which remove ammonia and nitrite from water. Testing for ammonia and nitrite is recommended. Chemical ammonia removers such as Fritz Aquatics A.C.C.R. may be used during treatment to prevent ammonia poisoning of the fish.
Note: Maracyn Two can cause water to appear cloudy and/or will have floating particles.

DIRECTIONS: One complete course of treatment will take 5 days. Treat for the full course of treatment even if visible signs of infection or illness disappear.

Day One: Add contents of two packets per 10 gallons of water to be treated.

Day Two through Day Five: Add contents of one packet per 10 gallons of water to be treated daily.


Day Six: Perform a 25% water change, if necessary, a second full course of treatment may be performed starting on Day Seven.

For treatment directions in unfiltered small aquariums, betta bowls and containers of less than 10 gallons please visit www.fritzaquatics.com and read instructions under FAQs for Maracyn.

ACTIVE INGREDIENT: Each packet contains 10 mg of Minocycline activity.

FRITZ

Item# 47001



0 80531 47001 5

MADE IN USA

www.fritzaquatics.com

Questions or Comments?

Email: info@fritzaquatics.com

FRITZ

MADE IN USA

www.fritzaquatics.com

Questions or Comments?

Email: info@fritzaquatics.com

FRITZ

FRITZ



PROVIDES RELIEF FROM BACTERIAL INFECTIONS

Maracyn Two

Fast Dissolving Powder Form

10 PACK

Net Contents: 10 - 0.035 oz (1 g) Powder Packets

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. For aquarium use only. WARNING: THIS PRODUCT IS INTENDED FOR THE EXCLUSIVE USE WITH ORNAMENTAL ORGANISMS INDICATED AND IS NOT INTENDED FOR USE WITH HUMANS OR PETS FOR HUMAN CONSUMPTION.

FRITZ AQUATICS
DROPSY POPEYE FIN & TAIL ROT
TREATMENT
FRESHWATER & SALTWATER

For expert advice on the use of this product scan the QR code.

Common Symptoms of Aeromonas and Flavobacterium infection.

Sepsicemia: Bleeding or red streaks on the body colors may also be present.

Fin & Tail Rot: Soft, ragged and deteriorating fin and/or tail.

Dropsy: Swollen, bloated body protruding scales and/or changed fins.

Popeye: Protruding eyes, may be cloudy or hazy.

GI Disease: Swollen, distended gill, gasping for air and decreased activity.

INDICATIONS: Use of a hospital tank is recommended. No water changes, pH or temperature adjustments necessary if readings are in ideal range (Freshwater pH 6.4 - 7.6 / Saltwater 8.0 - 8.4, temperature 68° to 85° F). Remove all chemical filtration (carbon, ion exchange resins) before and during use. Aeration and water flow should be maintained. In marine aquariums turn off protein skimmer during treatment; resume use after completion of treatment. Do not perform water changes during treatment if possible. Antibiotics can negatively affect beneficial bacteria which remove ammonia and nitrite from water. Testing for ammonia and nitrite is recommended. Chemical ammonia removers such as Fritz Aquatics A.C.C.R. may be used during treatment to prevent ammonia poisoning of the fish.
Note: Maracyn Two can cause water to appear cloudy and/or will have floating particles.

DIRECTIONS: One complete course of treatment will take 5 days. Treat for the full course of treatment even if visible signs of infection or illness disappear.
Day One: Add contents of two packets per 10 gallons of water to be treated.
Day Two through Day Five: Add contents of one packet per 10 gallons of water to be treated daily.
Day Six: Perform a 25% water change. If necessary, a second full course of treatment may be performed starting on Day Seven.

For treatment directions in unfiltered small aquariums, beta bowls and containers of less than 10 gallons please visit www.fritzaquatics.com read instructions under FAQs for Maracyn.

ACTIVE INGREDIENT: Each packet contains 10 mg of Minocycline activity.

FRITZ AQUATICS
Distributed by: Fritz Industries, Inc.
Mesquite, TX 75149
MADE IN USA
www.fritzaquatics.com
Questions or Comments?
Email: info@fritzaquatics.com

Item # 47002
0 805314 7002 2

FRITZ
Maracyn Two

PROVIDES RELIEF FROM BACTERIAL INFECTIONS

For the treatment and control of septicemia, bacterial fin and tail rot and cotton wool disease in freshwater and saltwater fish caused by *Aeromonas punctata*, *Aeromonas hydrophila* and *Flavobacterium columnare* bacteria.

Fast Dissolving Powder Form

20 PACK

Net Contents: 20 - 0.035 oz (1 g) Powder Packets

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.
For aquarium use only.
WARNING: THIS PRODUCT IS INTENDED FOR THE EXCLUSIVE USE WITH AQUARIUM FISH. IT IS NOT INTENDED FOR USE WITH HUMANS OR OTHER ANIMAL CONSUMPTION.

FRITZ AQUATICS MARDEL MARACYN TWO

minocycline powder

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:53745-012
Route of Administration	EXTRACORPOREAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MINOCYCLINE (UNII: FYY3R43WGO) (MINOCYCLINE - UNII:FYY3R43WGO)	MINOCYCLINE	10 mg in 0.6 g

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53745-012-08	8 in 1 CARTON		
1		0.6 g in 1 PACKET		
2	NDC:53745-012-24	24 in 1 CARTON		
2		0.6 g in 1 PACKET		
3	NDC:53745-012-10	10 in 1 PACKAGE		
3		0.6 g in 1 PACKET		
4	NDC:53745-012-20	20 in 1 PACKAGE		
4		0.6 g in 1 PACKET		

5	NDC:53745-012-28	280 g in 1 BOTTLE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
unapproved drug other		04/01/2014		

Labeler - Fritz Industries (131030991)

Establishment			
Name	Address	ID/FEI	Business Operations
Curia Italy S.r.l.		440365766	api manufacture

Establishment			
Name	Address	ID/FEI	Business Operations
Sulzbach Enterprises, Inc.		616846528	manufacture

Revised: 1/2024

Fritz Industries