

**MUCUS RELIEF MAXIMUM STRENGTH COLD FLU AND SORE THROAT-
acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid
EQUATE (Wal-Mart Stores, Inc.) (see also WAL-MART INC)**

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - sinus congestion and pressure
 - minor aches and pains
 - nasal congestion
 - headache
 - cough
 - sore throat
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 day
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in a 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- dose as follows or as directed by a doctor
- mL = milliliter
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not use

Other information

- **each 20 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, disodium EDTA, FD&C blue #1, FD&C yellow #10, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

1-888-287-1915

Principal Display Panel

Compare to Maximum Strength Mucinex® Fast-Max® Clear & Cool® Cold, Flu, & Sore Throat active ingredients*

Cold, Flu & Sore Throat

Acetaminophen 650 mg-Pain Reliever / Fever Reducer

Dextromethorphan HBr 20 mg-Cough Suppressant

Guaifenesin 400 mg-Expectorant

Phenylephrine HCl 10 mg-Nasal Decongestant

MAXIMUM STRENGTH

Multi-Symptom Relief

- Relieves headache & fever
- Controls cough
- Relieves nasal & chest congestion
- Thins and loosens mucus

For Ages 12+

Cooling Menthol Flavor

FL OZ (mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Clear & Cool® Cold, Flu & Sore Throat.

DISTRIBUTED BY: Walmart Inc.,

Bentonville, AR 72716

Package Label

equate™

NDC 49035-682-06



Cold, Flu & Sore Throat

Acetaminophen - Pain reliever/Fever reducer
Dextromethorphan HBr - Cough suppressant
Guaifenesin - Expectorant
Phenylephrine HCl - Nasal decongestant

MAXIMUM STRENGTH

Multi-Symptom Relief

- Relieves headache & fever
- Controls cough
- Relieves nasal & chest congestion
- Thins and loosens mucus
- Cooling menthol flavor



6 FL OZ (177mL)

For Ages 12+

PLD-8452B LB006394

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments please call 1-888-287-1915.

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Clear & Cool® Cold, Flu & Sore Throat.

483524



PARENTS:
Learn about teen medicine abuse

www.StopMedicineAbuse.org

PLD-8452B LB006394

PEEL CORNER FOR DRUG FACTS

Drug Facts

Active ingredients (in each 20 mL)

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms
 - sinus congestion and pressure
 - minor aches and pains
 - headache
 - cough
 - nasal congestion
 - sore throat
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

Drug Facts (continued)

- more than 4,000 mg of acetaminophen in 24 hours
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks daily while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
- Do not use**
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- Ask a doctor before use if you have**
- liver disease
 - heart disease
 - diabetes
 - thyroid disease
 - high blood pressure

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
 - persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
 - cough that occurs with too much phlegm (mucus)
- Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**
- When using this product, do not use more than directed.**
- Stop use and ask a doctor if**
- nervousness, dizziness, or sleeplessness occur
 - pain, nasal congestion, or cough gets worse, or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- If pregnant or breast-feeding, ask a health professional before use.**
- Keep out of reach of children.**
- Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for

Drug Facts (continued)

- adults as well as for children even if you do not notice any signs or symptoms.
- Directions**
- do not take more than directed (see Overdose warning)
 - do not take more than 6 doses in any 24-hour period
 - measure only with dosing cup provided. Do not use any other dosing device.
 - keep dosing cup with product
 - mL = milliliter
 - dose as follows or as directed by a doctor
 - adults and children 12 years of age and older: 20 mL every 4 hours
 - children under 12 years of age: do not use
- Other information**
- each 20 mL contains: sodium 12 mg
 - store between 20-25°C (68-77°F). Do not refrigerate.
- Inactive ingredients** citric acid, disodium EDTA, FD&C blue #1, FD&C yellow #10, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum
- Questions or comments?**
1-888-287-1915

PEEL CORNER FOR MORE DRUG FACTS

EQUATE Cold Flu & Sore Throat Maximum Strength

MUCUS RELIEF MAXIMUM STRENGTH COLD FLU AND SORE THROAT

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49035-682
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-682-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/30/2018	04/30/2025

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/30/2018	04/30/2025

Labeler - EQUATE (Wal-Mart Stores, Inc.) (see also WAL-MART INC) (051957769)

Revised: 3/2024

EQUATE (Wal-Mart Stores, Inc.) (see also WAL-MART INC)