

**BENZETHONIUM CHLORIDE AND DYCLONINE HYDROCHLORIDE- liquid  
bandage liquid  
CVS Pharmacy**

-----  
**CVS Health Liquid Bandage**

***Active ingredients***

Benzethonium chloride 0.2%

Dyclonine hydrochloride 0.75%

***Purpose***

First aid antiseptic

Topical analgesic

***Uses***

- first aid to prevent bacterial contamination in minor cuts, scrapes and burns
- for the temporary relief of pain

***Warnings***

**For external use only.**

**Flammable:**

Keep away from fire or flame.

**Do not use**

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

**Ask a doctor before use if you have**

- deep or puncture wounds
- deep cuts
- animal bites
- serious burns

**When using this product**

- avoid contact with eyes.

**Stop use and consult a doctor if**

- condition persists or gets worse
- symptoms persist for more than 7 days or clear up and occur again within a few days

- infection occurs

**Keep out of reach of children.**

If swallowed, get medical help or call a Poison Control Center right away.

***Directions***

- for adults and children 2 years of age and older
- clean affected area
- apply a small amount on the area 1 to 3 times daily and let dry
- a second coating may be applied for extra protection
- children under 2 years of age: consult a doctor

***Other information***

- cap tightly
- store between 20° to 25° C (68° to 77°F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off

***Inactive ingredient***

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

***Questions?***

**Call 1-866-964-0939**

**Principal Display Panel**

CVS Health

**Liquid Bandage**

**FOR SMALL CUTS & WOUNDS**

TOPICAL ANALGESIC & ANTISEPTIC SEALS TO PROTECT

- Prevent infection
- Invisible & flexible
- Easy to use

Actual Bottle Size on Side Panel

1 FL OZ (29.6 mL)

PRINT SIDE GLUE

CVS Health.

# Liquid Bandage

FOR SMALL CUTS & WOUNDS

TOPICAL ANALGESIC & ANTISEPTIC SEALS TO PROTECT

VALUE SIZE

CVS Health.

Compare to New-Skin<sup>®</sup>

NDC 69842-084-30

# Liquid Bandage

FOR SMALL CUTS & WOUNDS

TOPICAL ANALGESIC & ANTISEPTIC SEALS TO PROTECT

- Prevents infection
- Invisible & flexible
- Easy to use



Actual Bottle Size on Side Panel



1 FL OZ (29.6 mL)

Use CVS Health<sup>™</sup> Liquid Bandage for:

- Paper cuts
- Blisters
- Chapped & cracked fingertips
- Shaving nicks
- Hangnails
- To help prevent the formation of calluses

Used by:

Bowlers, golfers, tennis players, fishermen, musicians, runners, hikers and dancers

Package Contains One Bottle

Actual Size

DO NOT USE IF NECKBAND IS BROKEN

### Drug Facts

**Active ingredients Purpose**  
Benzethonium chloride 0.2% First aid antiseptic  
Dyclonine hydrochloride 0.75% Topical analgesic

**Uses** ■ first aid to prevent bacterial contamination in minor cuts, scrapes and burns ■ for the temporary relief of pain

### Warnings

For external use only.

**Flammable:** Keep away from fire or flame.

**Do not use**

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

**Ask a doctor before use if you have**

- deep or puncture wounds ■ deep cuts
- animal bites ■ serious burns

**When using this product**

- avoid contact with eyes.

**Stop use and consult a doctor if**

- condition persists or gets worse
- symptoms persist for more than 7 days or clear up and occur again within a few days
- infection occurs

**Keep out of reach of children.**

If swallowed, get medical help or call a Poison Control Center right away.

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2019 CVS/pharmacy  
CVS.com<sup>®</sup> 1-800-SHOP CVS  
Made in the U.S.A. with U.S. and imported parts  
V-16372



### Drug Facts (continued)

#### Directions

- for adults and children 2 years of age and older
- clean affected area
- apply a small amount on the area 1 to 3 times daily and let dry
- a second coating may be applied for extra protection
- children under 2 years of age: consult a doctor

#### Other information

- cap bottle tightly
- store between 20° to 25°C (68° to 77°F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off

#### Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

**Questions? Call 1-866-864-0939**

Care: Wipe top of bottle clean after each use.

\*This product is not manufactured or distributed by Moberg Pharma North America LLC, owner of the registered trademark New-Skin<sup>®</sup>.

#369239

FPO 80%  
UPC# 050428616710

X XXXXXX XXXXXX X

53-033CV-B01

NO VARNISH  
AREA  
(WHITE ONLY)

DO NOT USE IF NECKBAND IS BROKEN



NDC 69842-084-30

**Active Ingredients:** Benzethonium Chloride 0.2% and Dyclonine Hydrochloride 0.75%. **Purpose:** First aid antiseptic and topical analgesic. **Uses:** First aid to prevent bacterial contamination in minor cuts, scrapes, and burns. For the temporary relief of pain.

**Warnings:** For external use only. **Flammable:** Keep away from fire or flame. Do not use in eyes, over large areas of the body, and longer than 1 week unless directed by a doctor.

## Liquid Bandage

FOR SMALL CUTS & WOUNDS

TOPICAL ANALGESIC & ANTISEPTIC SEALS TO PROTECT

**Ask a doctor before use if you have** deep or puncture wounds, deep cuts, animal bites and serious burns.

**When using this product** avoid contact with eyes.

**Stop use and ask a doctor** if condition persists or gets worse, symptoms persist for more than 7 days or clear up and occur again within a few days, and infection occurs.

**Keep out of reach of children.** If swallowed, get medical help or call a Poison Control Center right away.

**Directions:** For adults and children 2 years of age and older. Clean affected area. Apply a small amount on the area 1 to 3 times daily and let dry. A second coating may be applied for extra protection. Children under 2 years of age: consult a doctor.

**Other information:** Cap bottle tightly. Store between 20° to 25°C (68° to 77°F). May temporarily sting upon application. To remove, apply more Liquid Bandage and quickly wipe off.

**Inactive ingredients:** acetone, amyli acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40


**Questions? Call 1-866-964-0939**

**Care:** Wipe top of bottle clean after each use.

**Distributed by:** CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2019 CVS/pharmacy  
CVS.com® 1-800-SHOP CVS  
Made in the U.S.A. with U.S. and imported parts  
V-16372

1 FL OZ  
(29.6 mL)





50-033CV-B01

## BENZETHONIUM CHLORIDE AND DYCLONINE HYDROCHLORIDE

liquid bandage liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69842-084
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZETHONIUM CHLORIDE</b> (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 10 mL
<b>DYCLONINE HYDROCHLORIDE</b> (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q300)	DYCLONINE HYDROCHLORIDE	0.75 mg in 10 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>PYROXYLIN</b> (UNII: KYR8BR2X6O)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>CASTOR OIL</b> (UNII: D5340Y2I9G)	
<b>ACETONE</b> (UNII: 1364PS73AF)	
<b>AMYL ACETATE</b> (UNII: 92Q24NH7AS)	
<b>ETHYL ACETATE</b> (UNII: 76845O8NMZ)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date

<b>1</b>	NDC:69842-084-30	10 mL in 1 BOTTLE, WITH APPLICATOR; Type 0: Not a Combination Product	06/01/2019	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M003		06/01/2019	

**Labeler -** CVS Pharmacy (062312574)

Revised: 11/2024

CVS Pharmacy