

RIMMEL LASTING BREATHABLE FOUNDATION SPF 20- octinoxate liquid
Rimmel Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rimmel Lasting Finish Breathable Foundation SPF 20

Active ingredients

Octinoxate 4%

Purpose

Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Do not use

on damaged or broken skin.

When using this product

keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor

if rash occurs.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply at least every 2 hours
- use a water resistant sunscreen if swimming or sweating
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including: [bullet] limit time in the sun, especially from 10 a.m.-2 p.m. [bullet] wear long-sleeved shirts, pants, hats, and sunglasses.

- Children under 6 months of age: ask a doctor

Other information

Protect the product in this container from excessive heat and direct sun.

Inactive ingredients

WATER, CYCLOPENTASILOXANE, TALC, ISODODECANE, GLYCERIN, VINYL DIMETHICONE/METHICONE SILSESQUIOXANE CROSSPOLYMER, PHENYL TRIMETHICONE, PEG-10 DIMETHICONE, TRIMETHYLSILOXYSILICATE, SODIUM CHLORIDE, AMMONIUM ACRYLATES COPOLYMER, PHENOXYETHANOL, DISTEARDIMONIUM HECTORITE, PENTYLENE GLYCOL, TRIETHOXYCAPRYLYLSILANE, CHLORPHENESIN, FRAGRANCE, ALCOHOL DENAT., ASCORBYL PALMITATE, CITRIC ACID, FRUCTOSE, PROPYLENE CARBONATE, SODIUM HYDROXIDE, TOCOPHERYL ACETATE, UREA, RETINYL PALMITATE, DISODIUM DECETH-6 SULFOSUCCINATE, DISODIUM EDTA, LAURETH-30, ALLANTOIN, MALTOSE, SODIUM LACTATE, SODIUM PCA, TREHALOSE, SODIUM DEHYDROACETATE, BHT, GLUCOSE, SODIUM HYALURONATE, TOCOPHEROL, [May Contain: TITANIUM DIOXIDE, IRON OXIDES]

Questions or comments?

Call 1-800-715-4023

9:00 am to 5:00 pm ET

Monday - Friday

RIMMEL

LONDON

LASTING

FINISH 25 HR

BREATHABLE

LONG WEAR FOUNDATION

HIGH DEFINITION FINISH

MEDIUM COVERAGE

BROAD SPECTRUM

SPF 20 SUNSCREEN

1 fl oz 30 ml



Shake before use. Lightweight skin perfecting foundation with medium coverage. Breathable skin technology that moisturizes for up to 25H. Broad Spectrum SPF 20 Sunscreen.

DIST. RIMMEL INC.
NEW YORK, N.Y. 10118
MADE IN MONACO

BAR CODE AREA
23 x 10mm

Drug Facts

Active ingredients	Purpose
Octinoxate 4%	Sunscreen ▶

Peel back for more information ▶

47,5

46,5

Drug Facts (continued)

Uses • helps prevent sunburn
• if used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If swallowed, get medical help ▶

Drug Facts (continued)

or contact a Poison Control Center right away.

Directions For sunscreen use:
• apply liberally 15 minutes before sun exposure • reapply at least every 2 hours • use a water resistant sunscreen if swimming or sweating • **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum value of 15 SPF or higher and other sun protection measures including: • limit time in the sun, especially from 10 a.m. – 2 p.m. • wear long-sleeved shirts, ▶

46,5

Drug Facts (continued)

pants, hats, and sunglasses
• Children under 6 months of age: ask a doctor

Other information Protect the product in this container from excessive heat and direct sun.

Inactive ingredients

water, cyclopentasiloxane, talc, isododecane, glycerin, vinyl dimethicone/methicone silsesquioxane crosspolymer, phenyl trimethicone, PEG-10 dimethicone, trimethylsiloxysilicate, sodium chloride, ammonium acrylates copolymer, phenoxylethanol, disteardimonium hectorite, pentylene glycol. ▶

46,5

Drug Facts (continued)

triethoxycaprylylsilane, chlorophenesin, fragrance, alcohol denat., ascorbyl palmitate, citric acid, fructose, propylene carbonate, sodium hydroxide, tocopheryl acetate, ursae, retinyl palmitate, disodium EDTA, laureth-30, allantoin, mallose, sodium lactate, sodium PCA, trehalose, sodium dehydroacetate, BHT, glucose, sodium hyaluronate, tocopherol. (may contain: titanium dioxide, iron oxides).

Questions or comments?

Call 1-800-715-4023
9:00 am to 5:00 pm ET
Monday – Friday





RIMMEL LASTING BREATHABLE FOUNDATION SPF 20

octinoxate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76485-1111
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	4 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ISODODECANE (UNII: A8289P68Y2)	
GLYCERIN (UNII: PDC6A3C0OX)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
TALC (UNII: 7SEV7J4R1U)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
PHENOXYETHANOL (UNII: HE492ZZ3T)	
CHLORPHENESIN (UNII: I670DAL4SZ)	
ASCORBYL PALMITATE (UNII: QN83US2B0N)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM LACTATE (UNII: TU7HW0W0QT)	

PROPYLENE CARBONATE (UNII: 8D08K3S51E)	
TRIMETHYLSILOXYSILICATE (M/Q 0.6-0.8) (UNII: 5041RX63GN)	
SODIUM DEHYDROACETATE (UNII: 8W46YN971G)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
WATER (UNII: 059QF0KO0R)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
PENTYLENE GLYCOL (UNII: 50C1307PZG)	
ALLANTOIN (UNII: 344S277G0Z)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76485-1111-0	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2019	
2	NDC:76485-1111-9	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2019	
3	NDC:76485-1111-1	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2019	
4	NDC:76485-1111-2	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2019	
5	NDC:76485-1111-4	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2019	
6	NDC:76485-1111-5	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2019	
7	NDC:76485-1111-6	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2019	
8	NDC:76485-1111-7	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2019	
9	NDC:76485-1111-8	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2019	
10	NDC:76485-1111-3	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2019	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	03/15/2019	

Labeler - Rimmel Inc. (401011325)

Establishment			
Name	Address	ID/FEI	Business Operations
Coty Lancaster S.A.M.		401011325	manufacture(76485-1111)