

**METRONIDAZOLE- metronidazole injection, solution**  
**Amneal Pharmaceuticals Private Limited**

-----  
**Metronidazole Injection, USP**  
**in Plastic Container**  
**500 mg/100 mL (5 mg/mL)**  
**Rx only**


To reduce the development of drug-resistant bacteria and maintain the effectiveness of metronidazole injection and other antibacterial drugs, metronidazole injection should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

**WARNING**

Metronidazole has been shown to be carcinogenic in mice and rats (see **PRECAUTIONS**). Unnecessary use of the drug should be avoided. Its use should be reserved for the conditions described in the **INDICATIONS AND USAGE** section below.

**DESCRIPTION**

Metronidazole injection, USP, is a parenteral formulation of the synthetic nitroimidazole antibacterial agent 2-methyl-5-nitro-1*H*-imidazole-1-ethanol.



**Metronidazole, USP**

Metronidazole injection USP, 500 mg/100 mL (5 mg/mL) is a sterile, clear, colorless to pale yellow color, nonpyrogenic, iso-osmotic, buffered solution supplied in 100 mL single-dose non-PVC plastic container. Each 100 mL contains 500 mg metronidazole, USP; 790 mg sodium chloride, USP; 47.6 mg dibasic sodium phosphate dried, USP; and 22.9 mg citric acid anhydrous, USP.

Metronidazole injection, USP has an osmolarity of 308.66 mOsmol/L (calc) and a pH of 5.5 (4.5 to 7.0). Each container contains 14 mEq of sodium.

## CLINICAL PHARMACOLOGY

In patients treated with intravenous metronidazole, using a dosage regimen of 15 mg/kg loading dose followed 6 hours later by 7.5 mg/kg every 6 hours, the average peak steady-state plasma concentrations ( $C_{max}$ ) and trough concentrations ( $C_{min}$ ) were 25 mcg/mL and 18 mcg/mL, respectively. Plasma concentrations of metronidazole are proportional to the administered dose. An eight-hour intravenous infusion of 100 mg to 4,000 mg of metronidazole in normal subjects showed a linear relationship between dose and peak plasma concentration. The average elimination half-life of metronidazole in healthy subjects is eight hours.

### Distribution

Metronidazole is the major component appearing in the plasma, with lesser quantities of metabolites also being present. Less than 20% of the circulating metronidazole is bound to plasma proteins. Metronidazole appears in cerebrospinal fluid, saliva and breast milk in concentrations similar to those found in plasma. Bactericidal concentrations of metronidazole have also been detected in pus from hepatic abscesses.

Following a single intravenous dose of metronidazole 500 mg, 4 healthy subjects who underwent gastrointestinal endoscopy had peak gastric juice metronidazole concentrations of 5 mcg/mL to 6 mcg/mL at one-hour post-dose. In patients receiving intravenous metronidazole in whom gastric secretions are continuously removed by nasogastric aspiration, sufficient metronidazole may be removed in the aspirate to cause a reduction in serum levels.

### Metabolism

The metabolites of metronidazole result primarily from side-chain oxidation [1-( $\beta$ -hydroxyethyl)-2-hydroxymethyl-5-nitroimidazole and 2-methyl-5-nitroimidazole-1-ylacetic acid] and glucuronide conjugation. Both the parent compound and the hydroxyl metabolite possess *in vitro* antimicrobial activity.

### Excretion

The major route of elimination of metronidazole and its metabolites is via the urine (60% to 80% of the dose), with approximately 20% of the amount excreted appearing as unchanged metronidazole. Renal clearance of metronidazole is approximately 10 mL/min/1.73 m<sup>2</sup>. Fecal excretion accounts for 6% to 15% of the dose.

### Renal Impairment

Decreased renal function does not alter the single-dose pharmacokinetics of metronidazole.

Subjects with end-stage renal disease (ESRD;  $CL_{CR} = 8.1 \text{ mL/min} \pm 9.1 \text{ mL/min}$ ) and who received a single intravenous infusion of metronidazole 500 mg had no significant change in metronidazole pharmacokinetics but had 2-fold higher  $C_{max}$  of hydroxy-metronidazole and 5-fold higher  $C_{max}$  of metronidazole acetate, compared to healthy subjects with normal renal function ( $CL_{CR} = 126 \text{ mL/min} \pm 16 \text{ mL/min}$ ). Thus, on account of the potential accumulation of metronidazole metabolites in ESRD patients, monitoring for metronidazole associated adverse events is recommended (see

**PRECAUTIONS**).

## Effect of Dialysis

Following a single intravenous infusion or oral dose of metronidazole 500 mg, the clearance of metronidazole was investigated in ESRD subjects undergoing hemodialysis or continuous ambulatory peritoneal dialysis (CAPD). A hemodialysis session lasting for 4 hours to 8 hours removed 40% to 65% of the administered metronidazole dose, depending on the type of the dialyzer membrane used and the duration of the dialysis session. If the administration of metronidazole cannot be separated from the dialysis session, supplementation of metronidazole dose following hemodialysis should be considered (see **DOSAGE AND ADMINISTRATION**). A peritoneal dialysis session lasting for 7.5 hours removed approximately 10% of the administered metronidazole dose. No adjustment in metronidazole dose is needed in ESRD patients undergoing CAPD.

## Hepatic Impairment

Following a single intravenous infusion of 500 mg metronidazole, the mean  $AUC_{24}$  of metronidazole was higher by 114% in patients with severe (Child-Pugh C) hepatic impairment, and by 54% and 53% in patients with a mild (Child-Pugh A) and moderate (Child-Pugh B) hepatic impairment, respectively, compared to healthy control subjects. There were no significant changes in the  $AUC_{24}$  of hydroxy-metronidazole in these hepatically impaired patients. A reduction in metronidazole dosage by 50% is recommended in patients with severe (Child-Pugh C) hepatic impairment (see **DOSAGE AND ADMINISTRATION**). No dosage adjustment is needed for patients with mild to moderate hepatic impairment. Patients with mild to moderate hepatic impairment should be monitored for metronidazole associated adverse events (see **PRECAUTIONS** and **DOSAGE AND ADMINISTRATION**).

## Geriatric Patients

Following a single 500 mg oral or intravenous dose of metronidazole, subjects > 70 years old with no apparent renal or hepatic dysfunction had a 40% to 80% higher mean AUC of hydroxy-metronidazole (active metabolite), with no apparent increase in the mean AUC of metronidazole (parent compound), compared to young healthy controls < 40 years old. In geriatric patients, monitoring for metronidazole associated adverse events is recommended (see **PRECAUTIONS**).

## Pediatric Patients

In one study newborn infants appeared to demonstrate diminished capacity to eliminate metronidazole. The elimination half-life, measured during the first three days of life, was inversely related to gestational age. In infants whose gestational ages were between 28 weeks and 40 weeks, the corresponding elimination half-lives ranged from 109 hours to 22.5 hours.

## Microbiology

### Mechanism of Action

Metronidazole, a nitroimidazole, exerts antibacterial effects in an anaerobic environment against most obligate anaerobes. Once metronidazole enters the organism by passive diffusion and is activated in the cytoplasm of susceptible anaerobic bacteria, it is

reduced; this process includes intra-cellular electron transport proteins such as ferredoxin, transfer of an electron to the nitro group of the metronidazole, and formation of a short-lived nitroso free radical. Because of this alteration of the metronidazole molecule, a concentration gradient is created and maintained which promotes the drug's intracellular transport. The reduced form of metronidazole and free radicals can interact with DNA leading to inhibition of DNA synthesis and DNA degradation leading to death of bacteria. The precise mechanism of action of metronidazole is unclear.

## **Drug Resistance**

A potential for development of resistance exists against metronidazole.

Resistance may be due to multiple mechanisms that include decreased uptake of the drug, altered reduction efficiency, overexpression of the efflux pumps, inactivation of the drug and/or increased DNA damage repair.

Metronidazole does not possess any clinically relevant activity against facultative anaerobes or obligate aerobes.

## **Activity *In Vitro* and in Clinical Infections**

Metronidazole has been shown to be active against most isolates of the following bacteria both *in vitro* and in clinical infections as described in the **INDICATIONS AND USAGE** section.

### **Gram-positive anaerobes**

*Clostridium* species

*Eubacterium* species

*Peptococcus* species

*Peptostreptococcus* species

### **Gram-negative anaerobes**

*Bacteroides fragilis* group (*B. fragilis*, *B. distasonis*, *B. ovatus*, *B. thetaiotaomicron*, *B. vulgatus*)

*Fusobacterium* species

The following *in vitro* data are available, **but their clinical significance is unknown.**

Metronidazole exhibits *in vitro* minimal inhibitory concentrations (MIC's) of 8 mcg/mL or less against most ( $\geq 90\%$ ) isolates of the following bacteria; however, the safety and effectiveness of metronidazole in treating clinical infections due to these bacteria have not been established in adequate and well-controlled clinical trials.

### **Gram-negative anaerobes**

*Bacteroides fragilis* group (*B. caccae*, *B. uniformis*)

*Prevotella* species (*P. bivia*, *P. buccae*, *P. disiens*)

## **Susceptibility Tests**

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see:

## INDICATIONS AND USAGE

### Treatment of Anaerobic Bacterial Infections

Metronidazole injection is indicated in the treatment of serious infections caused by susceptible anaerobic bacteria. Indicated surgical procedures should be performed in conjunction with metronidazole injection therapy. In a mixed aerobic and anaerobic infection, antibiotics appropriate for the treatment of the aerobic infection should be used in addition to metronidazole injection.

Metronidazole injection is effective in *Bacteroides fragilis* infections resistant to clindamycin, chloramphenicol and penicillin.

**Intra-Abdominal Infections**, including peritonitis, intra-abdominal abscess and liver abscess, caused by *Bacteroides* species including the *B. fragilis* group (*B. fragilis*, *B. distasonis*, *B. ovatus*, *B. thetaiotaomicron*, *B. vulgatus*), *Clostridium* species, *Eubacterium* species, *Peptococcus* species and *Peptostreptococcus* species.

**Skin and Skin Structure Infections** caused by *Bacteroides* species including the *B. fragilis* group, *Clostridium* species, *Peptococcus* species, *Peptostreptococcus* species and *Fusobacterium* species.

**Gynecologic Infections**, including endometritis, endomyometritis, tubo-ovarian abscess and postsurgical vaginal cuff infection, caused by *Bacteroides* species including the *B. fragilis* group, *Clostridium* species, *Peptococcus* species, *Peptostreptococcus* species and *Fusobacterium* species.

**Bacterial Septicemia** caused by *Bacteroides* species including the *B. fragilis* group and *Clostridium* species.

**Bone and Joint Infections**, as adjunctive therapy, caused by *Bacteroides* species including the *B. fragilis* group.

**Central Nervous System (CNS) Infections**, including meningitis and brain abscess, caused by *Bacteroides* species including the *B. fragilis* group.

**Lower Respiratory Tract Infections**, including pneumonia, empyema and lung abscess, caused by *Bacteroides* species including the *B. fragilis* group.

**Endocarditis** caused by *Bacteroides* species including the *B. fragilis* group.

### Prophylaxis

The prophylactic administration of metronidazole injection preoperatively, intraoperatively and postoperatively may reduce the incidence of postoperative infection in patients undergoing elective colorectal surgery which is classified as contaminated or potentially contaminated. Prophylactic use of metronidazole injection should be discontinued within 12 hours after surgery. If there are signs of infection, specimens for cultures should be obtained for the identification of the causative organism(s) so that appropriate therapy may be given (see **DOSAGE AND ADMINISTRATION**).

To reduce the development of drug-resistant bacteria and maintain the effectiveness of metronidazole injection and other antibacterial drugs, metronidazole injection should be

used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

## **CONTRAINDICATIONS**

### **Hypersensitivity**

Metronidazole injection is contraindicated in patients with a prior history of hypersensitivity to metronidazole or other nitroimidazole derivatives.

### **Psychotic Reaction with Disulfiram**

Use of oral metronidazole is associated with psychotic reactions in alcoholic patients who were using disulfiram concurrently. Do not administer metronidazole to patients who have taken disulfiram within the last two weeks (see **PRECAUTIONS-Drug Interactions**).

### **Interaction with Alcohol**

Use of oral metronidazole is associated with a disulfiram-like reaction to alcohol, including abdominal cramps, nausea, vomiting, headaches and flushing. Discontinue consumption of alcohol or products containing propylene glycol during and for at least three days after therapy with metronidazole (see **PRECAUTIONS-Drug Interactions**).

### **Cockayne Syndrome**

Metronidazole injection is contraindicated in patients with Cockayne syndrome. Severe irreversible hepatotoxicity/acute liver failure with fatal outcomes have been reported after initiation of metronidazole in patients with Cockayne syndrome (see **ADVERSE REACTIONS**).

## **WARNINGS**

### **Central and Peripheral Nervous System Effects**

Severe neurological disturbances, including encephalopathy, cerebellar symptoms, convulsive seizures, peripheral neuropathy, optic neuropathy, and aseptic meningitis, have been reported in patients treated with metronidazole.

Encephalopathy associated with metronidazole may manifest as confusion or decreased level of consciousness, and is associated with widespread lesions on magnetic resonance imaging (MRI) of the brain. Cerebellar toxicity associated with metronidazole may manifest as ataxia, dizziness, dysarthria, nystagmus and saccadic pursuit and is accompanied by T2 flair lesions within the dentate nuclei seen on MRI. Cerebellar toxicity may concurrently occur with encephalopathy, peripheral neuropathy or seizures. CNS symptoms and CNS lesions are generally reversible within days to weeks upon discontinuation of metronidazole injection. Peripheral neuropathy, usually symmetric and mainly of sensory type is characterized by numbness or paresthesia of an extremity. Symptoms may be prolonged after drug discontinuation. Aseptic meningitis may occur within hours of dose administration and generally resolve after metronidazole therapy is

discontinued (see **ADVERSE REACTIONS**).

Advise patients to report neurologic symptoms that occur during metronidazole administration. Discontinue metronidazole treatment if any abnormal neurologic symptoms occur such as ataxia, dizziness, confusion or any other CNS adverse reaction (see **ADVERSE REACTIONS**).

## **PRECAUTIONS**

### **General**

#### **Hepatic Impairment**

Patients with hepatic impairment metabolize metronidazole slowly, with resultant accumulation of metronidazole and increase the plasma concentrations. Reduce the dose of metronidazole injection by 50% in patients with severe hepatic impairment (Child-Pugh C). For patients with mild to moderate hepatic impairment, no dosage adjustment is needed but these patients should be monitored for metronidazole associated adverse events (see **CLINICAL PHARMACOLOGY** and **DOSAGE AND ADMINISTRATION**).

Patients with severe hepatic encephalopathy metabolize metronidazole slowly, with resultant accumulation of metronidazole. This may cause exacerbation of CNS adverse effects. Reduce the dose of metronidazole injection as necessary.

#### **Renal Impairment**

For patients with mild to moderate renal impairment dose adjustment is not considered necessary as elimination half-life is not significantly altered. In patients with severe renal impairment or end stage of renal disease, metronidazole and metronidazole metabolites may accumulate significantly because of reduced urinary excretion in those patients. Monitoring for metronidazole associated adverse events is recommended when metronidazole is administered in patients with severe renal impairment or end stage of renal disease who are not undergoing hemodialysis (see **CLINICAL PHARMACOLOGY**).

Hemodialysis removes significant amounts of metronidazole and its metabolites from systemic circulation. Therefore, supplementation of metronidazole following a hemodialysis session may be necessary.

Patients receiving peritoneal dialysis should be monitored for signs of toxicity due to the potential accumulation of metronidazole metabolites.

#### **Fungal Superinfections**

Known or previously unrecognized candidiasis may present more prominent symptoms during therapy with metronidazole injection and requires treatment with a candidicidal agent.

#### **Use in Patients with Blood Dyscrasias**

Metronidazole is a nitroimidazole and should be used with care in patients with evidence of or history of blood dyscrasia. Agranulocytosis, leukopenia and neutropenia have been associated with metronidazole administration. Monitor complete blood count in these patients.

## **Monitoring for Leukopenia**

Monitoring of complete blood count (CBC) is recommended before, during and after prolonged or repeated courses of metronidazole therapy.

## **Sodium Retention**

Metronidazole injection contains 790 mg of sodium per 100 mL. Care should be taken when administering metronidazole injection to patients receiving a controlled sodium diet or corticosteroids or to patients predisposed to edema.

## **Drug-Resistant Bacteria**

Prescribing metronidazole injection in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

## **Information for Patients**

### **Interaction with Alcohol**

Discontinue consumption of alcoholic beverages or products containing propylene glycol while taking metronidazole injection and for at least three days afterward because abdominal cramps, nausea, vomiting, headaches, and flushing may occur (see **CONTRAINDICATIONS, PRECAUTIONS-Drug Interactions**).

### **Treatment of Bacterial Infections**

Patients should be counseled that antibacterial drugs including metronidazole injection should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When metronidazole injection is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by metronidazole injection or other antibacterial drugs in the future.

## **Drug Interactions**

### **Disulfiram**

Psychotic reactions and confusion have been reported in alcoholic patients who are using metronidazole and disulfiram concurrently. Do not administer metronidazole injection to patients who have taken disulfiram within the last two weeks (see **CONTRAINDICATIONS**).

### **Alcoholic Beverages**

Abdominal cramps, nausea, vomiting, headaches, tachycardia and flushing may occur if alcoholic beverages or products containing propylene glycol are consumed during or following metronidazole therapy. Discontinue consumption of alcohol or products containing propylene glycol before, during and up to 72 hours after therapy with metronidazole injection (see **CONTRAINDICATIONS**).

### **Warfarin and other Oral Anticoagulants**

Metronidazole has been reported to potentiate the anticoagulant effect of warfarin and



other oral coumarin anticoagulants, resulting in a prolongation of prothrombin time and increased risk of hemorrhages. When metronidazole injection is prescribed for patients on this type of anticoagulant therapy, prothrombin time and international normalized ratio (INR) should be carefully monitored and their anticoagulant dose adjusted accordingly. Monitor patients for signs and symptoms of bleeding.

### **Lithium**

In patients stabilized on relatively high doses of lithium, short-term metronidazole therapy has been associated with elevation of serum lithium and, in a few cases, signs of lithium toxicity. Lithium toxicity may lead to renal damage. Frequent monitoring of serum lithium and serum creatinine levels is necessary.

### **Busulfan**

Metronidazole has been reported to increase plasma concentrations of busulfan, which can result in an increased risk for serious busulfan toxicity such as sinusoidal obstruction syndrome, gastrointestinal mucositis and hepatic veno-occlusive disease. Metronidazole injection should not be administered concomitantly with busulfan unless the benefit outweighs the risk. If no therapeutic alternatives to metronidazole are available, and concomitant administration with busulfan is medically needed, frequent monitoring of busulfan plasma concentration should be performed and the busulfan dose should be adjusted accordingly.

### **Drugs that Inhibit CYP450 Enzymes**

The simultaneous administration of drugs that decrease microsomal liver enzyme activity, such as cimetidine, may decrease metabolism and reduce plasma clearance of metronidazole which may result in metronidazole toxicity.

### **Drugs that Induce CYP450 Enzymes**

The simultaneous administration of drugs that induce microsomal liver enzyme activity, such as phenytoin or phenobarbital, may accelerate the elimination of metronidazole and therefore decrease its efficacy.

### **Cytochrome P450 3A4 (CYP3A4) substrates**

Concomitant use of metronidazole injection and CYP3A4 substrates (e.g., amiodarone, tacrolimus, cyclosporine, carbamazepine, phenytoin and quinidine) may increase respective CYP3A4-substrate plasma levels. Monitoring of plasma concentrations of CYP3A4 substrates may be necessary.

### **5-Fluorouracil**

Metronidazole injection decreases the clearance of 5-fluorouracil and may therefore cause 5-fluorouracil toxicity.

### **Vecuronium**

Metronidazole injection may potentiate the effects of vecuronium.

### **Drugs that Prolong the QT interval**

QT prolongation has been reported, particularly when metronidazole was administered with drugs with the potential for prolonging the QT interval.

### **Drug/Laboratory Test Interactions**

Metronidazole may interfere with certain types of determinations of serum chemistry values, such as aspartate aminotransferase (AST, SGOT), alanine aminotransferase (ALT, SGPT), lactate dehydrogenase (LDH), triglycerides and glucose hexokinase. Metronidazole causes an increase in ultraviolet absorbance at 340 nm resulting in falsely decreased values.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Tumors affecting the liver, lung, mammary and lymphatic tissues have been detected in several studies of metronidazole in rats and mice, but not hamsters.

Pulmonary tumors have been observed in all six reported studies in the mouse, including one study in which the animals were dosed on an intermittent schedule (administration during every fourth week only). Malignant tumors were increased in male mice treated at approximately 1,500 mg/m<sup>2</sup> (similar to the maximum recommended daily dose, based on body surface area comparisons). Malignant lymphomas and pulmonary neoplasms were also increased with lifetime feeding of the drug to mice. Mammary and hepatic tumors were increased among female rats administered oral metronidazole compared to concurrent controls. Two lifetime tumorigenicity studies in hamsters have been performed and reported to be negative.

Metronidazole has shown mutagenic activity in *in vitro* assay systems including the Ames test. Studies in mammals *in vivo* have failed to demonstrate a potential for genetic damage.

Metronidazole failed to produce any adverse effects on fertility or testicular function in male rats at doses up to 400 mg/kg/day (approximately 2 times the maximum recommended daily dose based on body surface area comparison) for 28 days. However, rats treated at the same dose for 6 weeks, or longer were infertile and showed severe degeneration of the seminiferous epithelium in the testes as well as marked decreases in testicular spermatid counts and epididymal sperm counts. Fertility was restored in most rats after an eight-week, drug-free recovery period.

Fertility studies have been performed in male mice at doses up to six times the maximum recommended human dose based on mg/m<sup>2</sup> and have revealed no evidence of impaired fertility. However, metronidazole was associated with reversible adverse effects on the male reproductive system (significantly decreased testes and epididymides weight, decreased sperm viability, and increased the incidence of abnormal sperm).

### **Pregnancy**

#### **Teratogenic Effects**

There are no adequate and well-controlled studies of metronidazole injection in pregnant women. There are published data from case-control studies, cohort studies and 2 meta-analyses that include more than 5,000 pregnant women who used metronidazole during pregnancy. Many studies included first trimester exposures. One study showed an increased risk of cleft lip, with or without cleft palate, in infants exposed to metronidazole *in utero*; however, these findings were not confirmed. In addition, more than ten randomized, placebo-controlled clinical trials enrolled more than 5,000 pregnant women to assess the use of antibiotic treatment (including metronidazole) for bacterial vaginosis on the incidence of preterm delivery. Most studies did not show an increased

risk for congenital anomalies or other adverse fetal outcomes following metronidazole exposure during pregnancy. Three studies conducted to assess the risk of infant cancer following metronidazole exposure during pregnancy did not show an increased risk; however, the ability of these studies to detect such a signal was limited.

Metronidazole crosses the placental barrier and its effects on the human fetal organogenesis are not known. Reproduction studies have been performed in rats, rabbits and mice at doses similar to the maximum recommended daily dose based on body surface area comparisons. There was no evidence of harm to the fetus due to metronidazole.

Healthcare provider should carefully consider the potential risks and benefits for each specific patient before prescribing metronidazole injection.

### **Nursing Mothers**

Metronidazole is present in human milk at concentrations similar to maternal serum levels, and infant serum levels can be close to or comparable to infant therapeutic levels. Because of the potential for tumorigenicity shown for metronidazole in mouse and rat studies, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother. Alternatively, a nursing mother may choose to pump and discard human milk for the duration of metronidazole therapy, and for 24 hours after therapy ends and feed her infant stored human milk or formula.

### **Geriatric Use**

In general, dose selection for an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

In geriatric patients, monitoring for metronidazole associated adverse events is recommended (see **CLINICAL PHARMACOLOGY, PRECAUTIONS**). Decreased liver function in geriatric patients can result in increased concentrations of metronidazole that may necessitate adjustment of metronidazole dosage (see **DOSAGE AND ADMINISTRATION**).

### **Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

## **ADVERSE REACTIONS**

The following reactions have been reported during treatment with metronidazole formulations:

*INFECTIONS AND INFESTATIONS:* Vaginal candidiasis

*BLOOD AND LYMPHATIC SYSTEM DISORDERS:* Agranulocytosis, Leukopenia, Neutropenia, Thrombocytopenia, Eosinophilia

*IMMUNE SYSTEM DISORDERS:* Anaphylactic reaction, Hypersensitivity

*METABOLISM AND NUTRITION DISORDERS:* Decreased appetite

*PSYCHIATRIC DISORDERS:* Confusional state, Depression, Insomnia, Decreased libido

*NERVOUS SYSTEM DISORDERS:* Encephalopathy, Seizure, Neuropathy peripheral, Ataxia, Dizziness, Hypoesthesia, Paresthesia, Dysgeusia, Headache, Nystagmus, Aseptic meningitis, Somnolence, Dysarthria, Numbness, Syncope

*EYE DISORDERS:* Optic neuropathy, Saccadic eye movement

*EAR AND LABYRINTH DISORDERS:* Vertigo

*CARDIAC DISORDERS:* QT prolongation has been reported, particularly when metronidazole was administered with drugs with the potential for prolonging the QT interval. Flattening of the T-wave may be seen in electrocardiographic tracings, Tachycardia, Palpitation

*RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS:* Dyspnea

*GASTROINTESTINAL DISORDERS:* Pancreatitis, Abdominal pain, Diarrhea, Nausea, Vomiting, Asthenia, Proctitis

*HEPATOBIILIARY DISORDERS:* Hepatotoxicity/Liver Failure in patients with Cockayne syndrome (see **CONTRAINDICATIONS**), Jaundice

*SKIN AND SUBCUTANEOUS DISORDERS:* Toxic epidermal necrolysis, Swelling face, Pruritus, Urticaria, Hyperhidrosis, Erythema, Rash; Stevens-Johnson syndrome, Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)

*MUSCULOSKELETAL AND CONNECTIVE TISSUE DISORDERS:* Muscle spasms, Arthralgia, Myalgia

*RENAL AND URINARY DISORDERS:* Chromaturia, Dysuria

**Hepatic:** Cases of severe irreversible hepatotoxicity/acute liver failure, including cases with fatal outcomes with very rapid onset after initiation of systemic use of metronidazole, have been reported in patients with Cockayne Syndrome (latency from drug start to signs of liver failure as short as 2 days) (see **CONTRAINDICATIONS**).

*REPRODUCTIVE:* Dyspareunia

*GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS:* Injection site reaction, Malaise, Face edema, Edema peripheral, Chest pain, Chills

*INVESTIGATIONS:* Hepatic enzyme increased.

Patients with Crohn's disease are known to have an increased incidence of gastrointestinal and certain extraintestinal cancers. There have been some reports in the medical literature of breast and colon cancer in Crohn's disease patients who have been treated with metronidazole at high doses for extended periods of time. A cause and effect relationship has not been established. Crohn's disease is not an approved indication for metronidazole injection.

## **OVERDOSAGE**

Signs and symptoms of an overdose may include: nausea, vomiting, and neurotoxic effects, including ataxia, confusion, disorientation, seizures and peripheral neuropathy.

## **Treatment of Overdosage**

Symptomatic treatment is recommended. There is no specific antidote for an overdose of metronidazole. Discontinue metronidazole administration in the event of an overdose. Hemodialysis removes significant amounts of metronidazole and its metabolites from systemic circulation.

## **DOSAGE AND ADMINISTRATION**

Dosage, rate of administration, and duration of treatment are to be individualized and depend upon the indication for use, the patient's age, weight, clinical condition and concomitant treatment, and on the patient's clinical and laboratory response to the treatment.

### **Treatment of Anaerobic Bacterial Infections**

The recommended dosage schedule for adults is:

Loading Dose	15 mg/kg infused intravenously over one hour (approximately 1 gram for a 70-kg adult).
Maintenance Dose	7.5 mg/kg infused intravenously over one hour every six hours (approximately 500 mg for a 70-kg adult). The first maintenance dose should be instituted six hours following the initiation of the loading dose.

Parenteral therapy may be changed to oral metronidazole when conditions warrant, based upon the severity of the disease and the response of the patient to metronidazole injection treatment. The usual adult oral dosage is 7.5 mg/kg every six hours (approximately 500 mg for a 70-kg adult).

A maximum of 4 grams should not be exceeded during a 24-hour period.

The usual duration of therapy is 7 days to 10 days; however, infections of the bone and joint, lower respiratory tract and endocardium may require longer treatment.

### **Dosage Adjustments**

#### **Patients with Severe Hepatic Impairment**

For patients with severe hepatic impairment (Child-Pugh C), the metronidazole dose should be reduced by 50% (see **CLINICAL PHARMACOLOGY** and **PRECAUTIONS**).

#### **Patients Undergoing Hemodialysis**

Hemodialysis removes significant amounts of metronidazole and its metabolites from systemic circulation. The clearance of metronidazole will depend on the type of dialysis membrane used, the duration of the dialysis session, and other factors. If the administration of metronidazole cannot be separated from a hemodialysis session, supplementation of metronidazole dosage following a hemodialysis session should be considered, depending on the patient's clinical situation (see **CLINICAL PHARMACOLOGY**).

#### **Prophylaxis**

For surgical prophylactic use, to prevent postoperative infection in contaminated or

potentially contaminated colorectal surgery, the recommended dosage schedule for adults is:

- a. 15 mg/kg infused over 30 minutes to 60 minutes and completed approximately one hour before surgery; followed by
- b. 7.5 mg/kg infused over 30 minutes to 60 minutes at 6 hours and 12 hours after the initial dose.

It is important that (1) administration of the initial preoperative dose be completed approximately one hour before surgery so that adequate drug levels are present in the serum and tissues at the time of initial incision and (2) metronidazole injection be administered, if necessary, at 6-hour intervals to maintain effective drug levels. Prophylactic use of metronidazole injection should be limited to the day of surgery only, following the above guidelines.

**Caution: Metronidazole injection is to be administered by slow intravenous drip infusion only, either as a continuous or intermittent infusion. Additives should not be introduced into metronidazole injection, unless compatibility is known. If used with a primary intravenous fluid system, the primary solution should be discontinued during metronidazole infusion. DO NOT USE EQUIPMENT CONTAINING ALUMINUM (e.g., NEEDLES, CANNULAE) THAT WOULD COME IN CONTACT WITH THE DRUG SOLUTION AS PRECIPITATES MAY FORM.**

Metronidazole injection is incompatible with (includes but is not limited to): Aztreonam, Cefamandole nafate, Cefoxitin, Penicillin G.

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products (see **DIRECTIONS FOR USE OF PLASTIC CONTAINER**).

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Do not administer unless the solution is clear and the seal is intact.

## **HOW SUPPLIED**

Metronidazole Injection USP, **500 mg/100 mL** (5 mg/mL) is a sterile clear, colorless to pale yellow color, nonpyrogenic, iso-osmotic, buffered solution supplied in 100 mL single-dose non-PVC plastic container. Each 100 mL contains 500 mg metronidazole, USP. It is available as follows:

### **500 mg/100 mL (5 mg/mL):**

1 Single-dose Container in an Overwrap: NDC 80830-2465-1

Carton of 24: NDC 80830-2465-9

Store at 20° to 25°C (68° to 77° F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature] and protect from light during storage.

Do not remove unit from overwrap until ready for use. The overwrap is a moisture barrier. The inner bag maintains the sterility of the product. After removing overwrap, check for minute leaks by squeezing inner bag firmly. If leaks are found, discard solution as sterility may be impaired.

## **DIRECTIONS FOR USE OF PLASTIC CONTAINER**

Metronidazole injection is a ready-to-use iso-osmotic solution. **No dilution or buffering is required.** Do not refrigerate. Each container of metronidazole injection contains 14 mEq of sodium.

Do not connect flexible plastic containers in series in order to avoid air embolism due to possible residual air contained in the primary container.

Pressurizing intravenous solutions contained in flexible plastic containers to increase flow rates can result in air embolism if the residual air in the container is not fully evacuated prior to administration.

Use of a vented intravenous administration set with the vent in the open position could result in air embolism. Vented intravenous administration sets with the vent in the open position should not be used with flexible plastic containers.

Discard any unused portion.

For single-dose only.

### **To open**

Tear overwrap at notch and remove solution container. Visually inspect the container. If the twist-off port is damaged, detached, or not present, discard container as solution path sterility may be impaired. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually. Check for leaks. **Do not add supplementary medication.**

### **Preparation for Administration**

1. Suspend container from eyelet support.
2. Remove twist-off port from bag.
3. Attach administration set. Refer to complete directions accompanying set.

**To report SUSPECTED ADVERSE REACTIONS, contact Amneal Pharmaceuticals at 1-877-835-5472 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

Manufactured by:

**Amneal Pharmaceuticals Pvt. Ltd.**

### **Parenteral Unit**

Ahmedabad 382110, INDIA

Distributed by:

**Amneal Pharmaceuticals LLC**

Bridgewater, NJ 08807

Rev. 10-2023-01

## **PRINCIPAL DISPLAY PANEL**

**NDC 80830-2465-1**

**Metronidazole Injection USP, 500 mg/100 mL (5 mg/mL)**

**Rx only**

**Intravenous Bag Label**

**Amneal Pharmaceuticals LLC**

NDC 80830-2465-1

Rx only

# Metronidazole Injection, USP

## 500 mg/100 mL (5 mg/mL)

**For Intravenous Infusion Only**  
**100 mL Sterile Single-Dose Container**  
**Discard Unused Portion**

Each 100 mL contains 500 mg metronidazole, USP; 790 mg sodium chloride, USP; 47.6 mg dibasic sodium phosphate dried, USP; and 22.9 mg citric acid anhydrous, USP.  
Each container contains 14 mEq of sodium.  
pH 5.5 (4.5 to 7.0). Osmolarity 308.66 mOsmol/L (Calc).

**Usual Dosage:** See insert.

**DO NOT ADD SUPPLEMENTARY MEDICATION.**

Store in moisture barrier overwrap at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature] and protect from light during storage.

Mfg. Lic. No.: G/28D/LVP/20

Distributed by: **Amneal Pharmaceuticals LLC**

Bridgewater, NJ 08807

MADE IN INDIA.

Rev. 06-2023-00



Lot:

Exp:

Space for overcoding  
at the time of packing  
(For Lot and Mfg. / Exp. Date)  
28 x 28 mm

**NDC 80830-2465-1**

**Metronidazole Injection USP, 500 mg/100 mL (5 mg/mL)**

**Rx only**

**Pouch Label**

**Amneal Pharmaceuticals LLC**



TO OPEN - TEAR AT NOTCH

Do not remove unit from overwrap until ready for use.

The overwrap is a moisture barrier. The inner bag maintains the sterility of the product.

NDC 80830-2465-1

Rx only

# Metronidazole Injection, USP

**500 mg/100 mL (5 mg/mL)**

**For Intravenous Infusion Only**  
**100 mL Sterile Single-Dose Container**  
**Discard Unused Portion**

Each 100 mL contains 500 mg metronidazole, USP; 790 mg sodium chloride, USP; 47.6 mg dibasic sodium phosphate dried, USP; and 22.9 mg citric acid anhydrous, USP. Each container contains 14 mEq of sodium. pH 5.5 (4.5 to 7.0).

Osmolarity 308.66 mOsmol/L (Calc).

**Usual Dosage:** See insert.

**DO NOT ADD SUPPLEMENTARY MEDICATION.**

Store in moisture barrier overwrap at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature] and protect from light during storage.

Manufactured by: **Amneal Pharmaceuticals Pvt. Ltd.**  
**Parenteral Unit**  
Ahmedabad 382110, INDIA

Distributed by: **Amneal Pharmaceuticals LLC**  
Bridgewater, NJ 08807

Mfg. Lic. No. G/28D/LVP/20

Rev. 10-2023-01

PMI-1527



Blank Area for  
Variable data of GTIN, LOT, EXP and  
UNIQUE SERIAL NO. on each pouch  
with 2D DATA MATRIX shall be printed.  
110 x 40 mm

**NDC 80830-2465-9**

**Metronidazole Injection USP, 500 mg/100 mL (5 mg/mL)**

**Rx only**

**Carton Label**

**Amneal Pharmaceuticals LLC**

NDC 80830-2465-9

**24 x 100 mL Sterile  
Single-Dose Plastic Container**

# Metronidazole Injection, USP

**500 mg/100 mL (5 mg/mL)**

- **For Intravenous Infusion Only**
- **Discard Unused Portion**

Each 100 mL contains 500 mg metronidazole, USP; 790 mg sodium chloride, USP; 47.6 mg dibasic sodium phosphate dried, USP; and 22.9 mg citric acid anhydrous, USP. Each container contains 14 mEq of sodium. pH 5.5 (4.5 to 7.0). Osmolarity 308.66 mOsmol/L (Calc).

**Usual Dosage:** See insert.

**DO NOT ADD SUPPLEMENTARY MEDICATION.**

Store in moisture barrier overwrap at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature] and protect from light during storage.

**Manufactured by:** Amneal Pharmaceuticals Pvt. Ltd.  
Parenteral Unit  
Ahmedabad 382110, INDIA

**Distributed by:** Amneal Pharmaceuticals LLC  
Bridgewater, NJ 08807

Mfg. Lic. No. G/28D/LVP/20

Rev. 10-2023-01

For Lot & Exp. please refer to sticker  
attached to carton.

**Rx only**



**METRONIDAZOLE**

metronidazole injection, solution

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:80830-2465
<b>Route of Administration</b>	INTRAVENOUS		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>METRONIDAZOLE</b> (UNII: 140QMO216E) (METRONIDAZOLE - UNII:140QMO216E)	METRONIDAZOLE	500 mg in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	790 mg in 100 mL
<b>SODIUM PHOSPHATE, DIBASIC, ANHYDROUS</b> (UNII: 22ADO53M6F)	47.6 mg in 100 mL
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	22.9 mg in 100 mL
<b>WATER</b> (UNII: 059QF0KO0R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:80830-2465-9	24 in 1 CARTON	06/17/2023	
1	NDC:80830-2465-1	1 in 1 POUCH		
1		100 mL in 1 BAG; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA217665	06/17/2023	

**Labeler** - Amneal Pharmaceuticals Private Limited (675474666)

**Registrant** - Amneal Pharmaceuticals of New York, LLC (123797875)

## Establishment

Name	Address	ID/FEI	Business Operations
Amneal Pharmaceuticals Private Limited		675474666	analysis(80830-2465) , label(80830-2465) , manufacture(80830-2465) , pack(80830-2465) , sterilize(80830-2465)

