BENGAY ULTRA STRENGTH REGULAR SIZE- menthol patch Johnson & Johnson Consumer Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Ultra Strength BENGAY® Pain Relieving Patch

Drug Facts

Active ingredient

Menthol 5%

Purpose

Topical analgesic

Uses

- temporarily relieves minor aches and pains of muscles and joints associated with:
 - simple backache
 - arthritis
 - strains
 - bruises
 - sprains

Warnings

For external use only.

Do not use

- on wounds or damaged skin
- with a heating pad
- on a child under 12 years of age with arthritis-like conditions

Ask a doctor before use if you have

redness over the affected area

When using this product

- avoid contact with eyes or mucous membranes
- do not bandage tightly

Stop use and ask a doctor if

- condition worsens or symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days
- excessive skin irritation occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately.

Directions

- open pouch and remove patch
- if desired, cut patch to size
- peel off protective backing and apply sticky side to affected area
- adults and children 12 years of age and older: apply to affected area not more than 3 to 4 times daily
- children under 12 years of age: consult a doctor

Other information

store at 20° to 25°C (68° to 77°F)

Inactive ingredients

calcined kaolin, cellulose gum, glycerin, methyl acrylate/2-ethylhexyl acrylate copolymer, methylparaben, polyacrylic acid, polysorbate 80, propylparaben, silica, sodium polyacrylate, sodium polyacrylate starch, sorbitan oleate, sorbitol, tartaric acid, titanium dioxide, water

Questions?

call toll-free 800-223-0182 or 215-273-8755 (collect)

Dist: Johnson & Johnson Consumer Products Company Division of Johnson & Johnson Consumer Companies, Inc. Skillman, NJ 08558 USA

PRINCIPAL DISPLAY PANEL - 5 Pouch Carton

ULTRA STRENGTH BENGAY® PAIN RELIEVING PATCH Menthol Topical Analgesic Patch • POWERFUL PAIN RELIEF

- DEEP, PENETRATING HEAT
- LONG LASTING, UP TO 8 HOURS

REGULAR SIZE

Good for Neck to Arm

5 INDIVIDUALLY SEALED PATCHES 3.9 in \times 5.5 in (10 cm \times 14 cm)



<form><form></form></form>	<section-header><form></form></section-header>	\\\c	
Active ingredient Phrpose Aenthol 5% Topical analysesic Vises members of muscles and joints associated with: asimple backache arthitis asimple backache arthitis wormings or catternal use only. Do not use on a child under 12 years of age with arthritis-like conditions lake a doctor before use if you have and control form or damaged skin and a child under 12 years of age with arthritis-like conditions add on to bandage tightly Map use and ask a doctor if acod contact with eyes or mucous membranes a do not bandage tightly Map use and ack a doctor if acod contact with eyes or mucous membranes a do not bandage tightly Map use and ack a doctor if acod contact with eyes or mucous membranes a do not bandage tightly Map use and ack a doctor if acod contact and cour agin within a few days accessive shin irritation occurs Geep out of reach of children. If swallowed, get medical help or contact a Poison Control Center inmediately. Directions Diopen pouch and remove patch if desred, cut patch to size apad of protective backing and apply sticky side to affected area addits and children. If swallowed tod ato area at 20° to 25°C (68° to 77°F) anactive longroffiendis	Lettre ingredient Phrpose Menthol 5% Topical analyseic Asso: a temporarily relieves minor aches and pairs of muscles and joints associated with: a single backache a trinitis strains bruises a printip for caternal use only. for caternal use only. for caternal use only. for caternal use only. to no wounds or damaged skin with a heating pad mon a child under 12 years of age with arthritis-like conditions for caternal use only. for caternal use only. to no wounds or damaged skin with a heating pad mon a child under 12 years of age with arthritis-like conditions for caternal use only. for caternal use only. to no wounds or damaged skin with a heating pad mon ot child use only. for caternal use only. for caternal use only. to not bandage tightly heat actorn before use if you have mon occurs for contact a Poison Control Center immediately. Directions noopen pocch and remove patch if desred, cut patch to size nopen of protective backing and apply sticky side to affected area for pactor for gree consult adord for pactor for consult and topic mactive log redients cating and apply sticky side to affected area for pactor for gree consult adord toreadord for pactore for pact	Benngan B	BENGAY
ISes: isimpos backache withits straine mouses seprains Warnings: or extend use only. on outse outse<	ISes: is importanily relieves minor aches and paire of muscles and joints associated with: is simple backache withits is simple backache withits is miple backache withits is major aches withits is major aches with a heating pad is on a child under 12 years of age with arthritis-like conditions is a doctor before use if you have is a doctor before use if you have is on ont tase is a doctor before use if you have is on ont tase is a doctor before use if you have is constitute is a doctor before use if you have is constitute is a doctor before use if you have is constitute is a doctor before use if you have is constitute is a doctor before use if you have is constitute is avoid constact a doctor if is constitute and tak a doctor if is constitute with any so mucous again withit is five days is constitute and tak a doctor if is constitute with and remove patche if desired, cut patch to size is constitut a doctor if desired, cut patch to size is age and older: apply to affected area is an ant more than 3 to 4 times daily is difter undref 12 years of age and older: apple of prot	1777 AND 1977	-
simple backache arthitis strains bruises esprains Warnings for external use only. Do not use on wounds or damaged skin with a heating pad on a child under 12 years of age with arthritis-like conditions like a doctor before use if you have arechess over the affected area When using this product anxid contact with eyes or mucous membranes and on to bandage tightly Non use and ask a doctor if condition worsens or symptoms persist for more than 7 days areginers clear up and occur again within a few days an excessive skin irritation occurs Geop out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately. Directions open pouch and remove patch if desired, cut patch to size peel off protective backing and apply sticky side to affected area peel off protective backing and apply sticky side to affected area peel off protective backing and apply sticky side to affected area peel off protective backing and apply sticky side to affected area peel off protective backing and apply sticky side to affected area peel off protective backing and apply sticky side to affected area peel off protective backing and apply sticky side to affected area peel off protective backing calculated adoctor Dither information store at 20° to 25°C (68° to 77°F) mactive ingredients cakined doubles gum, glycenin, methyl acrylate/2-ethylhexyl acrylate copolymer, nethylegraben, polyacrylic acid, polysorbate 80, propybaraben, silica, sodium polyacrylate starch, portain cleate, softault, tartaric acid, thanium dinoxde, water Cheeslions? Call toll-free 800-223-0182 or 215-273-8755 (collect)	a simple backache arthitis strains bruises exprains Varinings or external use only. Io no tuse Io no wounds or damaged skin with a heating pad Io na a child under 12 years of age with arthritis-like conditions In a child under 12 years of age with arthritis-like conditions In a child under 12 years of age with arthritis-like conditions In a child under 12 years of age with arthritis-like conditions In a child under 12 years of age with arthritis-like conditions In a child under 12 years of age with arthritis-like conditions In a child under 12 years of age with arthritis-like conditions In a child under 12 years of age with arthritis-like conditions In a child under 12 years of age with arthritis-like conditions In a child and remove patch If desired, cut patch to size Inped off protective backing and apply sticky side to affected area Inped off protective backing and apply sticky side to affected area Inped off protective backing and apply sticky side to affected area Inped off protective backing and apply sticky side to affected area Inped off protective backing and apply sticky side to affected area Inped off protective backing and apply sticky side to affected area Inped off protective backing and apply sticky side to affected area Inped off protective backing and apply sticky side to affected area Inped off protective backing and apply sticky side to affected area Internation Inter information Inter inform		
For external use only. Do not use ion wounds or damaged skin with a heating pad ion a child under 12 years of age with arthritis-like conditions Lak a doctor before use if you have archess over the affected area When using this product archidess over the affected area When using this product archess over the affected area When using this product archess over the affected area when using this product archest up and occur again within a few days a condition worsens or symptoms parsist for more than 7 days symptoms clear up and occur again within a few days = excessive skin irritation occurs Gree out of reach of children. If swallowed, get medical help or contact a Poison Control Center inmediately. Directions open pouch and remove patch airdesrid, cut patch to size pale off protective backing and apply sticky side to affected area adults and children 12 years of age consult a doctor Differ information = store at 20° to 25°C (68° to 77°F) machive ingredients calcined kackin, cellulose gum, glycenin, methyl acrylate?-ethylhexyl acrylate copolymer, nethylparaben, polyacrylate at 20° to 25°C (68° to 77°F) machive ingredients ca	For external use only. to not use non wounds or damaged skin m with a heating pad non a child under 12 years of age with arthritis-like conditions lak a doctor before use if you have rechess over the affectual area When using this product rechess over the affectual area When using this product rechess over the affectual area whon uses or symptoms parsist for more than 7 days rechest of children. If swallowed, get medical help or contact a Poison Control Center immediately. Directions rope of protective bacing and apply sticky side to affectad area adults and children 12 years of age and older: apply to affectad area adults and children 12 years of age: consult a doctor Differe information = store at 20° to 25°C (68° to 77°F) machtive longredients calcined kacin, calclulose gum, glycenin, methyl acrylate/2-ethylhexyl acrylate copolymer, nethylearaben, polyscrylic acid, polysorabe 80, propybaraben, silica, andium polyacrylate, sodium polyacrylate starch, othildren ic add, thainum dioxide, water Ruestions? call toll-free 800-223-0182 or 215-273-8755 (collect)	simple beckache 🗰 arthritis 🗰 strains 🗰 bruises 🗰 sprains	
non wounds or damaged skin with a heating pad on a child under 12 years of age with arthritis-like conditions lak a doctor before use if you have mechaes over the affected area When using this product a void contact with eyes or mucous membranes do not bandage tightly Nop use and ack a doctor if econdition worsane or symptoms persist for more than 7 days esymptoms clear up and occur again within a few days excessive skin irritation occurs Geop out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately. Diffections a point of more patch if desired, cut patch to size pael of protective backing and apply sticky side to affected area adults and children 12 years of age and okser: apply to affected area adults and children 12 years of age and okser: apply to affected area children under 12 years of age consult a doctor Dither information extra 20° to 25°C (68° to 77°F) mactive ingredients calculad kaolin, cellulose gum, glycesin, methyl acrylate 2-ethylhexyl acrylate scapolymer, nethylaraten, polyacrytic add, polysoritate 80, propykeraben, silica, sodium polyacrylate, sodium polyacrylate starch, orithan okate, softridi, tratratic add, thanium dioxdd, water Ruestions? call toll-free 800-223-0182 or 215-273-8755 (collect)	non wounds or damaged skin with a heating pad on a child under 12 years of age with arthritis-like conditions lak a doctor before use if you have mechaes over the affected area When using this product a void contact with eyes or mucous membranes do not bandage tightly Nop use and ack a doctor if econdition worsane or symptoms persist for more than 7 days esymptoms clear up and occur again within a few days excessive skin irritation occurs Geop out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately. Diffections a point of more patch if desired, cut patch to size pael of protective backing and apply sticky side to affected area adults and children 12 years of age and okser: apply to affected area adults and children 12 years of age and okser: apply to affected area children under 12 years of age consult a doctor Dither information extra 20° to 25°C (68° to 77°F) mactive ingredients calculad kaolin, cellulose gum, glycesin, methyl acrylate 2-ethylhexyl acrylate scapolymer, nethylaraten, polyacrytic add, polysoritate 80, propykeraben, silica, sodium polyacrylate, sodium polyacrylate starch, orithan okate, softridi, tratratic add, thanium dioxdd, water Ruestions? call toll-free 800-223-0182 or 215-273-8755 (collect)		
arechess over the affected area When using this product a root contact with eyes or mucous membranes a do not bandage tightly When using this product a condition worsens or symptoms persist for more than 7 days a condition worsens or symptoms persist for more than 7 days excessive skin irritation occurs a symptoms clear up and occur again within a few days excessive skin irritation occurs Geep out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately. Immediately Directions appel off protective backing and apply sticky side to affected area audus and children 12 years of age: consult a doctor Difter information estore at 20° to 25°C (68° to 77°F) Immedive ingredients calcine ad children. Active ingredients calcine ad children, cellulose gum, glycenin, methyl acrylate/2-ethylhexyl acrylate copolymer, nothlan oleate, sorthiol, tartaric acid, thanium dioxide, water Softwart at cole to 215-273-8755 (collect) Softwart at cole to 215-273-8755 (collect)	arechess over the affected area When using this product a root contact with eyes or mucous membranes a do not bandage tightly When using this product a condition worsens or symptoms persist for more than 7 days a condition worsens or symptoms persist for more than 7 days excessive skin irritation occurs a symptoms clear up and occur again within a few days excessive skin irritation occurs Geep out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately. Immediately Directions appel off protective backing and apply sticky side to affected area audus and children 12 years of age: consult a doctor Difter information estore at 20° to 25°C (68° to 77°F) Immedive ingredients calcine ad children. Active ingredients calcine ad children, cellulose gum, glycenin, methyl acrylate/2-ethylhexyl acrylate copolymer, nothlan oleate, sorthiol, tartaric acid, thanium dioxide, water Softwart at cole to 215-273-8755 (collect) Softwart at cole to 215-273-8755 (collect)	on wounds or damaged skin 🛛 🗰 with a heating pad	
a avoid contact with eyes or mucous membranes do not bandage tightly And use and ask a doctor if condition worsens or symptoms persist for more than 7 days expensions clear up and occur again within a few days does excessive skin irritation occurs deep out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately. Directions open pouch and remove patch if desred, cut patch to size pped off protective backing and apply sticky side to affected area outus and children under 12 years of age and okler: apply to affected area outus and children if gene at 20° to 25°C (68° to 77°F) mactive ingredients cakined kacin, celkukes gum, glycerin, methyl acrylate/2-ethylhexyl acrylate copolymer, orbitan oleate, sorbitol, tartaric acid, titanium dioxide, water Ruestions? call toll-free 800-223-0182 or 215-273-8755 (collect)	a avoid contact with eyes or mucous membranes do not bandage tightly And use and ask a doctor if condition worsens or symptoms persist for more than 7 days expensions clear up and occur again within a few days does excessive skin irritation occurs deep out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately. Directions open pouch and remove patch if desred, cut patch to size pped off protective backing and apply sticky side to affected area outus and children under 12 years of age and okler: apply to affected area outus and children if gene at 20° to 25°C (68° to 77°F) mactive ingredients cakined kacin, celkukes gum, glycerin, methyl acrylate/2-ethylhexyl acrylate copolymer, orbitan oleate, sorbitol, tartaric acid, titanium dioxide, water Ruestions? call toll-free 800-223-0182 or 215-273-8755 (collect)	A a doctor before use if you have redness over the affected area	
a condition worsens or symptoms persist for more than 7 days a symptoms clear up and occur again within a few days excessive skin irritation occurs Greep out of reach of children . If swallowed, get medical help or contact a Poison Control Center immediately. Directions open pouch and remove patch if desired, cut patch to size peel off protective backing and apply sticky side to affected area adults and children 12 years of age consult a doctor Diher information extors at 20° to 25°C (68° to 77°F) mactive ingredients cakined kaclin, cellulose gum, glycerin, methyl acrylate/2-ethylhexyl acrylate copolymer, nethylparaben, polyacrylic acid, polysorbate 80, propylparaben, silica, sodium polyacrylate, sodium polyacrylate starch, orbitario clear, sodii (t, tartaric acid, tutanium dioxide, water Ruestions? call toll-free 880-223-0182 or 215-273-8755 (collect)	a condition worsens or symptoms persist for more than 7 days a symptoms clear up and occur again within a few days excessive skin irritation occurs Greep out of reach of children . If swallowed, get medical help or contact a Poison Control Center immediately. Directions open pouch and remove patch if desired, cut patch to size peel off protective backing and apply sticky side to affected area adults and children 12 years of age consult a doctor Diher information extors at 20° to 25°C (68° to 77°F) mactive ingredients cakined kaclin, cellulose gum, glycerin, methyl acrylate/2-ethylhexyl acrylate copolymer, nethylparaben, polyacrylic acid, polysorbate 80, propylparaben, silica, sodium polyacrylate, sodium polyacrylate starch, orbitario clear, sodii (t, tartaric acid, tutanium dioxide, water Ruestions? call toll-free 880-223-0182 or 215-273-8755 (collect)	hen using this product avoid contact with eyes or mucous membranes 🛛 🗰 do not bandage tightly	<u>a</u> 2
peel off protective backing and apply sticky side to affected area peel off protective backing and apply sticky side to affected area adults and children 12 years of age and older: apply to affected area not more than 3 to 4 times daily children under 12 years of age: consult a doctor Other information = store at 20° to 25°C (68° to 77°F) inactive ingredients cakined kaolin, cellulose gum, glycerin, methyl acrylate/2-ethylhexyl acrylate copolymer, nethylparaben, polyacrylic acid, polyacrate 80, propybaraben, silica, sodium polyacrylate, sodium polyacrylate starch, portiation oleate, sorbitof, tartaric acid, titanium dioxide, water Questions? call toll-free 800-223-0182 or 215-273-8755 (collect)	a open pouch and remove patch if desred, cut patch to size peel off protective backing and apply sticky side to affected area adults and children 12 years of age and older: apply to affected area not more than 3 to 4 times daily children under 12 years of age: consult a doctor Other information = store at 20° to 25°C (68° to 77°F) mactive ingredients calcined kaolin, cellulose gum, glycerin, methyl acrylate/2-ethylhexyl acrylate copolymer, nothylopataben, polyacrylic acid, polyonate 80, propybaraben, silica, sodium polyacrylate, sodium polyacrylate starch, orbitan oleate, sorbitol, tartaric acid, titanium dioxide, water Questions? call toll-free 880-223-0182 or 215-273-8755 (collect)	condition worsens or symptoms persist for more than 7 days symptoms clear up and occurs	NG
Dther information m store at 20° to 25°C (68° to 77°F) nactive ingredients cakined kaolin, callulose gum, glycerin, methyl acrylate/2-ethylhexyl acrylate copolymer, nethylopraben, polyacrylate starch, polyacrylate starch, polyacrylate, softwork (in the start cald, titanium dioxide, water Questions? call toll-free 800-223-0182 or 215-273-8755 (collect)	Other information = store at 20° to 25°C (68° to 77°F) nactive ingredients cakined kaolin, callulose gum, glycerin, methyl acrylate/2-ethylhexyl acrylate copolymer, nethylparaben, polyacrylate starch, orbitan oleate, softivol, tartaric acid, titanium dioxide, water Questions? call toll-free 800-223-0182 or 215-273-8755 (collect)	open pouch and remove patch if desired, cut patch to size peel off protective backing and apply sticky side to affected area adults and children 12 years of age and older: apply to affected area not more than 3 to 4 times daily	m in the second second
nactive ingredients cakined kaolin, cellulose gum, glycerin, methyl acrylate/2-ethylhexyl acrylate copolymer, nethylperaben, polyacrylic acid, polyaorbate 90, propylaraben, silica, sodium polyacrylate, sodium polyacrylate starch, norbitan okate, sorbitol, tarlaric acid, titanium dioxide, water Questions? call toll-free 800-223-0182 or 215-273-8755 (collect)	nactive ingredients cakined kaolin, callulose gum, glycerin, methyl acrylata/2-ethylhexyl acrylate copolymer, nethylparaben, polyacrylic acid, polysorbate 80, propylparaben, silica, sodium polyacrylate, sodium polyacrylate starch, orbitan oleate, sorbitol, tarlaric acid, titanium dioxide, water Questions? call toll-free 800-223-0182 or 215-273-8755 (collect)		2
Questions? call toll-free 800-223-0182 or 215-273-8755 (collect)	Questions? call toll-free 800-223-0182 or 215-273-8755 (collect)	active ingredients calcined kaolin, cellulose gum, glycenin, methyl acrylata'2-ethylhexyl acrylate copolymer, ethylparaben, polyacrylic acid, polysorbate 90, propylparaben, silica, sodium polyacrylate, sodium polyacrylate starch, richan oleate, sorbitol, tartaric acid, titanium dioxide, water	
			S
///	///	D 0 3 8 9 1 Citiz Johnson & Johnson Consumer Products Company DMskin of Johnson & Johnson Consumer Companies, inc. Skillman, NJ (6556 USA CJUSJ CC) 2010 www.bengary.com Made in Taiwan	

\$1.00 COUPON INSIDE



BENGAY ULTRA STRENGTH REGULAR SIZE

menthol patch

Product Information			
Product T ype	HUMAN OTC DRUG	Item Code (Source)	NDC:58232-6000
Route of Administration	TOPICAL		

Active Ingredient/	Active	e Moiety					
		Ingredient Name		Basis of	Strength	Strength	
Menthol, Unspecified F UNII:L7T10EIP3A)	Form (U	NII: L7T10EIP3A) (Menthol, Unspecified Form	-	Menthol, Un Form	specified	50 mg	
Inactive Ingredien	its						
		Ingredient Name				Strength	
Carboxymethylcellulos	se Sodiı	um, Unspecified Form (UNII: K679OBS311)					
Glycerin (UNII: PDC6A3	BCOOX)						
Methylparaben (UNII: A	A2I8C7H	I9 T)					
Polysorbate 80 (UNII: 6	OZP39Z	ZG8 H)					
Propylparaben (UNII: Z	8IX2SC	10H)					
Silicon Dioxide (UNII: E	ETJ7Z6X	(BU4)					
Sorbitan Monooleate (UNII: 06	XEA2VD56)					
Sorbitol (UNII: 506T60)	A25R)						
Tartaric Acid (UNII: W4	8881119)H)					
Titanium Dioxide (UNI	I: 15FIX9	OV2JP)					
Water (UNII: 059QF0KC	00R)						
Packaging							
# Item Code		Package Description	Ma	rketing Start Date	Marketin	Marketing End Date	
1 NDC:58232-6000-5	5 in 1 C	ARTON	09/0) 1/20 12			
1	1 in 1 P	OUCH; Type 0: Not a Combination Product					
Marketing Info	rmat	ion					
Marketing Catego	ory	Application Number or Monograph Cita	ation	Marketing Start Dat	e Marketi	ng End Date	
OTC MONOGRAPH NOT	FINAL	part348		09/01/2012			

Labeler - Johnson & Johnson Consumer Inc. (002347102)

Revised: 12/2019

Johnson & Johnson Consumer Inc.