

DEXAMETHASONE- dexamethasone injection, solution MWI (Vet One)

DEXAMETHASONE INJECTION 2 mg/mL Dexamethasone Sterile Injection Solution for intravenous or intramuscular injection

Veterinary

Not For Use in Humans

CAUTION Federal Law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION

DEXAMETHASONE INJECTION 2 mg/mL is a synthetic analogue of prednisolone, having similar but more potent anti-inflammatory therapeutic action and diversified hormonal and metabolic effects. Modification of the basic corticoid structure as achieved in DEXAMETHASONE INJECTION 2 mg/mL offers enhanced anti-inflammatory effect compared to older corticosteroids. The dosage of DEXAMETHASONE INJECTION 2 mg/mL required is markedly lower than that of prednisone and prednisolone.

DEXAMETHASONE INJECTION 2 mg/mL is not species-specific; however, the veterinarian should read the sections on **INDICATIONS, DOSAGE, SIDE EFFECTS, CONTRAINDICATIONS, PRECAUTIONS,** and **WARNINGS** before this drug is used.

DEXAMETHASONE INJECTION 2 mg/mL is intended for *intravenous or intramuscular* administration. Each mL contains 2 mg dexamethasone, 500 mg polyethylene glycol 400, 9 mg benzyl alcohol, 1.8 mg methylparaben and 0.2 mg propylparaben as preservatives, 4.75% alcohol, HCl to adjust pH to approximately 4.9, water for injection q.s.

EXPERIMENTAL STUDIES

Experimental animal studies on dexamethasone have revealed it possesses greater anti-inflammatory activity than many steroids. Veterinary clinical evidence indicates dexamethasone has approximately twenty times the anti-inflammatory activity of prednisolone and seventy to eighty times that of hydrocortisone. Thymus involution studies show dexamethasone possesses twenty-five times the activity of prednisolone. In reference to mineralocorticoid activity, dexamethasone does not cause significant sodium or water retention. Metabolic balance studies show that animals on controlled and limited protein intake will exhibit nitrogen losses on exceedingly high dosages.

INDICATIONS

DEXAMETHASONE INJECTION 2 mg/mL is indicated for the treatment of primary bovine ketosis and as an anti-inflammatory agent in the bovine and equine.

As supportive therapy, DEXAMETHASONE INJECTION 2 mg/mL may be used in the management of various rheumatic, allergic, dermatologic, and other diseases known to be responsive to anti-inflammatory corticosteroids. DEXAMETHASONE INJECTION 2 mg/mL may be used intravenously as supportive therapy when an immediate hormonal

response is required.

Bovine Ketosis DEXAMETHASONE INJECTION 2 mg/mL is offered for the treatment of primary ketosis. The gluconeogenic effects of DEXAMETHASONE INJECTION 2 mg/mL, when administered intramuscularly, are generally noted within the first 6 to 12 hours. When DEXAMETHASONE INJECTION 2 mg/mL is used intravenously, the effects may be noted sooner. Blood sugar levels rise to normal levels rapidly and generally rise to above normal levels within 12 to 24 hours. Acetone bodies are reduced to normal concentrations usually within 24 hours. The physical attitude of animals treated with DEXAMETHASONE INJECTION 2 mg/mL brightens and appetite improves, usually within 12 hours. Milk production, which is suppressed as a compensatory reaction in this condition, begins to increase. In some instances, it may even surpass previous peaks. The recovery process usually takes from 3 to 7 days.

Supportive Therapy DEXAMETHASONE INJECTION 2 mg/mL may be used as supportive therapy in mastitis, metritis, traumatic gastritis, and pyelonephritis, while appropriate primary therapy is administered. In these cases, the corticosteroid combats accompanying stress and enhances the feeling of general well-being. DEXAMETHASONE INJECTION 2 mg/mL may also be used as supportive therapy in inflammatory conditions, such as arthritic conditions, snake bite, acute mastitis, shipping fever, pneumonia, laminitis, and retained placenta.

Equine DEXAMETHASONE INJECTION 2 mg/mL is indicated for the treatment of acute musculoskeletal inflammations, such as bursitis, carpalitis, osselets, tendonitis, myositis, and sprains. If bony changes exist in any of the conditions, joints, or accessory structures, responses to DEXAMETHASONE INJECTION 2 mg/mL cannot be expected. In addition, DEXAMETHASONE INJECTION 2 mg/mL may be used as supportive therapy in fatigue, heat exhaustion, influenza, laminitis, and retained placenta provided that the primary cause is determined and corrected.

ADMINISTRATION AND DOSAGE

Therapy with DEXAMETHASONE INJECTION 2 mg/mL, as with any other potent corticosteroid, should be individualized according to the severity of the condition being treated, anticipated duration of steroid therapy, and the animal's threshold or tolerance for steroid excess.

Treatment may be changed over to DEXAMETHASONE INJECTION 2 mg/mL from any other glucocorticoid with proper reduction or adjustment of dosage.

Bovine - DEXAMETHASONE INJECTION 2 mg/mL - 5 to 20 mg intravenously or intramuscularly.

Dexamethasone Powder may be administered or the parenteral dose repeated as needed.

Equine - DEXAMETHASONE INJECTION 2 mg/mL - 2.5 to 5 mg intravenously or intramuscularly.

Dexamethasone Powder may be administered or the parenteral dose repeated as needed.

CONTRAINDICATIONS

Except for emergency therapy, do not use in animals with chronic nephritis and hypercorticalism (Cushing's syndrome). Existence of congestive heart failure, diabetes, and osteoporosis are relative contraindications. Do not use in viral infections during the viremic stage.

PRECAUTIONS

Animals receiving DEXAMETHASONE INJECTION 2 mg/mL should be under close observation. Because of the anti-inflammatory action of corticosteroids, signs of infection may be masked and it may be necessary to stop treatment until a further diagnosis is made. Overdosage of some glucocorticoids may result in sodium retention, fluid retention, potassium loss, and weight gain.

DEXAMETHASONE INJECTION 2 mg/mL may be administered to animals with acute or chronic bacterial infections providing the infections are controlled with appropriate antibiotic or chemotherapeutic agents.

Doses greater than those recommended in horses may produce a transient drowsiness or lethargy in some horses. The lethargy usually abates in 24 hours.

Use of corticosteroids, depending on dose, duration, and specific steroid, may result in inhibition of endogenous steroid production following drug withdrawal. In patients presently receiving or recently withdrawn from systemic corticosteroid treatments, therapy with a rapid acting corticosteroid should be considered in unusually stressful situations.

WARNINGS

Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate parturition followed by dystocia, fetal death, retained placenta, and metritis.

Additionally, corticosteroids administered to dogs, rabbits, and rodents during pregnancy have produced cleft palate. Other congenital anomalies including deformed forelegs phocomelia, and anasarca have been reported in offspring of dogs which received corticosteroids during pregnancy.

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

SIDE EFFECTS

Side effects, such as SAP and SGPT enzyme elevations, weight loss, anorexia, polydipsia, and polyuria have occurred following the use of synthetic corticosteroids in dogs. Vomiting and diarrhea (occasionally bloody) have been observed in dogs and cats.

Cushing's syndrome in dogs has been reported in association with prolonged or repeated steroid therapy.

Corticosteroids reportedly cause laminitis in horses.

CONTACT INFORMATION

To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data sheet (SDS), contact Sparhawk Laboratories Inc at 1-800-255-6388 or 1-913-888-7500. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

HOW SUPPLIED

DEXAMETHASONE INJECTION 2 mg/mL

100 mL -NDC 13985-533-03

250 mL - NDC 13985-533-25

Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F).

Protect from freezing.

Approved by FDA under ANADA # 200-324

WARNING

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Usual Dose

Bovine-5 to 20 mg

Equine-2.5 to 5 mg

EACH mL CONTAINS: 2 mg dexamethasone, 500 mg polyethylene glycol 400, 9 mg benzyl alcohol, 1.8 methylparaben and 0.2 mg propylparaben as preservatives, 4.75% alcohol, HCl to adjust pH to approximately 4.9, water for injection q.s.

For intravenous or intramuscular injection.

Usual Dose:

Bovine - 5 to 20 mg
Equine - 2.5 to 5 mg

Warning: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (59°F and 86°F).

Protect from freezing.

READ ACCOMPANYING DIRECTIONS CAREFULLY

Lot No. Exp. Date

NDC 13985-533-25

250 mL



Dexamethasone

Injection 2 mg/mL

Dexamethasone Sterile Injection

Veterinary
NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-324

V1 501112

MULTIPLE DOSE VIAL
Net Contents: 250 mL

EACH mL CONTAINS: 2 mg dexamethasone; 500 mg polyethylene glycol 400; 9 mg benzyl alcohol, 1.8 mg methylparaben and 0.2 mg propylparaben as preservatives; 4.75% alcohol; HCl to adjust pH to approximately 4.9; water for injection q.s.



Distributed by: MWI
Boise, ID 83705
www.VetOne.net

D-2953-05
Rev. 04/24



Cushing's syndrome in dogs has been reported in association with prolonged or repeated steroid therapy. Corticosteroids reportedly cause laminitis in horses.

CONTACT INFORMATION: To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet, contact Sparhawk Laboratories Inc. at 1-800-255-6388 or 1-913-888-7500. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

HOW SUPPLIED: DEXAMETHASONE INJECTION 2 mg/mL 100 mL - NDC 13985-533-03
250 mL - NDC 13985-533-25
multiple dose vials.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Protect from freezing.



Distributed by: MWI
Boise, ID 83705
www.VetOne.net Rev. 04/24

Approved by FDA under ANADA # 200-324

7

EACH mL CONTAINS: 2 mg dexamethasone; 500 mg polyethylene glycol 400; 9 mg benzyl alcohol, 1.8 mg methylparaben and 0.2 mg propylparaben as preservatives; 4.75% alcohol; HCl to adjust pH to approximately 4.9; water for injection q.s.



Distributed by: MWI
Boise, ID 83705
www.VetOne.net D-2953-05
Rev. 04/24



DEXAMETHASONE INJECTION 2 mg/mL

Solution for intravenous or intramuscular injection

Veterinary

**NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN**

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION: DEXAMETHASONE INJECTION 2 mg/mL is a synthetic analogue of prednisolone, having similar but more potent anti-inflammatory therapeutic action and diversified hormonal and metabolic effects. Modification of the basic corticoid structure as achieved in DEXAMETHASONE INJECTION 2 mg/mL offers enhanced anti-inflammatory effect compared to older corticosteroids. The dosage of DEXAMETHASONE INJECTION 2 mg/mL required is markedly lower than that of prednisolone and prednisolone.

DEXAMETHASONE INJECTION 2 mg/mL is not species-specific; however, the veterinarian should read the sections on **INDICATIONS, DOSAGE, SIDE EFFECTS, CONTRAINDICATIONS, PRECAUTIONS, and WARNINGS** before this drug is used.

1

DEXAMETHASONE INJECTION 2 mg/mL is intended for intravenous or intramuscular administration. Each mL contains 2 mg dexamethasone, 500 mg polyethylene glycol 400, 9 mg benzyl alcohol, 1.8 mg methylparaben and 0.2 mg propylparaben as preservatives, 4.75% alcohol, HCl to adjust pH to approximately 4.9, water for injection q.s.

EXPERIMENTAL STUDIES: Experimental animal studies on dexamethasone have revealed it possesses greater anti-inflammatory activity than many steroids. Veterinary clinical evidence indicates dexamethasone has approximately twenty times the anti-inflammatory activity of prednisolone and seventy to eighty times that of hydrocortisone. Thymus involution studies show dexamethasone possesses twenty-five times the activity of prednisolone. In reference to mineralocorticoid activity, dexamethasone does not cause significant sodium or water retention. Metabolic balance studies show that animals on controlled and limited protein intake will exhibit nitrogen losses on exceedingly high dosages.

INDICATIONS: DEXAMETHASONE INJECTION 2 mg/mL is indicated for the treatment of primary bovine ketosis and as an anti-inflammatory agent in the bovine and equine.

As supportive therapy, DEXAMETHASONE INJECTION 2 mg/mL may be used in the management of various rheumatic, allergic, dermatologic, and other diseases known to be responsive to anti-inflammatory

2

corticosteroids. DEXAMETHASONE INJECTION 2 mg/mL may be used intravenously as supportive therapy when an immediate hormonal response is required.

Bovine Ketosis

DEXAMETHASONE INJECTION 2 mg/mL is offered for the treatment of primary ketosis. The gluconeogenic effects of DEXAMETHASONE INJECTION 2 mg/mL, when administered intramuscularly, are generally noted within the first 6 to 12 hours. When DEXAMETHASONE INJECTION 2mg/mL is used intravenously, the effects may be noted sooner. Blood sugar levels rise to normal levels rapidly and generally rise to above normal levels within 12 to 24 hours. Acetone bodies are reduced to normal concentrations usually within 24 hours. The physical attitude of animals treated with DEXAMETHASONE INJECTION 2 mg/mL brightens and appetite improves, usually within 12 hours.

Milk production, which is suppressed as a compensatory reaction in this condition, begins to increase. In some instances, it may even surpass previous peaks. The recovery process usually takes from 3 to 7 days.

Supportive Therapy

DEXAMETHASONE INJECTION 2 mg/mL may be used as supportive therapy in mastitis, metritis, traumatic gastritis, and pyelonephritis, while appropriate primary therapy is administered. In these cases, the corticosteroid combats accompanying stress and enhances the feeling of general well-being.

3

DEXAMETHASONE INJECTION 2 mg/mL may also be used as supportive therapy in inflammatory conditions, such as arthritic conditions, snake bite, acute mastitis, shipping fever, pneumonia, laminitis, and retained placenta.

Equine

DEXAMETHASONE INJECTION 2 mg/mL is indicated for the treatment of acute musculoskeletal inflammations, such as bursitis, carpalitis, osselets, tendonitis, myositis, and sprains. If bony changes exist in any of the conditions, joints, or accessory structures, responses to DEXAMETHASONE INJECTION 2 mg/mL cannot be expected. In addition, DEXAMETHASONE INJECTION 2 mg/mL may be used as supportive therapy in fatigue, heat exhaustion, influenza, laminitis, and retained placenta provided that the primary cause is determined and corrected.

ADMINISTRATION AND DOSAGE: Therapy with DEXAMETHASONE INJECTION 2 mg/mL, as with any other potent corticosteroid, should be individualized according to the severity of the condition being treated, anticipated duration of steroid therapy, and the animal's threshold or tolerance for steroid excess.

Treatment may be changed over to DEXAMETHASONE INJECTION 2 mg/mL from any other glucocorticoid with proper reduction or adjustment of dosage.

Bovine - DEXAMETHASONE INJECTION 2 mg/mL - 5 to 20 mg intravenously or intramuscularly.

4

Dexamethasone Powder may be administered or the parenteral dose repeated as needed.

Equine - DEXAMETHASONE INJECTION 2 mg/mL - 2.5 to 5 mg intravenously or intramuscularly.

Dexamethasone Powder may be administered or the parenteral dose repeated as needed.

CONTRAINDICATIONS: Except for emergency therapy, do not use in animals with chronic nephritis and hypercorticalism (Cushing's syndrome). Existence of congestive heart failure, diabetes, and osteoporosis are relative contraindications. Do not use in viral infections during the viremic stage.

PRECAUTIONS: Animals receiving DEXAMETHASONE INJECTION 2 mg/mL should be under close observation. Because of the anti-inflammatory action of corticosteroids, signs of infection may be masked and it may be necessary to stop treatment until a further diagnosis is made. Overdosage of some glucocorticoids may result in sodium retention, fluid retention, potassium loss, and weight gain.

DEXAMETHASONE INJECTION 2 mg/mL may be administered to animals with acute or chronic bacterial infections providing the infections are controlled with appropriate antibiotic or chemotherapeutic agents.

Doses greater than those recommended in horses may produce a transient drowsiness or lethargy in some horses. The lethargy usually abates in 24 hours.

5

Use of corticosteroids, depending on dose, duration, and specific steroid, may result in inhibition of endogenous steroid production following drug withdrawal. In patients presently receiving or recently withdrawn from systemic corticosteroid treatments, therapy with a rapid acting corticosteroid should be considered in unusually stressful situations.

WARNINGS: Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate parturition followed by dystocia, fetal death, retained placenta, and metritis.

Additionally, corticosteroids administered to dogs, rabbits, and rodents during pregnancy have produced cleft palate. Other congenital anomalies including deformed forelegs phocomelia, and anasarca have been reported in offspring of dogs which received corticosteroids during pregnancy.

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

SIDE EFFECTS: Side effects, such as SAP and SGPT enzyme elevations, weight loss, anorexia, polydipsia, and polyuria have occurred following the use of synthetic corticosteroids in dogs. Vomiting and diarrhea (occasionally bloody) have been observed in dogs and cats.

6




Quantity: 12 Bottles Dexamethasone Injection
2mg/mL, 250mL (12 x 250 mL each)

V1 501112

Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Protect from freezing

Approved by FDA under ANADA # 200-324



30 3 13985 53325 0

Rev. 04/2024
D-2953-05C

DEXAMETHASONE

dexamethasone injection, solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:13985-533
Route of Administration	INTRAMUSCULAR, INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXAMETHASONE (UNII: 7S5I7G3JQL) (DEXAMETHASONE - UNII:7S5I7G3JQL)	DEXAMETHASONE	2 mg in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13985-533-03	100 mL in 1 VIAL, MULTI-DOSE		

2	NDC:13985-533-25	250 mL in 1 VIAL, MULTI-DOSE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANADA	ANADA200324		07/27/2012	

Labeler - MWI (Vet One) (019926120)

Registrant - Sparhawk Laboratories, Inc. (147979082)

Establishment			
Name	Address	ID/FEI	Business Operations
Sparhawk Laboratories, Inc.		147979082	analysis, manufacture

Establishment			
Name	Address	ID/FEI	Business Operations
Pharmacia & Upjohn Company LLC		618054084	api manufacture

Revised: 7/2024

MW (Vet One)