

**ACTICON- dextbrompheniramine maleate, pseudoephedrine hydrochloride tablet**  
**Actipharma, Inc**

-----  
**ACTICON® COLD & ALLERGY Tablets**

***Drug Facts***

***Active ingredients (in each tablet)***

Dexbrompheniramine Maleate, USP 2 mg  
Pseudoephedrine HCl, USP 60 mg

***Purpose***

Antihistamine  
Nasal Decongestant

***Uses***

Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose • sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

***Warnings***

**Do not exceed recommended dosage.**

***Do not use***

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

***Ask a doctor before use if you have***

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma • heart disease
- high blood pressure
- thyroid disease
- diabetes mellitus
- difficulty in urination due to enlargement of the prostate gland

**Do not take this product if you are taking sedatives or tranquilizers, without**

**first consulting your doctor.**

**When using this product**

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- if symptoms do not improve within 7 days or are accompanied by fever
- new symptoms occur

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

***Directions***

**Do not exceed recommended dosage.**

Adults and children 12 years of age and over:	1 tablet every 4 to 6 hours not to exceed 4 tablets in 24 hours or as directed by a doctor
Children 6 to under 12 years of age:	1/2 tablet every 4 to 6 hours not to exceed 2 tablets in 24 hours or as directed by a doctor
Children under 6 years of age	Consult a doctor

***Other information***

Store at 15°-30°C (59°-86°F) [see USP Controlled Room Temperature].

***Inactive ingredients***

Magnesium stearate, microcrystalline cellulose, sodium starch glycolate

***Questions or Comments?***

call weekdays from 8AM to 4PM AST at **1.787.608.0882**

**Contains the same active ingredients as Conex<sup>®</sup> Tablets\***

Tamper evident by foil seal under cap.  
Do not use if foil seal is broken or missing.

Manufactured in USA with imported ingredients for ActiPharma, Inc. San Juan, PR 00917. www.actipharma.net

**\*Conex® Tablets is a registered trademark of Llorens Pharmaceutical Corp. This product is not manufactured, distributed or marketed by Llorens Pharmaceutical Corp.**

**Packaging**

**FRONT PANEL**



**DRUG FACTS TABLE**

<p><b>Drug Facts</b></p>	
<p><b>Active ingredients (in each tablet)</b></p>	
<p>Dexbrompheniramine Maleate, USP 2 mg ....</p> <p>Pseudoephedrine HCl, USP 60 mg.....</p>	<p><b>Purpose</b></p> <p>Antihistamine</p> <p>Nasal Decongestant</p>
<p><b>Uses</b> Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:</p> <ul style="list-style-type: none"> <li>■ runny nose</li> <li>■ sneezing</li> <li>■ itching of the nose or throat</li> </ul>	

<p><b>Drug Facts (continued)</b></p>	
<p>In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.</p>	
<p><b>Directions</b></p> <p><b>Do not exceed recommended dosage.</b></p>	
<p>Adults and children 12 years of age and over:</p>	<p>1 tablet every 4 to 6 hours not to exceed 4 tablets in 24 hours or as directed by a doctor</p>
<p>Children 6 to 11 years of age:</p>	<p>1/2 tablet every 4 to 6 hours not to exceed 2 tablets in 24 hours or as directed by a doctor</p>

- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

### Warnings

**Do not exceed recommended dosage.**

#### Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

#### Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma ■ heart disease
- high blood pressure
- thyroid disease
- diabetes mellitus
- difficulty in urination due to enlargement of the prostate gland

**Do not take this product if you are taking sedatives or tranquilizers, without first consulting your doctor.**

#### When using this product

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- if symptoms do not improve within 7 days or are accompanied by fever
- new symptoms occur

**If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children.**

Children 6 to under 12 years of age:

1/2 tablet every 4 to 6 hours not to exceed 2 tablets in 24 hours or as directed by a doctor

Children under 6 years of age

Consult a doctor

### Other information

Store at 15°-30°C (59°-86°F) [see USP Controlled Room Temperature].

### Inactive ingredients

Magnesium stearate, microcrystalline cellulose, sodium starch glycolate

### Questions or Comments?

call weekdays from 8AM to 4PM AST at **1.787.608.0882**

Manufactured in USA with imported ingredients for ActiPharma, Inc. San Juan, PR 00917. [www.actipharma.net](http://www.actipharma.net)

**\*Conex® Tablets is a registered trademark of Llorens Pharmaceutical Corp. This product is not manufactured, distributed or marketed by Llorens Pharmaceutical Corp.**

Rev. 09/23

NDC 63102-401-60

**ACTICON®**  
COLD & ALLERGY

**Tablets**

Contains the same active ingredients as Conex® Tablets\*

**ANTIHISTAMINE • NASAL DECONGESTANT**

Each tablet contains:

Dexbrompheniramine Maleate ... 2 mg

Pseudoephedrine HCl ..... 60 mg

60 Tablets

Store at 59°-86°F (15°-30°C) [see USP Controlled Room Temperature]

Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing.

Manufactured for:

**tiPharma**  
Effective Solutions  
San Juan, PR 00917



## ACTICON

dexbrompheniramine maleate, pseudoephedrine hydrochloride tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63102-401
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXBROMPHENIRAMINE MALEATE</b> (UNII: BPA9UT29BS) (DEXBROMPHENIRAMINE - UNII:75T64B71RP)	DEXBROMPHENIRAMINE MALEATE	2 mg
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	60 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	13mm
<b>Flavor</b>		<b>Imprint Code</b>	A401
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63102-401-60	60 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/13/2015	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/13/2015	

**Labeler** - Actipharma, Inc (079340948)

Revised: 10/2023

Actipharma, Inc