

MILK FEVER CP- milk fever cp injection, solution
Aspen Veterinary

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Milk Fever CP

CAUTION:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For Animal Use Only
Keep Out of Reach of Children

NET CONTENTS:
500 mL (16.91 fl oz)

Parenteral Solution

INDICATIONS:

For intravenous use in milk fever, and in calcium, phosphorus, magnesium and glucose deficiency in animals.

NOTE:

Intravenous administration is always preferable. If used subcutaneously, always distribute dose in several places and massage to facilitate absorption.

DOSAGE:

Cattle, 500 mL. For milk fever prophylaxis, give 200-500 mL at time of calving. Horses, 250-500 mL; Sheep 50-125 mL. Repeat in 2 to 6 hours if needed. **This product contains no preservative. Unused portion remaining in bottle must be discarded.**

CAUTION:

Large doses administered intravenously may have toxic action on heart if the blood level of calcium

is raised excessively. Doses should be carefully regulated according to severity of hypocalcemia and injected slowly so that administration may be stopped if toxic action becomes evident.

CONTAINS:

Calcium borogluconate, dextrose, magnesium borogluconate, calcium hypophosphite and base.

Total calcium chemically equivalent to calcium borogluconate.....26.0% w/v
Dextrose · H₂O.....15.0% w/v
Magnesium borogluconate.....6.0% w/v
Total phosphorus in amounts chemically equivalent to elemental phosphorus.....0.5% w/v

Store between 15 degrees C-30 degrees C (59degrees F-86 degrees F).

TAKE TIME OBSERVE LABEL DIRECTIONS

Distributed by:

Aspen Veterinary Resources^R, Ltd.
Liberty, MO 64068 USA
www.aspenveterinaryresources.com

Manufactured by:

Nova-Tech, Inc.
Grand Island, NE 68801 USA 18-811 RMS# 92-565

Lot No. Exp. Date

Parenteral Solution

INDICATIONS:

For intravenous use in milk fever, and in calcium, phosphorus, magnesium and glucose deficiency in animals.

NOTE:

Intravenous administration is always preferable. If used subcutaneously, always distribute dose in several places and massage to facilitate absorption.

DOSAGE:

Cattle, 500 mL. For milk fever prophylaxis, give 200-500 mL at time of calving. Horses, 250-500 mL; Sheep 50-125 mL. Repeat in 2 to 6 hours if needed. **This product contains no preservative. Unused portion remaining in bottle must be discarded.**

CAUTION:

Large doses administered intravenously may have toxic action on heart if the blood level of calcium is raised excessively. Doses should be carefully regulated according to severity of hypocalcemia and injected slowly so that administration may be stopped if toxic action becomes evident.

NDC # 46066-811-50



Milk Fever CP™



CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**For Animal Use Only
Keep Out of Reach of Children**

**NET CONTENTS:
500 mL (16.91 fl oz)**

CONTAINS:

Calcium borogluconate, dextrose, magnesium borogluconate, calcium hypophosphite and base.

Total calcium chemically equivalent to calcium borogluconate26.0% w/v
Dextrose · H₂O15.0% w/v
Magnesium borogluconate6.0% w/v
Total phosphorus in amounts chemically equivalent to elemental phosphorus.....0.5% w/v

Store between 15°C-30°C (59°F-86°F).

Distributed by:

Aspen Veterinary Resources, Ltd.
Liberty, MO 64068 USA A219NT 04/15
www.aspenveterinaryresources.com

Manufactured by:

Nova-Tech, Inc.
Grand Island, NE 68801 USA 18-811 RMS# 92-565

TAKE TIME OBSERVE LABEL DIRECTIONS



Lot No. Exp. Date

milk fever cp injection, solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:46066-811
Route of Administration	INTRAVENOUS, SUBCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Calcium Borogluconate (UNII: 6915G2PQI0) (Calcium Cation - UNII:2M83C4R6ZB)	Calcium Borogluconate	130 g in 500 mL
Dextrose Monohydrate (UNII: LX22YL083G) (Anhydrous Dextrose - UNII:5SL0G7R0OK)	Dextrose Monohydrate	75 g in 500 mL
Magnesium Gluconate (UNII: T42NAD2KHC) (Magnesium Cation - UNII:T6V3LHY838)	Magnesium Gluconate	30 g in 500 mL
Phosphorus (UNII: 27YLU75U4W) (Phosphorus - UNII:27YLU75U4W)	Phosphorus	2.5 g in 500 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46066-811-50	500 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		03/20/2013	

Labeler - Aspen Veterinary (627265361)

Registrant - Aspen Veterinary (627265361)

Revised: 9/2025

Aspen Veterinary