

**PROFEN IB INFANTS- ibuprofen suspension**  
**Publix Super Markets Inc**

-----  
**Publix Super Markets, Inc. Profen IB Drops Drug Facts**

**Active ingredient (in each 1.25 mL)**

Ibuprofen 50 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

### **Do not use**

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

### **Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- give with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
  - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area

- any new symptoms appear

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- mL = milliliter
- measure with the dosing device provided. Do not use with any other device.
- dispense liquid slowly into the child’s mouth, toward the inner cheek
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**

<b>Dosing Chart</b>		
<b>Weight (lbs)</b>	<b>Age (mos)</b>	<b>Dose (mL)</b>
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

**Other information**

- store at 68-77°F (20-25°C)
- do not freeze
- **do not use if printed neckband is broken or missing**

**Inactive ingredients**

anhydrous citric acid, FD&C red #40, glycerin, hypromellose, natural and artificial berry flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

**Principal Display Panel**

P  
 FOR AGES 6 TO 23 MONTHS  
 INFANTS’  
 profen IB drops  
 CONCENTRATED DROPS IBUPROFEN ORAL SUSPENSION 50 mg PER 1.25 mL

Pain Reliever

Fever Reducer (NSAID)

No High Fructose Corn Syrup

USE ONLY WITH ENCLOSED SYRINGE

• Lasts up to 8 hours

BERRY FLAVOR

Compare to Motrin® Infants' Drops active ingredient.

0.5 FL OZ (15 mL)

**FOR AGES 6 TO 23 MONTHS INFANTS'**  
**profenlBdrops**  
CONCENTRATED DROPS  
IBUPROFEN ORAL SUSPENSION  
50 mg PER 1.25 mL

**Pain Reliever  
Fever Reducer (NSAID)  
No High Fructose Corn Syrup**

• Lasts up to 8 hours

**BERRY FLAVOR**

0.5 FL OZ (15 mL)

USE ONLY WITH ENCLOSED SYRINGE

Compare to Motrin® Infants' Drops active ingredient.

**ACTUAL SIZE**

**Warning:** Do not use if child has had an allergic reaction to ibuprofen or any other pain reliever/fever reducer right before or after heart surgery.

**Directions:** Give with food or milk if stomach upset occurs.

**Stop use and ask a doctor if:**

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Drug Facts**

**Active ingredient (in each 1.25 mL)** Ibuprofen 50 mg (NSAID)\*

**Purposes** Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses temporarily:**

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Drug Facts (continued)**

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- give with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Drug Facts (continued)**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- mL = milliliter
- measure with the dosing device provided. Do not use with any other device.
- dispense liquid slowly into the child's mouth, toward the inner cheek
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day

**Dosing Chart**

Weight (lbs)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

**Other information**

- store at 68-77°F (20-25°C)
- do not freeze
- do not use if printed neckband is broken or missing

**Inactive ingredients** anhydrous citric acid, FD&C red #40, glycerin, hypromellose, natural and artificial berry flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

DISTRIBUTED BY  
PUBLIX SUPER MARKETS, INC.,  
3300 PUBLIX CORPORATE PARKWAY,  
LAKELAND, FL 33811 [publix.com](http://publix.com)  
1-888-267-3037

**PUBLIX GUARANTEE:**  
COMPLETE SATISFACTION  
OR YOUR MONEY BACK

**Publix.**

05705 63 C11

# PROFEN IB INFANTS

ibuprofen suspension

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:56062-057
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	50 mg in 1.25 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Product Characteristics

<b>Color</b>	PINK (light)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56062-057-05	1 in 1 CARTON	06/29/2001	
1		15 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

ANDA	ANDA075217	06/29/2001	
------	------------	------------	--

**Labeler** - Publix Super Markets Inc (006922009)

Revised: 7/2025

Publix Super Markets Inc