

**WAL ITIN- loratadine tablet**  
**Walgreen Company**

-----  
**Walgreen Co. Wal-itin® Drug Facts**

**Active ingredient (in each tablet)**

Loratadine 10 mg

**Purpose**

Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

**Warnings**

**Do not use**

if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

liver or kidney disease. Your doctor should determine if you need a different dose.

**When using this product**

do not take more than directed. Taking more than directed may cause drowsiness.

**Stop use and ask a doctor if**

an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

## Other information

- do not use if blister unit is broken or torn
- store between 20° to 25° C (68° to 77° F)
- protect from excessive moisture

## Inactive ingredients

Lactose Monohydrate, Magnesium Stearate, Providone

## Questions or comments?

**1-800-719-9260**

## Principal Display Panel

Well at Walgreens

NON-DROWSY\*

ORIGINAL PRESCRIPTION STRENGTH

Wal-itin®

24 Hour Allergy

Loratadine Tablets 10 mg • Antihistamine

- 24-hour relief of sneezing, runny nose, itchy, watery eyes & itchy throat or nose
- Indoor & outdoor allergies

60 TABLETS

24 HOUR

\*WHEN TAKEN AS DIRECTED. SEE DRUG FACTS PANEL.

**Drug Facts (continued)**

- you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech
- leg swelling, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 28 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**100% Satisfaction Guarantee**  
(888) 308-9039

**Drug Facts (continued)**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** do not take more than directed. The smallest effective dose should be used.

adults and children 12 years and older

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

ask a doctor

**Compare to the active ingredient of Advil® Ibuprofen Tablets\*\***

**DG health**

**Ibuprofen**  
Tablets, 200 mg  
Pain Reliever/  
Fever Reducer (NSAID)

200 mg      1000 Tablets  
Actual Tablet Size

**Drug Facts**

**Active ingredient (in each tablet)** Ibuprofen 200 mg (NSAID)\*. \*Pain reliever/fever reducer. \*\*nonsteroidal anti-inflammatory drug.

**Purposes**

- temporarily relieves minor aches and pains due to: headache, muscle aches, minor pain of arthritis, toothache, backache, the common cold, menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters.

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or blood drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.

**Drug Facts (continued)**

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer, right before or after heart surgery.

**Ask a doctor before use if:** you have problems or serious side effects from taking pain relievers or fever reducers, the stomach bleeding warning applies to you, you have a history of stomach problems, such as heartburn, you have high blood pressure, heart disease, liver or kidney disease, asthma, or had a stroke, you are taking a diuretic.

**Ask a doctor or pharmacist before use if you are:** taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin, under a doctor's care for any serious condition, taking any other drug.

**When using this product:** take with food or milk if stomach upset occurs.

**Stop use and ask a doctor if:** you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better.

**EXP. LOT**  
1604K1 VT F3

## WAL ITIN

loratadine tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-0699
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJ03BO7QN) (LORATADINE - UNII: 7AJ03BO7QN)	LORATADINE	10 mg

### Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	L612
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0699-46	10 in 1 CARTON	02/13/2009	09/30/2023
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

2	NDC:0363-0699-60	20 in 1 CARTON	02/19/2009	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:0363-0699-65	1 in 1 CARTON	02/17/2009	06/30/2023
3		30 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0363-0699-95	1 in 1 CARTON	05/01/2013	10/31/2023
4		45 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:0363-0699-72	1 in 1 CARTON	03/10/2009	
5		60 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:0363-0699-03	1 in 1 CARTON	02/04/2014	
6		70 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:0363-0699-75	1 in 1 CARTON	03/10/2009	10/31/2023
7		90 in 1 BOTTLE; Type 0: Not a Combination Product		
8	NDC:0363-0699-47	1 in 1 CARTON	03/21/2009	07/31/2023
8		150 in 1 BOTTLE; Type 0: Not a Combination Product		
9	NDC:0363-0699-87	1 in 1 CARTON	02/04/2011	09/30/2023
9		300 in 1 BOTTLE; Type 0: Not a Combination Product		
10	NDC:0363-0699-58	1 in 1 CARTON	08/22/2016	08/31/2023
10		40 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076301	02/13/2009	

**Labeler** - Walgreen Company (008965063)

Revised: 12/2023

Walgreen Company