

**MUCINEX CHILDRENS FREEFROM MULTI-SYMPATOM COLD AND FLU NIGHTTIME- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, and triprolidine hydrochloride solution**  
**RB Health (US) LLC**

-----  
**Mucinex® Children's FreeFrom Multi-Symptom Cold & Flu Nighttime**

***Drug Facts***

<b><i>Active ingredients (in each 10 mL)</i></b>	<b><i>Purposes</i></b>
<b>Acetaminophen 325 mg</b>	<b>Pain reliever/fever reducer</b>
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant
Tripolidine HCl 1.25 mg	Antihistamine

**Uses**

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - runny nose
  - sneezing
  - cough
  - stuffy nose
- temporarily reduces fever
- controls cough to help your child get to sleep

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 1625 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

## **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleepy
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

## **Ask a doctor before use if the child has**

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

## **Ask a doctor or pharmacist before use if the child is**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

## **When using this product**

- **do not use more than directed (see Overdose warning)**
- excitability may occur, especially in children
- may cause marked drowsiness
- sedatives and tranquilizers may increase drowsiness

## **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

## **Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

### **Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see Overdose warning)**
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 4 doses in any 24-hour period
- children under 6 years of age: do not use

### **Other information**

- each 10 mL contains: **sodium 6 mg**
- store at 20-25°C (68-77°F)
- do not refrigerate

### **Inactive ingredients**

ammonium glycyrrhizate, anhydrous citric acid, cochineal carmine, edetate disodium, flavors, glycerin (soy), maltodextrin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

### **Questions?**

**1-866-MUCINEX (1-866-682-4639)**

You may also report side effects to this phone number.

**Dist. by: RB Health (US)  
Parsippany, NJ 07054-0224**

### **PRINCIPAL DISPLAY PANEL - 118 mL Bottle Carton**

Pediatrician Recommended †

NIGHTTIME

NDC 63824-959-64

Mucinex®  
Children's

FREEFROM

MULTI-SYMP TOM  
COLD & FLU

NIGHTTIME

ACETAMINOPHEN – PAIN RELIEVER/FEVER REDUCER

DEXTROMETHORPHAN HBr – COUGH SUPPRESSANT

PHENYLEPHRINE HCl – NASAL DECONGESTANT

TRIPROLIDINE HCl – ANTIHISTAMINE

FREEFROM

ARTIFICIAL FLAVORS, COLORS, & DYES • ALCOHOL & SUGAR

ELDERBERRY

& CHERRY NATURAL FLAVOR

AGES

6+ YRS.

4 FL OZ (118 mL)

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

**Mucinex<sup>®</sup>**  
**Children's**

**FREEFROM**  
MULTI-SYMPOM  
**COLD & FLU NIGHTTIME**  
ACETAMINOPHEN – PAIN RELIEVER/FEVER REDUCER  
DEXTROMETHORPHAN HBr – COUGH SUPPRESSANT  
PHENYLEPHRINE HCl – NASAL DECONGESTANT  
TRIPROLODINE HCl – ANTIHISTAMINE

### Drug Facts (continued)

#### When using this product

- do not use more than directed (see Overdose warning)
- excitability may occur, especially in children
- may cause marked drowsiness
- sedatives and tranquilizers may increase drowsiness

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

#### Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

#### Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 4 doses in any 24-hour period
- children under 6 years of age: do not use

#### Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

**Inactive ingredients** ammonium glycyrrhizate, anhydrous citric acid, cochineal camine, edetate disodium, flavors, glycerin (soy), maltodextrin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

**Questions?** 1-866-MUCINEX (1-866-682-4639)

Pediatrician Recommended†

**NIGHTTIME**

NDC 63824-959-64

**Mucinex<sup>®</sup>**  
**Children's**

**FREEFROM**

MULTI-SYMPOM  
**COLD & FLU**  
**NIGHTTIME**

ACETAMINOPHEN – PAIN RELIEVER/FEVER REDUCER  
DEXTROMETHORPHAN HBr – COUGH SUPPRESSANT  
PHENYLEPHRINE HCl – NASAL DECONGESTANT  
TRIPROLODINE HCl – ANTIHISTAMINE





You may also report side effects to this phone number.

**\*vs Private Label OTC**  
as an OTC brand in the Children's Cough/Cold category  
(AlphaImpactRxProVoice Survey)

Dist. by: RB Health (US)  
Parsippany, NJ 07054-0224  
©2020 RB Health

010820 3132570 US Patents: [www.rb.com/patents](http://www.rb.com/patents)

Keep carton for full information.

**FREEFROM**

ARTIFICIAL FLAVORS, COLORS, & DYES • ALCOHOL & SUGAR

# ELDERBERRY

& CHERRY NATURAL FLAVOR



AGES  
6+ YRS.

4 FL OZ (118 mL)



3 63824 60070 3

LOT CODE:

EXP. DATE:

MADE IN:



3132570



Pharma  
Code

**What this has is just as important as what it doesn't.**

Leading Children's OTC Cough & Cold brand\* with the effective symptom

### Drug Facts

**Active ingredients (in each 10 mL)**

**Purposes**

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant
Triprolidine HCl 1.25 mg.....	Antihistamine

relief you trust – FREEFROM the unwanted additives:

Caffeine Free · No Artificial Colors, Dyes, or Flavors · Alcohol & Sugar Free  
No High Fructose Corn Syrup

## POWERFUL SYMPTOM RELIEF

- ✓ Nighttime Relief for a Better Morning
- ✓ Relieves Sore Throat & Stuffy Nose
- ✓ Controls Cough
- ✓ Relieves Runny Nose & Sneezing
- ✓ Reduces Fever

DOSING CUP INCLUDED



**ELDERBERRY**  
& CHERRY NATURAL FLAVOR

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



Questions or comments?  
1-866-Mucinex (1-866-682-4639)  
or [www.mucinex.com](http://www.mucinex.com)

HEALTH > HYGIENE > HOME

### Uses

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - sore throat
  - runny nose
  - cough
  - headache
  - nasal congestion
  - sneezing
  - stuffy nose
- temporarily reduces fever
- controls cough to help your child get to sleep

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 1625 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
  - blisters
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleepy
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

### Ask a doctor before use if the child has

- liver disease
- diabetes
- thyroid disease
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)
- heart disease
- high blood pressure
- glaucoma

### Ask a doctor or pharmacist before use if the child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers



# MUCINEX CHILDRENS FREEFROM MULTI-SYMPATOM COLD AND FLU NIGHTTIME

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, and triprolidine hydrochloride solution

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63824-959
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 10 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 10 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 10 mL
<b>TRIPROLIDINE HYDROCHLORIDE</b> (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	1.25 mg in 10 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>AMMONIUM GLYCYRRHIZATE</b> (UNII: 3VRD35U26C)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>CARMINIC ACID</b> (UNII: CID8Z8N95N)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging



#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63824-959-64	1 in 1 CARTON	06/01/2020	
1		118 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2020	

**Labeler** - RB Health (US) LLC (081049410)

Revised: 12/2024

RB Health (US) LLC