

**PEPTO-BISMOL DIARRHEA- bismuth subsalicylate tablet**  
**Procter & Gamble Manufacturing México, S. de R.L. de C.V.**

-----  
**Pepto Bismol<sup>®</sup> Diarrhea Caplets**

***Drug Facts***

**Active ingredient (in each caplet)**

Bismuth subsalicylate 525 mg

**Purpose**

Antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea

**Warnings**

**Reye's syndrome:**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:**

Contains salicylate .

Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist before use if you are**

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

### **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur

### **Stop use and ask a doctor if**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

### **If pregnant or breast feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- swallow with water, do not chew
- adults and children 12 years and over:
- 1 caplet every ½ hour or 2 caplets every hour as needed
- do not exceed 8 caplets in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### **Other information**

- each caplet contains: calcium 54 mg
- salicylate 199 mg
- very low sodium
- avoid excessive heat (over 104°F or 40°C)

### **Inactive ingredients**

calcium carbonate, D&C Red No. 27 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, polysorbate 80, povidone, silicon dioxide, sodium starch glycolate

### **Questions?**

1-800-717-3786

**TAMPER EVIDENT: Do not use if inner printed seal on bottle is broken or missing.**

**MADE IN MEXICO  
DIST. BY PROCTER & GAMBLE,  
CINCINNATI OH 45202**

**PRINCIPAL DISPLAY PANEL - 12 Caplet Carton**

**Pepto<sup>®</sup>  
Diarrhea**

Bismuth Subsalicylate

Antidiarrheal

**2x CONCENTRATED FORMULA\***

COATS & KILLS BACTERIA THAT CAUSE DIARRHEA

**12 CAPLETS**

TAMPER EVIDENT: Do not use if inner printed seal on bottle is broken or missing.

# Pepto Bismol<sup>®</sup> Diarrhea

## Pepto<sup>®</sup> Diarrhea

Bismuth Subsalicylate Antidiarrheal

2x  
CONCENTRATED  
FORMULA\*



**COATS & KILLS  
BACTERIA THAT  
CAUSE DIARRHEA**

12 Caplets



### Drug Facts (continued)

- anticoagulation (thinning the blood)
- diabetes • gout • arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

#### Stop use and ask a doctor if

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- swallow with water, do not chew
- adults and children 12 years and over:
  - 1 caplet every ½ hour or 2 caplets every hour as needed
- do not exceed 8 caplets in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years; ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### Drug Facts (continued)

#### Other information

- each caplet contains: calcium 54 mg
- salicylate 199 mg
- very low sodium
- avoid excessive heat (over 104°F or 40°C)

**Inactive ingredients** calcium carbonate, D&C Red No. 27 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, polysorbate 80, povidone, silicon dioxide, sodium starch glycolate

**Questions?** 1-800-717-3786

## Pepto<sup>®</sup> Diarrhea

MADE IN MEXICO  
DIST. BY PROCTER & GAMBLE,  
CINCINNATI OH 45202

**P&G**

www.pg.com www.pepto-bismol.com  
Patents: www.pg.com/patents

Also sold as Pepto Bismol Ultra<sup>®</sup>

\*vs Original Pepto Bismol<sup>®</sup>

91194130

### Drug Facts

Active ingredient (in each caplet)	Purpose
Bismuth subsalicylate 525 mg...	Antidiarrheal

Bismuth subsalicylate 525 mg... Antidiarrheal

**Uses** relieves • travelers' diarrhea • diarrhea

#### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate.**

**Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have** • an ulcer

- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever • mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for**

## PEPTO-BISMOL DIARRHEA

bismuth subsalicylate tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58933-391
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>D&amp;C RED NO. 27 ALUMINUM LAKE</b> (UNII: ZK64F7XSTX)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

## Product Characteristics

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	OVAL (Caplet)	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	Pepto
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-391-12	1 in 1 CARTON	03/01/2019	06/30/2026
1		12 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	03/01/2018	06/30/2026

**Labeler** - Procter & Gamble Manufacturing México, S. de R.L. de C.V. (812807550)

Revised: 10/2024

Procter & Gamble Manufacturing México, S. de R.L. de C.V.