

TYLENOL EXTRA STRENGTH- acetaminophen tablet, film coated
Morning Star OTC

TYLENOL Extra Strength

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert:acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 caplets every 6 hours while symptoms last▪ do not take more than 6 caplets in 24 hours, unless directed by a doctor▪ do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store between 20-25°C (68-77°F)
- **do not use if carton is opened. Do not use if foil inner seal imprinted with "TYLENOL" is broken or missing**

Inactive ingredients

carnauba wax ¹, corn starch ¹, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, modified starch ¹, polyethylene glycol ¹, powdered cellulose, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, titanium dioxide

1 contains one or more of these ingredients

Questions or comments?

call **1-877-895-3665**(toll-free) or **215-273-8755**(collect)

Repackaged by:

Morning Star OTC

145 S. Anderson St, Los Angeles,

CA 90033

PRINCIPAL DISPLAY PANEL-2 Tablets



Distributed by: JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2020
Visit us at www.tylenol.com
or call toll-free 1-877-TYLENOL
(1-877-895-3665)

This product is repackaged by: Morning Star OTC
145 S. Anderson St., Los Angeles, CA 90033
Hours of operation 9 am - 4 pm, 323.354.4838

DO NOT USE IF INNER POUCH OR CARTON IS OPEN OR DAMAGED!

Drug Facts	
Active ingredient (in each caplet)	Purpose
Acetaminophen 500mg	Pain reliever/fever reducer
Uses	
temporarily relieves minor aches and pains due to:	
the common cold	headache
backache	minor pain of arthritis
toothache	muscular aches
premenstrual and menstrual cramps	
temporarily reduces fever	
Warnings	
Liver Warning: This product contains acetaminophen. Severe liver damage may occur if you take:	
more than 4,000 mg of acetaminophen in 24 hours	
with other drugs containing acetaminophen	
5 or more alcoholic drinks every day while using this product	
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash	
If a skin reaction occurs, stop use and seek medical help right away.	
Do not use:	
with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	
If you are allergic to acetaminophen or any of the inactive ingredients in this product.	
Ask a doctor or pharmacist before use if you have liver disease.	
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.	

LIFT PANEL FOR CONTINUED DRUG FACTS

Drug Facts (continued)	
Stop use and ask a doctor if:	
pain gets worse or lasts more than 10 days	
fever gets worse or lasts more than 3 days	
new symptoms occur	
redness or swelling is present	
These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children.	
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-522-1223). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions	
do not take more than directed (see overdose warning)	
adults and children 12 years and over	
take 2 caplets every 6 hours while symptoms last.	
do not take more than 6 caplets in 24 hours, unless directed by a doctor.	
do not use for more than 10 days unless directed by a doctor.	
children under 12 years	
ask a doctor	
Other information	
store between 20-25°C (68-77°F)	
do not use if pouch is torn or damaged	
Inactive ingredients	
carnauba wax*, corn starch*, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, modified starch*, polyethylene glycol*, powdered cellulose, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, titanium dioxide	
*contains one or more of these ingredients	
Questions?	
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)	

PRINCIPAL DISPLAY PANEL-2 Tablets x 25 Pouches



TYLENOL EXTRA STRENGTH

acetaminophen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53209-1001(NDC:50580-449)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ALUMINUM OXIDE (UNII: LMI26O6933)	

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	TYLENOL;500
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53209-1001-2	25 in 1 POUCH	06/19/2025	
1	NDC:53209-1001-1	2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/19/1984	

Labeler - Morning Star OTC (078589357)

Registrant - Morning Star OTC (078589357)

Establishment

Name	Address	ID/FEI	Business Operations
Morning Star OTC		078589357	repack(53209-1001)