

**PAIN RELIEF PM- acetaminophen and diphenhydramine hydrochloride tablet,
coated
Meijer**

MEI-1095-2019-0904

Drug Facts

| <i>Active ingredients (in each caplet)</i> | <i>Purpose</i> |
|---|------------------------|
| Acetaminophen 500 mg | Pain reliever |
| Diphenhydramine HCl 25 mg | Nighttime sleep aid |

Uses

temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease

- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

| | |
|---------------------------------------|--|
| adults and children 12 years and over | <ul style="list-style-type: none"> ▪ take 2 caplets at bedtime ▪ do not take more than 2 caplets of this product in 24 hours |
| children under 12 years | <ul style="list-style-type: none"> ▪ do not use |

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

colloidal silicon dioxide, copovidone, croscarmellose sodium, FD&C blue #1, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

1-844-705-4384

PRINCIPAL DISPLAY PANEL

NDC 41250-195-02

Compare to Extra Strength Tylenol® PM active ingredients†

Meijer®

For Adults

pain relief PM


Acetaminophen, Diphenhydramine HCl

Extra Strength

Pain Reliever, Nighttime Sleep Aid

50 CAPLETS

actual size



PAIN RELIEF PM

acetaminophen and diphenhydramine hydrochloride tablet, coated

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:41250-195 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------------------|-----------------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 500 mg |
| DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M) | DIPHENHYDRAMINE HYDROCHLORIDE | 25 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|-----------------|
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| COPOVIDONE K25-31 (UNII: D9C330MD8B) | |
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| HYPROMELLOSES (UNII: 3NXW29V3WO) | |
| LACTOSE (UNII: J2B2A4N98G) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) | |
| STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ) | |
| SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |

Product Characteristics

| | | | |
|-----------------|------|---------------------|----------|
| Color | blue | Score | no score |
| Shape | OVAL | Size | 17mm |
| Flavor | | Imprint Code | AAA;1031 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|----------|------------------|---|-----------------------------|---------------------------|
| 1 | NDC:41250-195-02 | 1 in 1 CARTON | 05/01/2017 | 09/30/2026 |
| 1 | | 50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |
| 2 | NDC:41250-195-03 | 1 in 1 CARTON | 05/01/2017 | 09/30/2026 |
| 2 | | 100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |
| 3 | NDC:41250-195-09 | 2 in 1 CARTON | 05/01/2017 | 09/30/2026 |
| 3 | | 100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M013 | 05/01/2017 | 09/30/2026 |

Labeler - Meijer (006959555)

Revised: 10/2024

Meijer