

GILTUSS MULTISYMPATOM COLD AND FLU- acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hydrochloride solution
Giltuss Pharmaceutical Corp

Giltuss Multi-Symptom Cold & Flu

Drug Facts

Active ingredients (in each 10 mL)

Acetaminophen 650 mg

Chlorpheniramine maleate 4 mg

Dextromethorphan hydrobromide 13 mg

Phenylephrine hydrochloride 10 mg

Purposes

Pain reliever/fever reducer

Antihistamine

Cough Suppressant

Nasal decongestant

Uses

- temporarily relieves:
- minor aches and pains
- headache
- minor sore throat pain
- runny nose
- itchy nose or throat
- sneezing
- itchy watery eyes due to hay fever
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 6 doses in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are allergic to acetaminophen or any inactive ingredient in this product
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor or pharmacist before use if the user has

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- chronic cough that lasts such as occurs with smoking, asthma, or emphysema
- trouble urinating due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if the user is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- marked drowsiness may occur
- excitability may occur, especially in children
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you get nervousness, dizziness or sleeplessness
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

- new symptoms occur
- cough comes back or occurs with fever, rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not give more than directed
- use enclosed dosing cup
- keep dosing cup for use with this product
- mL = milliliter

adults and children 12 years and over	10 mL every 4-6 hours, not to exceed 6 doses in 24 hours
children 6 to under 12 years of age	5 mL every 4-6 hours, not to exceed 5 doses in 24 hours
children under 6 years of age	Do not use

Other information

- store at room temperature 15° - 30°C (59° - 86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients citric acid, disodium EDTA, flavor, glycerin, methylparaben, polyethylene glycol, propylene glycol, propylparaben, purified water and sucralose

Questions or comments?

Call **1-787-848-9114**, Monday-Friday 9AM - 5PM EST. Call your doctor for medical advice in the event of side effects.

Tamper evident: do not use if safety seal is broken or missing

MANUFACTURED FOR

GIL PHARMACEUTICAL CORP.

PONCE, PUERTO RICO 00717-1565

MAXIMUM STRENGTH

Giltuss® MULTI-SYMPTOM

COLD & FLU

ANALGESIC, ANTITUSSIVE, ANTIHISTAMINE & NASAL DECONGESTANT

ALCOHOL FREE. SODIUM FREE. SUGAR FREE. DYE FREE

RELIEVES

- FEVER, HEADACHE AND BODY ACHES

- COUGH AND COLD
- SNEEZING
- RUNNY NOSE
- ITCHY AND WATERY EYES
- STUFFY NOSE
- SINUS PRESSURE

STRAWBERRY FLAVOR

4FL OZ (118 mL)

Drug Facts

Active Ingredients (In each 15 mL)	Purpose
Acetaminophen 150 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 4 mg	Antihistamine
Dextromethorphan HBr 12 mg	Cough Suppressant
Pseudoephedrine HCl 10 mg	Nasal Decongestant

Uses

- temporarily relieves
 - minor aches and pains
 - minor sore throat pain
 - itchy nose or throat
 - itchy, watery eyes due to hay fever
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation as may occur with a cold
 - temporarily reduces fever
- headache
- runny nose
- sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 6 doses in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are allergic to acetaminophen or any inactive ingredient in this product
- with any other drug containing acetaminophen (prescription or over-the-counter)

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are taking a prescription monoamine oxidase inhibitor (MAOI) for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor or pharmacist before use if the user has:

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- chronic cough that lasts such as occurs with smoking, asthma or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if the user is:

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- marked drowsiness may occur
- excitability may occur, especially in children
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1585

giltuss.com

Drug Facts (continued)

Stop use and ask a doctor if:

- you get nervousness, dizziness or sleeplessness
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- nasal or swelling is present
- new symptoms occur
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not give more than directed
- use enclosed dosing cup
- keep dosing cup for use with this product
- 6L-milliliters

adults and children 12 years and over	10 mL every 4-6 hours, not to exceed 6 doses in 24 hours.
children 6 to under 12 years of age	5 mL every 4 to 6 hours, not to exceed 5 doses in 24 hours.
children under 6 years of age	Do not use

Other information

- store at room temperature 15°-30°C (59°-86°F)
- close caps tightly
- SAVE CARTON FOR COMPLETE DRUG FACTS

Inactive Ingredients
Citric acid, disodium EDTA, flavor, glycerin, methylparaben, polyethylene glycol, propylene glycol, propylparaben, purified water and sucralose.

Questions or comments?
Call 1-787-848-0114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

Tamper evident: do not use if safety seal is broken or missing

DO NOT USE IF SAFETY SEAL IS BROKEN OR MISSING.
NO LO USE SI LA TRILLA EN LA TAPA ESTÁ ROTA O FALTA.

G

ANALGESIC, ANTITUSSIVE, ANTIHISTAMINE & NASAL DECONGESTANT
ANALGÉSICO, ANTITUSSIVO, ANTIHISTAMÍNICO & DESCONGESTIONANTE NASAL

**MAXIMUM STRENGTH
MÁXIMA CONCENTRACIÓN**

G

**Giltuss.
MULTI-SYMP TOM
COLD & FLU
MULTI-SÍNTOMA
CATARRO Y GRIPE**

Analgesic effectiveness and proven safety to reduce fever, headache, and body aches
Efectividad analgésica y seguridad comprobada para reducir la fiebre, dolor de cabeza y dolores de cuerpo

Antitussive strength to control cough and cold symptoms
Fuerza antitussiva para controlar la tos y el catarro

Fast antihistamine efficacy to relieve sneezing, runny nose, and itchy and watery eyes
Rápida eficacia antihistamínica para el alivio de estornudos, goteo nasal y picor y ojos llorosos

Decongestant efficacy to relieve sinus pressure and nasal congestion
Eficacia descongestionante para el alivio de presión por sinusitis y congestión nasal

RELIEVES ALIVIA

FEVER	✓ FIEBRE, DOLOR DE CABEZA Y CUERPO
HEADACHE AND BODY ACHES	✓ TOS Y CATARRO
COUGH AND COLD	✓ SNEEZING
✓ SNEEZING	✓ ESTORNUDOS
✓ RUNNY NOSE	✓ GOTEO NASAL
✓ ITCHY AND WATERY EYES	✓ PICOR Y OJOS LLOROSOS
✓ STUFFY NOSE	✓ NARIZ TAPADA
✓ SINUS PRESSURE	✓ PRESIÓN POR SINUSITIS

ALCOHOL FREE. SODIUM FREE. SUGAR FREE. DYE FREE.
SIN ALCOHOL. SIN SODIO. SIN AZÚCAR. SIN COLORANTES.

ALCOHOL FREE. SODIUM FREE. SUGAR FREE. DYE FREE.
SIN ALCOHOL. SIN SODIO. SIN AZÚCAR. SIN COLORANTES.

ALCOHOL FREE. SODIUM FREE. SUGAR FREE. DYE FREE.
SIN ALCOHOL. SIN SODIO. SIN AZÚCAR. SIN COLORANTES.

STRAWBERRY FLAVOR
SABOR A FRESA.

4FL OZ (118 mL)



N 5855-2136-04 4

Exp. Date:

GILTUSS MULTISYMPPTOM COLD AND FLU

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hydrochloride solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58552-136
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	13 mg in 10 mL
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
METHYLPARABEN (UNII: A2I8C7HI9T)	

Product Characteristics

Color	yellow (Light yellow)	Score	
Shape		Size	
Flavor	STRAWBERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58552-136-04	1 in 1 CARTON	10/02/2018	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/02/2018	

Labeler - Giltuss Pharmaceutical Corp (176826592)

Establishment

Name	Address	ID/FEI	Business Operations
Dextrum Laboratories Inc		007392322	manufacture(58552-136)

Revised: 10/2025

Giltuss Pharmaceutical Corp