

CHLORAPREP ONE-STEP- chlorhexidine gluconate and isopropyl alcohol solution
CareFusion 213, LLC

BD Chloraprep™ FREPP™ Clear 1.5mL Applicators
Sterile Solution

Active ingredients

Chlorhexidine gluconate 2% w/v

Isopropyl alcohol 70% v/v

Purposes

Antiseptic

Antiseptic

Use

for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only.

Flammable, keep away from fire or flame.

- do not use with electrocautery procedures

Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing • shock • facial swelling • hives • rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product

keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain.

If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if

irritation, sensitization, or allergic reaction occurs.

These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age.

These products may cause irritation or chemical burns.

- maximal treatment area for one applicator is approximately **2.5 in. x 2.5 in. (42 cm²)**
- remove applicator from package; do not touch sponge
- hold the applicator horizontally with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds. Allow the area to air dry for approximately 30 seconds.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes. Allow the area to air dry for approximately 1 minute.
- do not blot or wipe away
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other Information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive Ingredient

USP purified water

Questions?

• www.chloraprep.com • call **1-800-523-0502** (M-F 8 a.m.-5 p.m. CST)

BD Chloraprep™ FREPP™ Clear 1.5 mL Applicator Principal Display Panel-Package Label

Principal Display Panel-Package Label

BD Chloraprep™ FREPP™ Clear

1.5 mL Applicator

2% w/v chlorhexidine gluconate (CHG)

and 70% v/v isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

Sterile Solution

Applicator is sterile if package is intact

NDC 54365-400-30

REF 930299

 **BD Chloraprep™**
FREPP™ Clear
1.5 mL Applicator
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Sterile Solution
Applicator is sterile if package is intact
WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.
CareFusion 213, LLC, El Paso, TX 79912,
subsidiary of Becton, Dickinson and Co.
1-800-523-0502 (M-F 8 a.m.- 5 p.m. CST)
Artwork No. 6-930213 • A050619
Made in the USA of US and imported parts

REF 930299
NDC 54365-400-30

 **STERILE SOLUTION**

  

STERILE | **EO**

Bend and Peel

BD Chloraprep™ FREPP™ Clear 1.5 mL Applicators Principal Display Panel-Carton

Principal Display Panel-Carton
 BD Chloraprep™ FREPP™ Clear
 1.5 mL Applicators
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
 Sterile Solution
 Applicator is sterile if package is intact
 20 Applicators
 NDC 54365-400-30
 REF 930299

CHLORAPREP ONE-STEP				
chlorhexidine gluconate and isopropyl alcohol solution				
Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54365-400	
Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)		CHLORHEXIDINE GLUCONATE	20 mg in 1 mL	
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)		ISOPROPYL ALCOHOL	0.7 mL in 1 mL	
Inactive Ingredients				
Ingredient Name		Strength		
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC 54365			

1	NDC:54365-400-30	20 in 1 CARTON	05/24/2019	
1		1 in 1 POUCH		
1		1.5 mL in 1 APPLICATOR; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020832	05/24/2019	

Labeler - CareFusion 213, LLC (826496312)

Registrant - Becton, Dickinson and Company (832696038)

Establishment

Name	Address	ID/FEI	Business Operations
CareFusion 213, LLC		826496312	analysis(54365-400) , manufacture(54365-400) , label(54365-400) , pack(54365-400) , sterilize(54365-400)

Revised: 11/2025

CareFusion 213, LLC