

**CHLORAPREP ONE-STEP- chlorhexidine gluconate and isopropyl alcohol solution**  
**CareFusion 213, LLC**

-----  
**BD ChloroPrep™ Clear 3 mL Applicator with Sterile Solution**

**WARNING FLAMMABLE**

**Keep away from fire or flame.**

**To reduce risk of fire, PREP CAREFULLY:**

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (**minimum of 3 minutes on hairless skin; up to 1 hour in hair**)
- do not allow solution to pool
- remove wet materials from prep area

***Active ingredients***

Chlorhexidine gluconate 2% w/v

Isopropyl alcohol 70% v/v

***Purposes***

Antiseptic

Antiseptic

***Use***

for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

***Warnings***

**For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:**

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (**minimum of 3 minutes on hairless skin; up to 1 hour in hair**)
- do not allow solution to pool
- remove wet materials from prep area

## **Allergy alert:**

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing • shock • facial swelling • hives • rash

If an allergic reaction occurs, stop use and seek medical help right away.

## **Do not use**

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

## **When using this product**

keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

## **Stop use and ask a doctor if**

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

## **Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

## **Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately **4 in. x 5 in. (130 cm<sup>2</sup>)**
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for

approximately 30 seconds. Do not blot or wipe away.

- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.
- **if using an ignition source, allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair).** Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

### **Other Information**

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

### **Inactive Ingredient**

USP purified water

### **Questions?**

- [www.chloraprep.com](http://www.chloraprep.com)
- call **1-800-523-0502** (M-F 8 a.m.-5 p.m. CST)

### **BD Chloraprep™ Clear 3 mL Applicator with Sterile Solution Principal Display Panel-Package Label**

Principal Display Panel-Package Label

BD Chloraprep™ Clear

3 mL Applicator

2% w/v chlorhexidine gluconate (CHG)

and 70% v/v Isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

Sterile Solution

NDC 54365-400-32

REF 930400



# 3 mL Applicator

2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Sterile Solution  
Applicator is sterile if package is intact

REF 930400 NDC 54365-400-32

External Use Only  
Professional Use Only



STERILE EO



STERILE SOLUTION

Clear

## ⚠ WARNING FLAMMABLE



Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (**minimum of 3 minutes on hairless skin; up to 1 hour in hair**)
- do not allow solution to pool
- remove wet materials from prep area

### Drug Facts

#### Active ingredients

Chlorhexidine gluconate 2% w/v.....	Antiseptic
Isopropyl alcohol 70% v/v.....	Antiseptic

#### Purposes

**Use** for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

#### Warnings

**For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:**

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (**minimum of 3 minutes on hairless skin; up to 1 hour in hair**)
- do not allow solution to pool
- remove wet materials from prep area

**Allergy alert:** This product may cause a severe allergic reaction. Symptoms may include:  
• wheezing/difficulty breathing • shock • facial swelling • hives • rash  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use** • on patients allergic to chlorhexidine gluconate or any other ingredient in this product • for lumbar puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser

**When using this product** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children** If swallowed get medical help or contact a Poison

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions** • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • use in a well ventilated area • maximal treatment area for one applicator is approximately **4 in. x 5 in. (130 cm<sup>2</sup>)** • remove applicator from package; do not touch sponge • hold the applicator with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic. • wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin • completely wet the treatment area with antiseptic • **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away. • **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away. • **if using an ignition source, allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)**. Do not blot or wipe away. • discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

**Other information** • store between 15–30 °C (59–86 °F) • avoid freezing and excessive heat above 40 °C (104 °F)

**Inactive ingredient** USP purified water

**Questions?** • [www.chloraprep.com](http://www.chloraprep.com) • call **1-800-523-0502** (M-F 8 a.m.–5 p.m. CST)

CareFusion 213, LLC, El Paso, TX 79912, subsidiary of Becton, Dickinson and Co.

Artwork No. 6-930420 • A050619 • Made in the USA of US and imported parts • [www.bd.com](http://www.bd.com)

## **BD ChloraPrep™ Clear 3 mL Applicator with Sterile Solution Principal Display Panel-Package Carton Label**

Principal Display Panel-Carton

BD ChloraPrep™ Clear

3 mL Applicators

2% w/v chlorhexidine gluconate (CHG)

and 70% v/v Isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

Sterile Solution

Applicator is sterile if package is intact

25 Applicators

NDC 54365-400-32

REF 930400



## CHLORAPREP ONE-STEP

chlorhexidine gluconate and isopropyl alcohol solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:54365-400
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CHLORHEXIDINE GLUCONATE</b> (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	20 mg in 1 mL
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54365-400-32	25 in 1 CARTON	05/24/2019	
1		1 in 1 POUCH		
1		3 mL in 1 APPLICATOR; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020832	05/24/2019	

**Labeler** - CareFusion 213, LLC (826496312)

**Registrant** - Becton, Dickinson and Company (832696038)

**Establishment**

Name	Address	ID/FEI	Business Operations
CareFusion 213, LLC		826496312	analysis(54365-400) , label(54365-400) , manufacture(54365-400) , pack(54365-400) , sterilize(54365-400)

Revised: 10/2025

CareFusion 213, LLC