

CHLORAPREP ONE-STEP- chlorhexidine gluconate and isopropyl alcohol solution
CareFusion 213, LLC

BD ChloraPrep™ Clear 10.5 mL Applicator with Sterile Solution

WARNING. FLAMMABLE.

Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- **solution contains alcohol and gives off flammable vapors**
- **avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.**
- **do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)**
- **do not allow solution to pool**
- **remove wet materials from prep area**

Active ingredients

Chlorhexidine gluconate 2% w/v

Isopropyl alcohol 70% v/v

Purposes

Antiseptic

Antiseptic

Use

for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (**minimum of 3 minutes on hairless skin; up to 1 hour in hair**)
- do not allow solution to pool

- remove wet materials from prep area

Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing • shock • facial swelling • hives • rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product

keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse

with cold water right away and contact a doctor.

Stop use and ask a doctor if

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately **8.4 in. x 8.4 in. (457 cm²)**
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth

- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- allow the solution to completely dry (**minimum of 3 minutes on hairless skin; up to 1 hour in hair**). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

• www.chloraprep.com • **call 1-800-523-0502** (M-F 8 a.m.-5 p.m. CST)



BD Chloraprep™ Clear
ANTISEPTICALLY PREPARED PREPARE-TO-USE SOLUTION

Drug Facts	
Active ingredient(s): 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)	Preparation: Sterile Solution
How to use: See package insert for complete instructions. For use on intact skin only.	Indication: For use on intact skin only.
Warnings: Do not use on broken skin or mucous membranes. Avoid contact with eyes, nose, mouth, and ears. If contact occurs, flush thoroughly with water. Do not use if the cap is missing or the solution is discolored.	
Directions: Apply a small amount of solution to the skin. Rub thoroughly for 30 seconds. Allow to dry. Do not use on broken skin or mucous membranes.	
Other information: Contains 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA). For use on intact skin only.	
Keep out of reach of children. See back of package for complete instructions.	
© 2014 BD. All rights reserved. BD, the BD logo, and Chloraprep are trademarks of BD.	



BD Chloraprep™ Clear
10.5 mL Applicators

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation Sterile Solution

Applicator is sterile if package is intact

Store between 15-30°C (59-86°F)
Avoid freezing and excessive heat above 40°C (104°F)

© 2014 BD, Inc. (715-642) 0003
25 Applicators



BD Chloraprep™ Clear
10.5 mL Applicators

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation Sterile Solution

Applicator is sterile if package is intact

Store between 15-30°C (59-86°F)
Avoid freezing and excessive heat above 40°C (104°F)

© 2014 BD, Inc. (715-642) 0003
25 Applicators



BD Chloraprep™ Clear
10.5 mL Applicators

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation Sterile Solution

Applicator is sterile if package is intact

Store between 15-30°C (59-86°F)
Avoid freezing and excessive heat above 40°C (104°F)

© 2014 BD, Inc. (715-642) 0003
25 Applicators

Clear

NDC 54365-400-34

REF 930700

WARNING

FLAMMABLE

Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (**minimum of 3 minutes on hairless skin; up to 1 hour in hair**)
- do not allow solution to pool
- remove wet materials from prep area

CHLORAPREP ONE-STEP

chlorhexidine gluconate and isopropyl alcohol solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54365-400
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	20 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54365-400-34	25 in 1 CARTON	05/24/2019	
1		1 in 1 POUCH		
1		10.5 mL in 1 APPLICATOR; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020832	05/24/2019	

Labeler - CareFusion 213, LLC (826496312)

Registrant - Becton, Dickinson and Company (832696038)

Establishment

Name	Address	ID/FEI	Business Operations
CareFusion 213, LLC		826496312	analysis(54365-400) , label(54365-400) , manufacture(54365-400) , pack(54365-400) , sterilize(54365-400)

Revised: 7/2025

CareFusion 213, LLC